

Synthesis, Evaluation of Anticancer Activity and QSAR Study of Heterocyclic Esters of Caffeic Acid

Shima Hajmohamad Ebrahim Ketabforoosh^a, Mohsen Amini^b, Mohsen Vosooghi^a,
Abbas Shafiee^a, Ebrahim Azizi^c and Farzad Kobarfard^{d*}

^aDepartment of Medicinal Chemistry, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran 14176, Iran. ^bDepartment of Medicinal Chemistry, Faculty of Pharmacy and Drug Design and Development Research Center, Tehran University of Medical Sciences, Tehran 14176, Iran. ^cMolecular Research Laboratory, Department of Pharmacology and Toxicology, Faculty of Pharmacy, Medical Sciences, University of Tehran, Tehran, Iran. ^dDepartment of Medicinal Chemistry, School of Pharmacy, and Phytochemistry Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Abstract

Caffeic acid phenethyl ester (CAPE) suppresses the growth of transformed cells such as human breast cancer cells, hepatocarcinoma, myeloid leukemia, colorectal cancer cells, fibrosarcoma, glioma and melanoma. A group of heterocyclic esters of caffeic acid was synthesized using Mitsunobu reaction and the esters were subjected to further structural modification by electrooxidation of the catechol ring of caffeic acid esters in the presence of sodium benzenesulfinate and sodium toluenesulfinate as nucleophiles. Both heterocyclic esters of caffeic acid and their arylsulfonyl derivatives were evaluated for their cytotoxic activity against HeLa, SK-OV-3, and HT-29 cancer cell lines.

HeLa cells showed the highest sensitivity to the compounds and heterocyclic esters with no substituent on catechol ring showed better activity compared to their substituted counterparts. QSAR studies reemphasized the importance of molecular shape of the compounds for their cytotoxic activity.

Keywords: Caffeic acid; Mitsunobu; Electrooxidation; Cytotoxic activity.

Introduction

Cancer is one of the most important causes of mortality worldwide. Although there are many treatment options for this disease such as radiotherapy, surgical operation, immunotherapy and chemotherapy, cancer still remains a serious clinical problem (1). Accordingly we need new approach for cancer therapy. Among the wide spectrum of cancer treatment strategies, chemotherapy has significant role in management of cancer. Therefore discovery of

novel anti cancer agents is one of the necessities in cancer research. The main objectives of anti cancer drug development are decreasing the toxicity and maximizing the efficacy.

Caffeic acid (CA) and caffeic acid phenethyl ester (CAPE) (Figure 1) have been identified as biologically significant components of honey bee propolis (2). Existing reports indicate that CAPE has various biological properties including antibacterial, antiviral, anti-inflammatory, antioxidant, anti-HIV, anti thrombosis, and tumor cell arrest (3-5). CAPE suppresses the growth of transformed cells such as human breast cancer cells, hepatocarcinoma, myeloid leukemia, colorectal cancer cells, fibrosarcoma,

* Corresponding author:

E-mail: kobarfard@sbm.ac.ir




Figure 1. Structure of Caffeic acid phenethyl ester.

glioma and melanoma (6-11). As a result CAPE is an appropriate lead compound for anti cancer studies.

Structure-activity relationship studies for caffeic acid derivatives suggest that catechol ring is essential for cytotoxicity of CAPE and it has been proven that part of its anticancer effect is due to antioxidant mechanism of catechol ring. Moreover methylation of hydroxyl groups reduces the activity of CAPE. Conjugated double bond is also required for inhibition of cell growth (12).

Aliphatic esters of caffeic acid with 2, 5- or 3, 5- or 2, 3- dihydroxy benzene rings had comparable cytotoxicity to doxorubicin. Replacement of the ester by amide or hydroxy groups substantially decreased or abolished cytotoxicity (13). QSAR studies on aliphatic esters of CA showed the dependence of cytotoxicity of this compound on hydrophobicity and steric factors (14). Based on the previous studies on CAPE analogues, it has been discovered that replacing phenyl with cyclohexyl increases the cytotoxic activity. Elongation of alkyl chain of phenethyl up to eight carbons, increases strength of compound. Furthermore in aliphatic esters, elongation of the chain up to 17 carbons enhances potency (15).

Despite the interesting trends observed in SAR studies of CAPE, only few analogues of this compound have been synthesized and studied mostly because of the surprisingly difficult synthesis of these groups of compounds.

The first part of this study was aimed at the synthesis of heterocyclic esters of caffeic acid and evaluation of their cytotoxic activity in compared to that of caffeic acid phenethyl ester (CAPE).

Phenolic carboxylic acids or alcohols

are notorious for their unamenability to most of methods of esterification. The tough reaction conditions which is needed in Fischer esterification method (strong protic acid) along with the need to use excess alcohol makes this strategy of limited applicability for making caffeic acid esters.

Lewis acid-catalyzed acylations will not solve the problem either (16). Using acylhalides to make esters via acyl nucleophilic substitution requires protection of phenolic hydroxyl groups since carboxylic acids will not discriminate between aliphatic hydroxyl (alcoholic) and aromatic hydroxyl (phenolic) groups.

Stuwe *et al.* reported a method based on the reaction of cesium salt of phenolic carboxylic acid with proper alkyl halide to give the desired ester but unsatisfactory yield and the need for excess of halide are still major issues for this method (17).

The enzymatic and ultrasound-accelerated enzymatic synthesis of caffeic acid phenethyl ester from caffeic acid and phenethyl alcohol have been also investigated but these methods have not gone beyond simple alkyl esters and phenethyl esters (18-19).

Knoevenagel reaction has also been used as an indirect method for the synthesis of caffeic acid 3, 4-dihydroxyphenethyl ester (20). The major drawback for this method is unavailability of the atypical starting materials which are required to implement the method.

Perhaps the most successful method in phenolic ester synthesis is the report by Appendino *et al.* in which Mitsunobu reaction has been employed to make a few polyphenolic esters (21).

CAPE and its analogues have a diphenolic ring (catechol) in their structures with could be electrochemically converted to 1, 2-benzoquinone by a reversible two-electron oxidation (Figure 2).

The electrochemically generated *o*-benzoquinones are quite reactive intermediate which in a proper condition can be attacked by a variety of nucleophiles and undergo various reactions such as Micheal addition. Sodium benzenesulfinate and Sodium toluene-4-sulfinate are two strong nucleophiles which could add to quinone ring via a nucleophilic




Figure 2. Oxidation of catechol ring.

addition reaction. Electrochemical methods are considered as green methods for the synthesis of organic compounds and therefore the second part of this study was aimed at the preparation of a few novel CA analogues with a benzenesulfonyl or toluene-4-sulfonyl group attached to their catechol ring using electrochemical method (22-23).

Chemistry

Caffeic acid and alcohols b-f were condensed in equimolar ratio in presence of triphenylphosphine (TPP) and diisopropylazodicarboxylate (DIAD) in dry tetrahydrofuran (THF) as solvent at room temperature. (Figure 3)

Liquid-liquid extraction was used to extract the crude product and final purification was performed by silicagel column chromatography using chloroform-methanol or ethylacetate-petroleum ether as the eluent solvent. The synthesized compounds were characterized

using IR, ¹HNMR, ¹³C-NMR spectroscopies, ESI-Mass spectrometry and elemental analysis.

All the heterocyclic esters of caffeic acids (2-6) and CAPE (1) were subjected to electrochemical oxidation in the presence of sodium benzenesulfinate or sodium benzenesulfinate or sodium toluene-4-sulfinate as nucleophiles (Figure 4). Compounds (1-6) were thus electro-oxidized to their corresponding quinonic forms (1a-6a) on the surface of glassy carbon electrode. When the electro-oxidation process takes place in the presence of sodium benzenesulfinate or toluene-4-sulfinate, the sulfinate group can attack the quinonic ring in a Michael type addition reaction. The quinonic ring could then be reduced electrochemically using a cyclic voltammetric process to generate the reduced form of CA esters. The electrochemical behavior of CA esters (1-6) were studied in the absence and presence of the nucleophiles sodium benzenesulfinate and sodium toluene-4-




Figure 3. Synthesis of caffeic acid esters (2-6) using Mitsunobu reaction.




Figure 4. Electrooxidation of CA esters (1-6) and reaction with sodium benzenesulfinate or sodium toluene-4-sulfinate.

sulfinate (Figure 5).

Voltammograms of 2 in the absence and presence of sodium toluene-4-sulfinate (Figure 5) show anodic peaks at 359 mV and 449 mV and cathodic peaks at 239 mV and 169 mV, respectively.

In this condition, part (c) of Figure 5 is related to the electrochemical oxidation of sodium toluene-4-sulfinate. Anodic

peaks indicate the oxidation of 3a-e to their corresponding orthobenzoquinones which are reduced back to their initial catechols form by reversing the voltage. The ratio of the oxidation and reduction current amplitudes for all in the oxidation and reduction process were equal to unity. This phenomenon shows the stabilization of o-benzoquinone produced on the surface of electrode. Side reactions, such as dimerization or hydroxylation of (compounds 1-6) are too slow to be observed on the time scale of cyclic voltammograms.

When compounds sodium benzenesulfinate or sodium toluene-4-sulfinate were added to the solution containing (compounds 1-6), the anodic peaks shifted positively in all cases (part (b) of Figure 5) and the cathodic peak currents decreased. The positive shift of anodic potential could be related to the formation of a thin film on the surface of the electrode. The formation of toluene-4-sulfinate adducts (1b-6b) and benzenesulfinate adducts (1c-6c) was confirmed by thin layer chromatography (TLC) and electrospray ionization mass spectrometry (ESI-MS).




Figure 5. Comparative cyclic voltammograms of 0.25 mM compound 2 without (a), with (b) 0.25 mM sodium toluene-4-sulfinate and 0.25 mM sodium toluene-4-sulfinate (c) at a glassy-carbon electrode ($S = \pi \text{ mm}^2$) in aqueous solution ($\text{H}_2\text{O}:\text{AN}$, 80:20), containing acetate buffer ($\text{pH} = 7$, $c = 0.2 \text{ M}$). Scan rate: 50 mV/s.

Pharmacology

Cell Culture

Human ovarian adenocarcinoma cell line SK-OV-3 (ATCC no. HTB-77), human




Figure 6. Mechanism of Mitsunobu reaction.

cervix adenocarcinoma cell line HeLa (ATCC no.CCL-2), and human colon adenocarcinoma HT-29 (ATCC no. HTB-38) were obtained from American Type Culture Collection. Cells were grown on 75 cm² cell culture flasks with EMEM (Eagle's minimum essential medium), supplemented with 10% fetal bovine serum, and 1% penicillin/streptomycin solution (10,000 units of penicillin and 10 mg of streptomycin in 0.9% NaCl) in a humidified atmosphere of 5% CO₂, 95% air at 37 °C.

Cell proliferation assay

Cell proliferation assay was carried out using CellTiter 96 aqueous one solution cell proliferation assay kit (Promega, USA). Briefly, upon reaching about 75-80% confluency, 5000

cells/well were plated in 96-well microplate in 100 μL media. After seeding for 72 h, the cells were treated with 50 μM compound in triplicate. Doxorubicin (10 μM) was used as the positive control. At the end of the sample exposure period (72 h), 20 μL CellTiter 96 aqueous solution was added. The plate was returned to the incubator for 1 h in a humidified atmosphere at 37 °C. The absorbance of the formazan product was measured at 490 nm using microplate reader. The blank control was recorded by measuring the absorbance at 490 nm with wells containing medium mixed with CellTiter 96 aqueous solution but no cells. Results were expressed as the percentage of the control (without compound set at 100%). The results are shown as the percentage of the control DMSO that has no

Table 1. The yields and mp for Mitsunobu reaction of caffeic acid with alcohols 2a-e.

Alcohol	yield	mp°C
b	10%	138.5-140
c	25%	167-169
d	15%	70-72
e	18%	154-157
f	20%	200-203

compound (set at 0%). All the experiments were performed in triplicate.

Result and Discussion

Chemistry

Esters are traditionally produced by condensing carboxylic acids and alcohols in the presence of an acid catalyst. The esterification reaction is both slow and reversible. There are many ways to drive the equilibrium toward the ester product: a: addition of an excess of one of the reactants (usually the alcohol), b: removal of the ester or the water by distillation, c: removal of water by azeotropic distillation and d: removal of water by use of dehydrating agent or molecular sieve.

When carboxylic acids and/or alcohols have other functional groups which are sensitive to the esterification conditions, unconventional methods should be employed to circumvent the vulnerability of these functional groups. Phenolic hydroxyl groups are among the interfering functional groups with the ordinary process of esterification.

Caffeic acid is a polyphenolic α - β unsaturated carboxylic acid with more issues in esterification reaction than just being a simple phenol. Therefore special conditions should be used for its esterification. In the present study we compared the following esterification methods for caffeic acid:

I. Acid catalyzed esterification with H_2SO_4 .

II. Microwave assisted esterification in presence of a protic acid.

III. Ultrasound assisted esterification in presence of a protic acid.

IV. Esterification with the aid of thionyl

chloride.

V. Esterification with the aid of dicyclohexyl dicarbodiimide(DCC).

VI. Wittig reaction between 3, 4-dihydroxy benzaldehyde and phosphonium chloride salt of alcohol.

VII. Knoevenagel reaction.

VIII. Esterification after protection of phenolic hydroxyl groups by BBr_3 and tetrahydropyrane.

IX. Mitsunobu reaction.

None of the above mentioned (except Mitsunobu reaction) led to the satisfactory results either because the reaction did not give the desired product at all (I, II, III, IV, V, VIII) or due to the formation of a great number of byproducts which made the purification of the desired product impossible (VI, VII, IX).

Mitsunobu reaction on the other hand was successful in giving the desired products for the reaction of caffeic acid with the alcohols b-f.

Mitsunobu reaction is a method for converting molecules that bear poor leaving group into a derivative that can react via S_N2 displacement, even when the nucleophile is as weak as a carboxylate anion. The process involves reaction of diisopropylazodicarboxylate (DIAD) with triphenylphosphine to form h. This dipolar ion reacts with $RCOOH$, which is present in the initial reaction to give i. The alcohol is then added to this phosphonium salt, and the subsequent reaction generates alkoxyphosphonium salt j. The $RCOO^-$ group displaces triphenylphosphine oxide to give the desired product.

It is generally accepted that the acids with a $pK_a < 11$ will be successful in Mitsunobu reaction. In light of this fact, it is not surprising that the acidic phenolic hydroxyl groups of caffeic acid would not interfere with this reaction. The yields for the Mitsunobu reaction conducted for caffeic acid reaction with alcohols b-f are presented in Table 1.

The purification of esters 2-6-e was performed on silica gel. This is more convenient than the use of Sephadex LH-20 which has been used by Appendino et al. in their report (21).

In general due to the mild conditions and the possibility of purification polyphenolic esters on silica gel column, the Mitsunobu reaction is particularly suitable for the synthesis

Table 2. Cytotoxic activity (IC_{50} , μm) of compounds 1-6, 1b-6b and 1c-6c.







Compound	Structure	HeLa	SK-OV-3	HT-29
1		12.3	42.1	25.1
1b		21.7	53.8	51.5
1c		23.8	49.1	48.5
2		17.4	73.2	37.4
2b		47.1	171.5	84.1
2c		38.9	190.6	53.7

Table 2. (Continued)








3		143.7	121.7	187
3b		140.2	>200	240.5
3c		191.5	>200	324
4		82.7	121.1	127.1
4b		67.1	>200	200.5
4c		111.2	>200	182
5		37.2	54.1	43.2

Table 2. (Continued)






				74.1
5b		40.1	>200	
5c		53.1	>200	182
6		121	147	57.3
6b		137	>200	96.1
6c		141	>200	73.2

Table 3. Correlation coefficient (R^2) matrix for some of descriptors used for HeLa.

	ASP	GATS3e
ASP	1	0.234
GATS3e		1

Table 4. Correlation coefficient (R^2) matrix for some of descriptors used for Ht-29.

	GATS1e	GATS3v
GATS1e	1	0.248
GATS3v		1

of polyphenolic compound such as caffeic acid esters, a group of compound with interesting biological activities.

Cytotoxic activity

The results of cytotoxic activity were summarized as IC_{50} (μ molar) of compounds in Table 2.

In general all compounds had good inhibitory activity ($IC_{50}=10-200 \mu$ M) against HeLa and HT-29 but they did not show significant inhibitory activity ($IC_{50} > 200 \mu$ M) against SK-OV-3. All these esters except 6, 6b, and 6c showed the strongest activity toward colon HT-29 cell line. All the benzenesulfonyl (1b-6b) and toluene-4-sulfonyl (1b-6b) derivatives showed decreased activity. It could be speculated that the arylsulfonyl group causes an electron deficiency

and/or steric hindrance on catechol ring and therefore interfere with the normal oxidation/reduction of catechol ring. Comparing the IC_{50} values of structures 1-6 indicates that thiophen is more successful than pyridine.

2-(2-pyridyle) ethyl analogues of CAPE (5, 5b, 5c) have stronger cytotoxicity than 2-pyridyle methyl analogues (6, 6b, 6c). This is in agreement with the previous finding for CAPE which shows that phenethyl esters is more potent than phenmethyl ester of caffeic acid.

QSAR study

In order to evaluate the effects of the structural parameters of the investigated compounds 1-6, 1b-6b and 1c-6c on their cytotoxic activities, Quantitative-Structure Activity Relationship (QSAR) analysis with different types of

Table 5. Data of the selected descriptors used in this study and predicted values of pIC_{50} (HeLa).

Compound	ASP	GATS3e	Predicted pIC_{50} (HeLa)
1	0.600	1.802	4.056
1b	0.239	1.784	3.904
1c	0.238	1.788	3.904
2	0.628	1.971	4.090
2b	0.274	1.891	3.932
2c	0.308	1.902	3.947
3	0.662	1.817	4.084
3b	0.224	1.791	3.898
3c	0.279	1.805	3.923
4	0.659	1.799	4.080
4b	0.227	1.779	3.898
4c	0.282	1.793	3.923
5	0.559	1.996	4.064
5b	0.260	1.912	3.929
5c	0.258	1.922	3.929
6	0.657	1.823	4.082
6b	0.247	1.794	3.908
6c	0.297	1.808	3.931




Figure 7. Plots of cross-validated predicted values of activity by MLR against the experimental values for different cell lines: (A) HeLa and (B) HT-29.

molecular descriptors was performed.

Several physicochemical descriptors such as hydrophobicity, topological indices, electronic parameters and steric factors are usually used in QSAR studies in order to find the effects of different structural properties on the biological activity of compound of interest. Since the calculated values of some electronic descriptors depend on the three-dimensional molecular

geometry, the optimum 3-D geometry of the molecules were obtained by Hyperchem software (Hypercube Inc, USA), using AM1 semi-empirical method. The resulting structures were used to calculate constitutional, functional, geometrical and topological descriptors by Dragon software. Meanwhile some electronic descriptors such as frontier molecular orbital (HOMO, LUMO), dipole moment and partial charges were calculated by the

Table 6. Data of the selected descriptors used in this study and predicted values of pIC_{50} (HT-29)

Compound	GATS1e	GATS3v	Predicted pIC_{50} (Ht-29)
1	1.746	2.221	4.521
1b	1.381	2.244	4.376
1c	1.390	2.146	4.387
2	1.773	2.274	4.528
2b	1.400	2.283	4.380
2c	1.412	2.180	4.393
3	1.803	2.446	4.525
3b	1.427	2.394	4.382
3c	1.441	2.293	4.396
4	1.803	2.506	4.521
4b	1.427	2.431	4.379
4c	1.441	2.332	4.392
5	1.780	2.099	4.545
5b	1.416	2.169	4.396
5c	1.427	2.060	4.409
6	1.803	2.255	4.541
6b	1.427	2.276	4.391
6c	1.441	2.166	4.406

Hyperchem software. Activity data were converted to logarithmic scale (*i.e.* pIC_{50} for cytotoxicity in different cell lines). For each set of descriptors, the best multi-linear regression equations were obtained by the stepwise selection methods of multiple linear regression (MLR) subroutine of SPSS software. The correlation coefficient (r^2), standard error of regression (SE), correlation coefficient for cross-validation significance (q^2), root mean square error (RMS) and significant level (p-value) were employed to judge the validity of regression equation. As colinearity degrades the performances of the MLR-based QSAR equation, at the first, correlation analysis was performed to detect the co-linear descriptors (24). Thus, the correlation of descriptors with each other and with activity data was examined and among the co-linear descriptors one of them that represented the highest correlation with activity was retained and the rest were omitted. The resulted correlation matrix is represented in Table 3 and 4 for the remaining descriptors.

In the first step, it was tried to find an appropriate model for cytotoxic activity in HeLa cell line. The obtained equation is shown by Equation 1:

$$\text{pIC}_{50} = 0.416 \pm (0.021) \text{ASP} + 0.129 \pm (0.054) \text{GATS3e} + 3.574 \pm (0.098) \quad (\text{Equation 1})$$

$$n = 18 \quad r^2 = 0.968 \quad \text{RMScv} = 0.019 \quad \text{SE} = 0.015 \quad q^2 = 0.941 \quad F = 226$$

In this equation, the values in the parenthesis represent the standard deviation of the coefficients.

This model contains asphericity (ASP), Geary autocorrelation of lag 3 weighted by Sanderson electronegativity (GATS3e).

The next equation was obtained for cytotoxicity against Ht₂₉ cell line:

$$\text{pIC}_{50} = 0.393 \pm (0.016) \text{GATS1e} - 0.081 \pm (0.023) \text{GATS3v} + 4.015 \pm (0.052) \quad (\text{Equation 2})$$

$$n = 18 \quad r^2 = 0.976 \quad \text{RMscv} = 0.013 \quad \text{SE} = 0.011 \quad q^2 = 0.966 \quad F = 311$$

Equation 2 includes Geary autocorrelation of lag 1 weighted by Sanderson electronegativity

(GATS1e), Geary autocorrelation of lag 3 weighted by van der Waals volume (GATS3v). The values of the descriptors used by Equations 1 and 2 together with the predicted $-\log \text{IC}_{50}$ (pIC_{50}) are listed in Table 3, 4.

The correlation between the observed and predicted activities of all the compounds using Equation 1 and Equation 2 is represented graphically in Figure 7A and Figure 7B.

These two equations show that spatial and topological descriptors play important role in cytotoxicity of these compounds. The autocorrelation vectors represent the degree of similarity between molecules. Therefore, cytotoxic activity of these series of compounds is mainly dependant on molecular shape of structure.

Conclusion

A few heterocyclic esters of caffeic acid were synthesized using Mitsunobu reaction. Other methods of esterification failed to give the desired compounds possibly due to the sensitive nature of catechol ring in the structure of caffeic acid. The esters were further modified by attaching benzensulfonyl or toluene-4-sulfonyl group to their catechol ring using cyclic voltammetry method.

Evaluation of cytotoxic activity for the compounds indicates that the compounds are more active against HeLa cancer cell line compared to SK-OV-3 and HT-29.

QSAR analysis reveals that the cytotoxic activity is mainly under the influence of molecular shape of these compounds.

Incorporation of arylsulfonyl groups diminishes the activity, perhaps due to the electron withdrawing and/or steric effect.

Experimental

General method

Chemicals and all solvents used in this study were purchased from Merck AG and Aldrich Chemicals. CAPE (1) was prepared using Wittig reaction according to literature report (12). Melting points were determined on a Kofler hot stage apparatus. The IR spectra were obtained on a Shimadzu 470 spectrophotometer (potassium

bromide disks). ^1H NMR spectra were recorded on a Varian unity 500 spectrometer and chemical shifts (d) are reported in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard. Elemental analyses were carried out on a Costec rapid elemental analyzer Model 4010 (GmbH-Germany) for C, H, N and S, and the results are within $\pm 0.4\%$ of the theoretical values. Merck silica gel 60 F254 plates were used for analytical TLC; column chromatography was performed on Merck silica gel (70–230 mesh).

The electrospray mass spectra (ESI) were recorded on an Agilent 4610 triple quadrupole mass spectrometer.

Cyclic voltammetry was performed, using Metrom computerized voltammetric analyzer model 746 VA Trace Analyzer/747 VA Stand. Controlled-potential coulometry and preparative electrolysis were performed using BHP2050 potentiostat/galvanostat. The working electrode used in the voltammetry studies was a glassy carbon disc (1.8 mm diameter). The potential were measured versus the Ag/AgCl/KCl (3M) as a reference electrode and platinum wire was used as the counterelectrode.

In macroscale electrolysis, four carbon rods (8 mm diameter and 5 cm length) were used as working electrodes.

General procedure for the synthesis of 2-6

To a solution of alcohols b-f (2.5 mmole) and caffeic acid (2.5 mmole) in dry THF under N_2 atmosphere, were added TPP (2.3 mmole) and DIAD (2.3 mmole) at 0°C . The reaction mixture was stirred at room temperature for 2 days. Then, the reaction was worked up by removal of the solvent, redissolving the residue in ethyl acetate and washing with 1N NaHCO_3 (ca. 50 mL/250mg of alcohol). The product obtained after evaporation of ethylacetate was purified by silica gel column chromatography, eluting with ethyl acetate-petroleum ether (20:80 for 3-6) or chloroform-methanol (95:5 for 4).

(E)-2-(thiophen-2-yl)ethyl 3-(3,4-dihydroxyphenyl) acrylate (2)

IR, ν/cm^{-1} : 3465, 3224, 2924, 2854, 1671, 1603; ^1H NMR (DMSO- d_6), δ : 9.5 (bs, 2H, OHs),

7.48 (d, $J=16\text{Hz}$, 1H, olefinic H), 7.35 (d, $J=5\text{ Hz}$, 1H, H-5 of thiophene ring), 7.04 (s, 1H, H-2 of catechol ring), 6.99 (d, $J=8\text{ Hz}$, 1H, H-6 of catechol ring), 6.96 (m, 2H, H-3 and H-4 of thiophene ring), 6.75 (d, $J=8\text{Hz}$, 1H, H-5 of catechol ring), 6.25 (d, $J=16\text{ Hz}$, 1H, olefinic H), 4.30 (t, $J=6.5\text{ Hz}$, 2H, CH_2 -2), 3.17 (t, $J=6.5\text{ Hz}$, 2H, CH_2 -1); ^{13}C NMR (DMSO- d_6), δ : 166.4, 148.5, 145.5, 145.3, 140, 126.9, 125.7, 125.4, 124.4, 121.4, 115.7, 114.8, 113.7, 64, 21.9. MS, m/z ($I_r/\%$): 291.06 (M^+). Anal. Calcd. For $\text{C}_{15}\text{H}_{14}\text{O}_4\text{S}$: C, 62.05; H, 4.86. Found: C, 62.27; H, 4.72.

(E)-pyridin-4-ylmethyl 3-(3,4-dihydroxyphenyl) acrylate (3)

IR, ν/cm^{-1} : 3437, 3033, 2924, 1711, 1635, 1602; ^1H NMR (DMSO- d_6), δ : 9.66 (s, 1H, OH), 9.17 (s, 1H, OH), 8.56 (d, $J=6\text{Hz}$, 2H, H-2 and H-6 of pyridine ring), 7.57 (d, $J=15.5\text{Hz}$, 1H, olefinic H), 7.38 (d, $J=6\text{Hz}$, 2H, H-3 and H-5 of pyridine ring), 7.08 (d, $J=2\text{Hz}$, 1H, H-2 of catechol), 7.04 (dd, $J=8\text{Hz}$, 2H, 1H, H-6 of catechol), 6.76 (d, $J=8\text{Hz}$, 1H, H-5 of catechol ring), 6.37 (d, $J=15.5\text{Hz}$, olefinic H), 5.24 (s, 2H);

^{13}C NMR (DMSO- d_6), δ : 166.2, 149.7, 149.2, 148.6, 146, 145.5, 125.4, 121.8, 121.6, 121, 115.7, 114.9, 113.2, 63.5, 61.4. MS, m/z ($I_r/\%$): 272.08 (M^+). Anal. Calcd. For $\text{C}_{15}\text{H}_{13}\text{NO}_4$: C, 66.41; H, 4.83; N, 5.16. Found: C, 66.58; H, 5.02; N, 5.31.

(E)-pyridin-3-ylmethyl 3-(3,4-dihydroxyphenyl) acrylate (4)

IR, ν/cm^{-1} : 3441, 3060, 2928, 1713, 1632, 1602; ^1H NMR (DMSO- d_6), δ : 9.5 (bs, 2H, OHs), 8.63 (s, 1H, H-2 of pyridine ring), 8.54 (d, $J=4.5\text{Hz}$, 1H, H-6 of pyridine ring), 7.83 (d, $J=7.5\text{Hz}$, 1H, H-4 of pyridine ring), 7.53 (d, $J=15.5\text{Hz}$, 1H, olefinic H), 7.41 (dd, $J=7.5\text{Hz}$, 4.5Hz, 1H, H-5 of pyridine ring), 7.06 (d, $J=2\text{Hz}$, 1H, H-2 of catechol ring), 7.01 (dd, $J=8.5\text{Hz}$, 2H, 1H, H-6 of catechol ring), 6.75 (d, $J=8.5\text{Hz}$, 1H, H-5 of catechol ring), 6.32 (d, $J=15.5\text{Hz}$, 1H, olefinic H), 5.23 (s, 2H, CH_2); ^{13}C NMR (DMSO- d_6), δ : 166.3, 149.3, 148.5, 148.1, 145.8, 136, 132.1, 125.4, 123.6, 121.5, 115.7, 114.9, 113.4, 63, 60.6. MS, m/z ($I_r/\%$): 272.08 (M^+). Anal. Calcd. For $\text{C}_{15}\text{H}_{13}\text{NO}_4$: C, 66.41; H, 4.83; N, 5.16. Found: C, 66.79; H,

4.62; N, 5.44.

(E)-2-(pyridin-2-yl) ethyl 3-(3,4-dihydroxyphenyl) acrylate (5)

IR, ν/cm^{-1} : 3449, 3214, 2897, 1689, 1637, 1599; ^1H NMR (DMSO- d_6), δ : 9.4(bs, 2H, OHs), 8.5 (d, $J=5$ Hz, 1H, H-6 of pyridine ring), 7.71 (t, $J=7.5$ Hz, 1H, H-4 of pyridine ring), 7.42 (d, $J=16$ Hz, 1H, olefinic H), 7.32 (d, $J=7.5$ Hz, 1H, H-3 of pyridine ring), 7.23 (t, $J=5$ Hz, 1H, H-5 of pyridine ring), 7.01 (s, 1H, H-2 of catechol ring), 6.97 (d, $J=8$ Hz, 1H, H-5 of catechol ring), 6.74 (dd, $J=8$ Hz, 2 Hz, 1H, H-6 of catechol ring), 6.2 (dd, $J=16$ Hz, 2 Hz, 1H, olefinic H), 4.47 (t, $J=6.5$ Hz, 2H, CH_2-2), 3.09 (t, $J=6.5$ Hz, 2H, CH_2-1); ^{13}C NMR (DMSO- d_6), δ : 166.5, 157.9, 149.1, 148.4, 145.5, 145.2, 136.5, 125.4, 123.4, 121.7, 121.4, 115.7, 114.8, 113.8, 62.9, 36.6. MS, m/z ($I_r\%$): 286.10 (M^+). Anal. Calcd. For $\text{C}_{16}\text{H}_{15}\text{NO}_4$: C, 67.36; H, 5.30; N, 4.91. Found: C, 67.01; H, 5.42; N, 4.73.

(E)-pyridin-2-ylmethyl 3-(3,4-dihydroxyphenyl) acrylate (6)

IR, ν/cm^{-1} : 3476, 3056, 2953, 2563, 1693, 1600; ^1H NMR (DMSO- d_6), δ : 9.63 (s, 1H, OH), 8.55 (d, $J=4$ Hz, 1H, H-6 of pyridine ring), 7.82 (t, $J=8$ Hz, 1H, H-4 of pyridine ring), 7.55 (d, $J=16$ Hz, 1H, olefinic H), 7.43 (d, $J=7.5$ Hz, 1H, H-3 of pyridine ring), 7.33 (dd, $J=8$ Hz, 4 Hz, 1H, H-5 of pyridine ring), 7.07 (s, 1H, H-2 of catechol ring), 7.03 (d, $J=8$ Hz, 1H, H-6 of catechol ring), 6.76 (d, $J=8$ Hz, 1H, H-5 of catechol ring), 6.37 (d, $J=16$ Hz, 1H, olefinic H), 5.25 (s, 2H, CH_2); ^{13}C NMR (DMSO- d_6), δ : 166.2, 155.9, 149.1, 148.5, 145.8, 145.6, 136.9, 125.4, 122.9, 121.5, 115.7, 114.9, 113.5, 65.9, 63.9. MS, m/z ($I_r\%$): 272.08 (M^+). Anal. Calcd. For $\text{C}_{15}\text{H}_{13}\text{NO}_4$: C, 66.41; H, 4.83; N, 5.16. Found: C, 66.64; H, 4.52; N, 5.31.

General procedure for electroorganic synthesis of 1b-6b and 1c-6c

The mixture of water-acetonitrile (AN) (80:20), containing phosphate buffer (pH= 7.0, C=0.2M), was preelectrolyzed at 0.3V.

Equimolar amounts of caffeic acid eters and sodium benzenesulfinate or (sodium toluene-4-sulfinate) were added to the cell with 4 graphit

rodes as working electrode and Pt electrode as counter electrode.

The potentials of working electrode were measured versus the Ag/AgCl/KCl as a reference electrode. The electrolysis was interrupted many times, when the current reached to 5% of the starting value, to wash the anodic electrode with acetone to reactivate it. The precipitated products were filtered off and washed with water/acetone mixture.

Compounds 1b-6b and 1c-6c

Compounds were characterized by ESI mass spectrometry and all of the compounds were confirmed by observing their pseudomolecular ion as hydrogen adduct.

Acknowledgment

Authors would like to thank the research council of Medical Sciences/University of Tehran for facilitating this research project. We are also thankful to Dr. A. Bayandori Moghaddam from department of Engineering Science, College of Engineering, University of Tehran for his scientific advice on electrochemical reactions.

References

- (1) El-Refaei MF and El-Naab MM. Inhibitory effect of caffeic acid phenethyl ester on mice bearing tumor involving angiostatic and apoptotic activities. *Chemico-Biological Interactions* (2010) 186: 152-156.
- (2) Grunberger D, Banerjee R, Eisinger K, Oltz EM, Efros L, Caldwell M, Estevez V and Nakanishi K. Preferential cytotoxicity on tumor cells by caffeic acid phenethyl ester isolated from propolis. *Experientia* (1988) 44: 230-232.
- (3) Orsolio N, Knezevic A H, Sver L, Terzic S and Basic I. Immunomodulatory and antimetastatic action of propolis and related polyphenolic compounds. *J. Ethnopharmacol.* (2004) 94: 307-315.
- (4) Icklaus MC, Neamati N, Hong H, Mazumder A, Sunder S, Chen J, Milne GWA and Pommier Y. HIV-1 integrase pharmacophore: discovery of inhibitors through three-dimensional database searching. *J. Med. Chem.* (1997) 40: 920-929.
- (5) Zaemzadeh N, Hemmati A, Arzi A, Jalali M, Rashidi I. *Iran. J. Pharm. Res.* (2011) 10: 321-328.
- (6) -Wang D, Xiang D, He Y, Li Z, Wu X, Mou J, Xiao H and Zhang Q. Effect of caffeic acid phenethyl ester on proliferation and apoptosis of colorectal cancer cells in vitro. *World J Gastroenterol* (2005) 11: 4008-4012.

- (7) Chung T, Moon S, Chang Y, Ko J, Lee Y, Cho G, Kim S, Kim J, Kim A. Novel and therapeutic effect of caffeic acid and caffeic acid phenyl ester on hepatocarcinoma cells: complete regression of hepatoma growth and metastasis by dual mechanism. *FASEB J.* (2004) 18: 1670–1681.
- (8) Lee Y, Kuob H, Chuc C, Wangb C, Lin W and Tseng T. Involvement of tumor suppressor protein p53 and p38 MAPK in caffeic acid phenethyl ester-induced apoptosis of C6 glioma cells. *Biochem. Pharmacol.* (2003) 66: 2281-2289.
- (9) Hye J, Park H, Chung H, Min H, Park E, Hong J and Lee S. Inhibitory effects of caffeic acid phenethyl ester on cancer cell metastasis mediated by the down-regulation of matrix metalloproteinase expression in human HT1080 fibrosarcoma cells. *J. Nutr. Biochem.* (2006) 17: 356-362.
- (10) Kuo H, Kuo W, Lee Y, Wang C and Tseng T. Enhancement of caffeic acid phenethyl ester on all-trans retinoic acid-induced differentiation in human leukemia HL-60 cells. *Toxicol. Appl. Pharmacol.* (2006) 216: 80-88.
- (11) Kampa M, Alexaki VI, Notas G, Nifli AP, Nistikaki A, Hatzoglou A, Bakogeorgou E, Kouimtoglou E, Blekas G, Boskou D, Gravanis A and Castanas E. Antiproliferative and apoptotic effects of selective phenolic acids on T47D human breast cancer cells: potential mechanisms of action. *Breast Cancer Res.* (2004) 6: R63-R74.
- (12) Lee Y, Liao P, Chen W and Yang. Preferential cytotoxicity of caffeic acid phenethyl ester analogues on oral cancer cells. *Cancer Lett.* (2000) 153: 51-56.
- (13) Nam N, You Y, Kim Y, Hong D, Kim H and Z.Ahn B. Syntheses of certain 3-aryl-2-propenoates and evaluation of their cytotoxicity. *Bioorg. Med. Chem. Lett.* (2001) 11: 1173-1176.
- (14) Etzenhouser B, Hansch C, Kapur S and Selassie CD. Mechanism of toxicity of esters of caffeic and dihydrocaffeic acids. *Bioorg. Med. Chem.* (2001) 9: 199-209.
- (15) Nagaoka T, Banskota AH, Tezuka Y, Saiki I and Kadota S. Selective antiproliferative activity of caffeic acid phenethyl ester analogues on highly liver-metastatic murine colon 26-15 carcinoma cell line. *Bioorg. Med. Chem.* (2002) 10: 3351-3359.
- (16) Chandra KL, Saravanan P, Singh RK and Singh VK. Lewis acid catalyzed acylation reactions: scope and limitations. *Tetrahedron* (2002) 58: 1369-1374.
- (17) Stuwe HT, Bruhn G, Konig WA and Hausen BM. The synthesis of caffeic acid esters, a new group of naturally occurring contact allergens. *Naturwissenschaften* (1989) 76: 426-427.
- (18) Chen H, Chen J, Chang C and Shieh C. Optimization of ultrasound-accelerated synthesis of enzymatic caffeic acid phenethyl ester by response surface methodology. *Ultrasonic Sonochem.* (2011) 18: 455-459.
- (19) Widjaja A, Yeh T and Ju Y. Enzymatic synthesis of caffeic acid phenethyl ester. *J. Chin. Ins. Chem. Engin.* (2008) 39: 413-418.
- (20) Zhang Z, Xiao B, Chen Q and Lian X. Synthesis and biological evaluation of caffeic acid 3,4-dihydroxyphenethyl ester. *J. Nat. Prod.* (2010) 73: 252-257.
- (21) Appendino G, Minassi A, Daddario N, Bianchi F and Tron GC. Chemoselective esterification of phenolic acids and alcohols. *Org. Lett.* (2002) 4: 3839-3841.
- (22) Bayandori Moghaddam A, Ganjali MR, Norouzi P and Niasari M. A green method on the electro-organic synthesis of new caffeic acid derivatives: Electrochemical properties and LC–ESI–MS analysis of products. *J. Electroanal. Chem.* (2007) 601: 205-210.
- (23) Bayandori Moghaddam A, Ganjali MR, Dinarvand R, Norouzi P, Saboury AA and Moosavi-Movahedi AA. Electrochemical behavior of caffeic acid at single-walled carbon nanotube:graphite-based electrode. *Biophys. Chem.* (2007) 128: 30-37
- (24) Fernández M and Caballero J. QSAR modeling of matrix metalloproteinase inhibition by *N*-hydroxy-*a*-phenylsulfonylester derivatives. *Bioorg. Med. Chem.* (2007) 15: 6298-6310

This article is available online at <http://www.ijpr.ir>
