

CO₂ Ocean Bistability on Terrestrial ExoplanetsR. J. Graham¹ , Tim Lichtenberg¹ , and Raymond T. Pierrehumbert¹ ¹Atmospheric, Oceanic and Planetary Physics, Department of Physics, University of Oxford, Oxford, UK

Key Points:

- At low instillations within the circumstellar habitable zone, rocky planets with H₂O oceans may also develop oceans of condensed CO₂
- CO₂-ocean-bearing climate states are bistable with more traditional Earth-like climates where only water condenses at the planetary surface
- CO₂ oceans may occur even in the presence of the negative feedback on planetary climate provided by silicate weathering

Correspondence to:

R. J. Graham,
robert.graham@physics.ox.ac.uk

Citation:

Graham, R. J., Lichtenberg, T., & Pierrehumbert, R. T. (2022). CO₂ ocean bistability on terrestrial exoplanets. *Journal of Geophysical Research: Planets*, 127, e2022JE007456. <https://doi.org/10.1029/2022JE007456>

Received 13 JUL 2022

Accepted 29 SEP 2022

Abstract Cycling of carbon dioxide between the atmosphere and interior of rocky planets can stabilize global climate and enable planetary surface temperatures above freezing over geologic time. However, variations in global carbon budget and unstable feedback cycles between planetary sub-systems may destabilize the climate of rocky exoplanets toward regimes unknown in the Solar System. Here, we perform clear-sky atmospheric radiative transfer and surface weathering simulations to probe the stability of climate equilibria for rocky, ocean-bearing exoplanets at instillations relevant for planetary systems in the outer regions of the circumstellar habitable zone. Our simulations suggest that planets orbiting G- and F-type stars (but not M-type stars) may display bistability between an Earth-like climate state with efficient carbon sequestration and an alternative stable climate equilibrium where CO₂ condenses at the surface and forms a blanket of either clathrate hydrate or liquid CO₂. At increasing instillation and with ineffective weathering, the latter state oscillates between cool, surface CO₂-condensing and hot, non-condensing climates. CO₂ bistable climates may emerge early in planetary history and remain stable for billions of years. The carbon dioxide-condensing climates follow an opposite trend in *p*CO₂ versus instillation compared to the weathering-stabilized planet population, suggesting the possibility of observational discrimination between these distinct climate categories.

Plain Language Summary On Earth, water is the only molecule that occurs in large quantities as both vapor and condensed phases, but that may not be the case for every planet. We simulate terrestrial exoplanets that receive less light from their stars than the Earth does. We find that CO₂ may build up to such high levels that it condenses out onto the planetary surface, allowing for oceans of liquid CO₂ and/or sheets of CO₂ clathrate to accumulate. Depending on factors like CO₂ outgassing rate, the kinds of silicate rocks present at the planetary surface, and the level of irradiation received by a given planet from its parent star, such oceans can remain stable for geologic-scale time periods, or a planet may oscillate back and forth between an Earth-like, non-CO₂-condensing state and an exotic CO₂-condensing state. Planets with liquid (or solid) CO₂ at their surface can have a profoundly different evolution than Earth, with important implications for their potential to host life.

1. Introduction

Earth's surface is dominated by a liquid water ocean in direct contact with the lithosphere. This state of affairs seems to be crucial for Earth's long-term climate stability and habitability, with the carbonate-silicate cycle modulating and stabilizing the planet's atmospheric CO₂ inventory through a set of water-rock chemical reactions taking place on the continents and seafloor (Berner et al., 1983; Coogan & Gillis, 2013; Penman et al., 2020; Walker et al., 1981). The carbonate-silicate cycle acts as a thermostat when CO₂ is a net greenhouse gas; however, when CO₂ builds to high enough levels, it increases planetary Rayleigh scattering and behaves as a coolant, which can convert the carbonate-silicate cycle into a destabilizing positive feedback. This suggests that otherwise Earth-like planets with large enough carbon inventories might be able to support climate configurations with high enough *p*CO₂ for either a hot, supercritical, Venus-like atmosphere at high instillations or an exotic, subcritical atmosphere with surface liquid CO₂ condensation coexisting with a liquid water ocean at low instillations. Planets of the latter variety might be difficult to remotely distinguish from more traditionally "Earth-like" planets lacking surface CO₂ condensation at equivalent orbits, but the geochemistry and potential habitability of these worlds would be radically different, even with a temperate surface climate. Most previous examinations of surface CO₂ condensation on terrestrial (exo)planets have focused on cold, glaciated climates where CO₂ would only condense as a solid (Bonati & Ramirez, 2021; Kadoya & Tajika, 2019; Turbet et al., 2017); waterworlds with high pressure ice mantles (Marounina & Rogers, 2020; Ramirez & Levi, 2018); or the potential for CO₂ condensation on Mars in the deep past (Forget et al., 2013; Kasting, 1991; Soto et al., 2015). In this study, we focus on surface CO₂ condensation on rocky exoplanets with temperate climates in different end-member weathering regimes that

© 2022 The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial License](https://creativecommons.org/licenses/by-nc/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

inform the anticipated diversity of potentially habitable planets (Kasting et al., 1993; Wordsworth, Forget, Selsis, et al., 2010; von Paris et al., 2013).

From an astronomical perspective, carbon compounds are strongly depleted on the terrestrial planets of the Solar System relative to the nominal values in the Sun or the interstellar medium (Öberg & Bergin, 2021) as a result of processes operating in the protoplanetary disk (Krijt et al., 2020; Li et al., 2021) and on planetesimals (Hirschmann et al., 2021; Lichtenberg & Krijt, 2021). In addition, volatile partitioning into metal and melt phases can redistribute major carbon and hydrogen carriers between core, mantle, and atmospheric reservoirs and partly decouple the initially accreted volatile reservoir (Fischer et al., 2020; Grewal et al., 2020) from the atmospheric composition of rocky exoplanets. A recent example is provided by the outer TRAPPIST-1 planets, for which mass-radius constraints suggest volatile mass fractions on the order of several weight per cent (Agol et al., 2021). On a statistical level, the larger sub-Neptune cluster of the Kepler radius valley suggests that at least a fraction of systems accrete substantial volatile mass budgets during their formation (Venturini et al., 2020; Zeng et al., 2019). The anticipated variation in carbon abundance suggests that the majority of rocky exoplanets may exhibit diverse climate regimes, for which the thermodynamic limits to maintain clement surface states are poorly understood (Lichtenberg et al., 2022). Future exoplanet surveys that will aim to probe the atmospheres of temperate exoplanets to test the range of climate diversity (Checlair et al., 2021; Gaudi et al., 2020; Quanz, Absil, et al., 2021; Quanz, Ottiger, et al., 2021) will rely on physically motivated theories to interpret their findings.

Here, we study the interplay between silicate weathering and CO₂ pressure variations to probe the limits of clement climates on terrestrial exoplanets. We use 1-D, two-stream radiative transfer and carbon cycle simulations to investigate the behavior of climates with high partial pressure of CO₂ ($p\text{CO}_2$) and low irradiation from the central star (S_{eff}) without (Sections 3.1 and 3.2) and with (Section 3.3) weathering feedbacks. Our simulations suggest that terrestrial planets at low instillations in the classical circumstellar habitable zone (HZ; Kasting et al., 1993) may emerge from their accretionary period directly into stable climate states with long-lasting periods of liquid CO₂ surface condensation, even in cases where they are not initially globally glaciated.

2. Methods

In this study we combine global-mean, clear-sky climate and silicate weathering calculations to examine the interplay between radiative and geochemical feedbacks on ocean-bearing, high- $p\text{CO}_2$ planets in the outer reaches of the classical HZ. Here we briefly outline the procedure we follow and the models we use for the radiative calculations and the weathering calculations.

2.1. Radiative Transfer

We carry out radiative transfer calculations using the SOCRATES code (Edwards & Slingo, 1996), solving the plane-parallel, two-stream approximated radiative transfer equation with scattering (see the extensive description in Lichtenberg et al., 2021, though note that the implementation in that paper does not include scattering). Opacity coefficients are tabulated and derived from the HITRAN database, making use of the line-by-line and collision-induced absorption coefficients for H₂O (Gordon et al., 2017), CO₂ (Gordon et al., 2017; Gruszka & Borysow, 1997), N₂ (Gordon et al., 2017; Karman et al., 2015), and the H₂O continuum (Mlawer et al., 2012). We note that the CO₂ continuum spectrum is uncertain at high temperatures and pressures, which introduces a potentially significant source of error into our calculations (e.g., Halevy et al., 2009; Wordsworth, Forget, & Eymet, 2010).

Rayleigh scattering cross-sections for CO₂ and N₂ are calculated following Vardavas and Carver (1984).

$$\sigma_{R,i} = \frac{0.2756}{\mu_i} \times \frac{6 + 3\Delta}{\lambda^4(6 - 7\Delta)} \left[A \left(1 + \frac{B}{\lambda^2} \right) \right]^2, \quad (1)$$

where the subscript i iterates over the species present, $\sigma_{R,i}$ (m² kg⁻¹) is the Rayleigh scattering cross-section, μ_i is the molar mass of species i (kg mol⁻¹), λ (μm) is wavelength, coefficients A and B are taken from Cox (2015), Δ is the depolarization factor, and the numerical values we use are given in Table 1. For H₂O, as far as we are aware, values for the coefficients A and B have not been published at the relevant wavelengths. For this reason,

Table 1
Tabulation of Rayleigh Scattering Data Used in This Study

Molecule	A (units of 10^{-6})	B (units of 10^{-3})	Δ	σ_0 ($10^{-7} \text{ m}^2 \text{ kg}^{-1}$)
CO ₂	4.39	6.4	0.0805	–
N ₂	2.906	7.7	0.0305	–
H ₂ O	–	–	–	9.32

Note. A and B values come from Cox (2015). σ_0 for H₂O is drawn from Pierrehumbert (2010). Δ values come from Vardavas and Carver (1984).

and because H₂O is a minor constituent in the atmospheres we simulate, we use a simple λ^{-4} scaling to calculate H₂O's Rayleigh scattering cross-section,

$$\sigma_{R,H_2O} = \sigma_{R,0} \frac{\lambda_0^4}{\lambda^4}, \quad (2)$$

where $\sigma_{R,0} = 9.32 \times 10^{-7} \text{ m}^2 \text{ kg}^{-1}$ (Pierrehumbert, 2010) and $\lambda_0 = 1 \text{ }\mu\text{m}$. Here we take the opportunity to note that some previous papers (Kopparapu et al., 2013; Pluriel et al., 2019), have erroneously used H₂O Rayleigh scattering coefficients calculated using a depolarization ratio that was drawn from a study (Marshall & Smith, 1990) of the scattering properties of *liquid* H₂O, not water vapor. The total Rayleigh scattering cross-section is calculated by summing the cross-sections of the individual species, weighted by volumetric mixing ratio,

$$\sigma_{R,tot} = \sum_i x_i \sigma_{R,i}, \quad (3)$$

where x_i represents the volumetric mixing ratio of a given species ($\sum_i x_i = 1$). SOCRATES does not allow vertically-varying Rayleigh scattering coefficients, so we take the mixing ratios at the surface to calculate the total scattering cross-sections.

The primary stellar spectrum we use in the presented calculations is based on measurements of the Sun's spectral irradiance (Kurucz, 1995), and thus represents irradiation from a G2V star. To show how the climate behaviors we identify depend on stellar type and age, we also present sets of simulations using spectra from AD Leonis, an M3.5V star (Segura et al., 2005), and Sigma Boötis, an F2V star (Segura et al., 2003). The different spectra result in different planetary albedo values for a given atmospheric composition and climate. We also tested the effects of a change in the solar spectrum with time: at 4.5 and 3.8 Ga before present (Claire et al., 2012) our simulations produced results that differed negligibly from the fiducial, modern case.

We apply a generalized pseudoadiabat formula (Graham et al., 2021) to calculate atmospheric temperature-pressure profiles with a variety of compositions and surface boundary conditions. This pseudoadiabat formula incorporates the fraction of retained condensate as a freely tunable parameter which can significantly impact the specific heat capacity and lapse rate in atmospheres with non-dilute condensable species. In our calculations, we assume that all condensate is instantaneously rained out upon condensation. We also assume H₂O saturation in the main set of simulations discussed in this paper. This formula is useful because it allows us to self-consistently calculate atmospheric profiles with any combination of condensable (e.g., H₂O, CO₂) and non-condensable (e.g., N₂) gases, though the atmospheres we focus on in this paper are simply CO₂ + H₂O (sensitivity tests with up to 10 bar of N₂ produced qualitatively identical results, as expected). Throughout this paper, unless otherwise noted, atmospheres are taken to have isothermal stratospheres with $T_{\text{strat}} = 150 \text{ K}$. This is comparable to the stratospheric temperatures hypothesized for high-CO₂ planets in the outer reaches of the classical circumstellar HZ (Kopparapu et al., 2013), and time-stepped radiative-convective calculations in high- $p\text{CO}_2$ atmospheres have recovered a stratospheric temperature of 150 K (Wordsworth & Pierrehumbert, 2013). Condensation of H₂O and CO₂ is assumed to cease within the stratosphere, so the mixing ratios remain constant at pressure levels below the highest pressure level (lowest altitude level) with a temperature of 150 K. Assuming a constant stratospheric temperature considerably simplifies our climate calculations (described further below), at the expense of neglecting the feedback between instellation and stratospheric temperature, which can change a planet's outgoing longwave radiation (OLR) and thus its surface temperature. Sensitivity tests carried out with an increased stratospheric temperature of up to 200 K demonstrate that the effect on climate is minor, with no qualitative changes to our results.

For rapid simulation of a wide range of surface conditions, we take an “inverse climate modeling” approach (Kasting, 1991). This entails choosing a $p\text{CO}_2$ and a surface temperature (which in turn specifies $p\text{H}_2\text{O}$ by the Clausius-Clapeyron relation), using those values as boundary conditions to integrate the pseudoadiabat from the surface up to the 150 K isothermal stratosphere, and running the radiative transfer code to get the OLR and albedo for that specific atmospheric temperature/pressure/composition combination. With those values, we can calculate the instellation necessary to maintain global-mean energetic balance between incoming and outgoing radiation,




Figure 1. Energetic properties of terrestrial planetary climates at low instellation as a function of surface temperature T_{surf} (y-axes) and CO_2 partial pressure (x-axes). In each panel, the black line represents the CO_2 saturation vapor pressure curve; climates with CO_2 greater than that of the vapor pressure curve at a given temperature are super-saturated at their surface (gray region in all three panels). The white dotted line in each panel represents the approximate contour location of the S_{eff} value separating climates that can support CO_2 bistability versus those that may limit cycle between condensing and non-condensing states (see Sections 3.2 and 3.3 for discussion of those climate types). The black dotted line in each panel marks the temperature below which CO_2 clathrate hydrates may be stable. In panel A, the contours represent the top-of-atmosphere stellar irradiation required to maintain a climate with a given surface temperature and $p\text{CO}_2$, normalized by the instellation received at Earth's orbit, that is, $S/S_{\text{Earth}} = S_{\text{eff}}$. Panel (b) shows the global-mean outgoing longwave radiation. Panel (c) shows the global-mean planetary albedo.

$$(1 - \alpha(T, p\text{CO}_2)) \frac{S}{4} = F_{\text{out}}(T, p\text{CO}_2), \quad (4)$$

$$S_{\text{eff}}(T, p\text{CO}_2) = \frac{S}{S_{\text{Earth}}} = \frac{4F_{\text{out}}(T, p\text{CO}_2)}{S_{\text{Earth}}(1 - \alpha(T, p\text{CO}_2))}, \quad (5)$$

where S is top-of-atmosphere instellation, $S_{\text{Earth}} = 1368 \text{ W m}^{-2}$ is Earth's present-day instellation, S_{eff} is the fraction of present-day Earth's instellation (e.g., $S_{\text{eff}} = 0.3$ is equivalent to 30% of present-day Earth instellation), F_{out} is the OLR, and α is the global-mean planetary albedo, with the cosine of the stellar zenith angle assumed to be the instellation-weighted global mean of $\frac{2}{3}$ in all calculations (Cronin, 2014). We also set the surface albedo to 0.0, similar to the albedo of a cloudless sea surface, which would be 2%–4% at the chosen stellar zenith angle (Li et al., 2006). Low-lying marine stratocumulus clouds that would increase the near-surface albedo to levels above that of the sea surface are expected to dissipate at CO_2 levels far lower than 1 bar, the lowest $p\text{CO}_2$ evaluated in this study, due to the inhibition of cloud-top radiative cooling and subsequent shutdown of cloud-sustaining lower-tropospheric convection (e.g., Schneider et al., 2019), though of course other processes could still cause low-lying clouds or hazes that would affect near-surface albedo on these planets.

We carried out a grid of these inverse climate calculations for $p\text{CO}_2$ levels ranging from 1 to 73 bar with increments of 1 bar and for T_{surf} from 250 to 365 K with increments of 5 K. Figure 1 shows equilibrium S_{eff} (panel a), OLR (panel b), and albedo (panel c), all as functions of T_{surf} and $p\text{CO}_2$. Linear interpolation of OLR and albedo to temperatures and $p\text{CO}_2$ levels between the climate grid points allows for fast climate calculations to examine a wide variety of scenarios.

Table 2
Model Parameters Used in This Study

Parameter	Units	Definition	Fiducial value
γ	–	Land fraction	0.3
a_g	–	Surface albedo	0.0
R_{planet}	meters (m)	Planetary radius	6.37×10^6
T_{ref}	Kelvin (K)	Reference global-avg. temperature	288
$p\text{CO}_{2,\text{ref}}$	bar	Reference CO_2 partial pressure	280×10^{-6}
q_{ref}	m yr^{-1}	Modern global-avg. runoff	0.20 (Oki et al., 2001)
ϵ	1/K	Fractional change in precip. per K change in temp.	0.03
V_{ref}	mol yr^{-1}	Modern global CO_2 outgassing	7.5×10^{12} (Gerlach, 2011; Haqq-Misra et al., 2016)
ν	$\text{mol m}^{-2} \text{yr}^{-1}$	Modern CO_2 outgassing per m^2 planetary area	0.0147
Λ	Variable	Thermodynamic coefficient for C_{eq}	1.4×10^{-3}
n	–	Thermodynamic $p\text{CO}_2$ dependence	0.316
α^*	–	$L\phi\rho_{\text{st}}AX\mu$ (see Section 2.2 and below)	3.39×10^5
$k_{\text{eff,ref}}^*$	$\text{mol m}^{-2} \text{yr}^{-1}$	Reference rate constant	8.7×10^{-6}
β	–	Kinetic weathering $p\text{CO}_2$ dependence	0.2 (Rimstidt et al., 2012)
T_e	Kelvin	Kinetic weathering temperature dependence	11.1 (Berner, 1994)

Note. This table lists parameters used in our calculations, their units, their definitions, and the default values they take. A single asterisk (*) means the default parameter value was drawn from Table S1 of the Supplement to Maher and Chamberlain (2014). For default parameters drawn from other sources, the citation is given in the “Value” column.

2.2. Carbon Cycling

To examine how the carbon cycle might operate on abiotic terrestrial planets under very high $p\text{CO}_2$ conditions, we apply an idealized global-mean weathering formulation (Graham & Pierrehumbert, 2020) based on work that accounts for the impact of clay precipitation on weathering solute concentrations (Maher & Chamberlain, 2014) on the global-mean weathering flux,

$$W = \gamma \frac{\alpha}{[k_{\text{eff}}]^{-1} + mA t_s + \alpha [qC_{\text{eq}}]^{-1}}, \quad (6)$$

where W ($\text{mol m}^{-2} \text{yr}^{-1}$) is the global-mean weathering flux, that is, the total number of divalent cations (which react with oceanic carbon to form carbonate minerals, ultimately removing CO_2 from the atmosphere) delivered to the ocean from the land and/or seafloor in a year, divided by the surface area of the planet; γ is the fraction of planetary surface that is weatherable; α is a parameter that captures the effects of various weathering zone properties like characteristic water flow length scale, porosity, ratio of mineral mass to fluid volume, and the mass fraction of minerals in the weathering zone that are weatherable; $k_{\text{eff}} = k_{\text{eff,ref}} \exp\left(\frac{T_{\text{surf}} - T_{\text{surf,ref}}}{T_e}\right) \left(\frac{p\text{CO}_2}{p\text{CO}_{2,\text{ref}}}\right)^\beta$ ($\text{mol m}^{-2} \text{yr}^{-1}$) is the effective kinetic weathering rate, that is, the weathering rate in the absence of chemical equilibration with clay precipitation (Walker et al., 1981); m (kg mol^{-1}) is the average molar mass of minerals being weathered; A ($\text{m}^2 \text{kg}^{-1}$) is the average specific surface area of the minerals being weathered; t_s (yr) is the mean age of the material being weathered; $q = q_{\text{ref}}(1 + \epsilon(T_{\text{surf}} - T_{\text{surf,ref}}))$ (m yr^{-1}) is the volume-weighted global-mean flux of water through the planet’s weathering zones (in this study we apply a linear temperature dependence to runoff, based on the behavior of surface H_2O precipitation on Earth, but we note that the functional form for q could be very different when modeling seafloor weathering); and $C_{\text{eq}} = \Lambda(p\text{CO}_2)^n$ (mol m^{-3}) is the maximum concentration of divalent cations in the water passing through weathering zones, as determined by chemical equilibrium between silicate dissolution and clay precipitation. The values of all constants are given in Table 2. A more thorough set of weathering calculations would carefully account for the lithology of minerals being weathered (Hakim et al., 2021) and the differences between continental and seafloor weathering (Hakim et al., 2021; Hayworth & Foley, 2020), but our calculations are meant to be illustrative of the qualitative behavior

of the carbon cycle under non-terrestrial conditions, so we restrict our simulations to the simplified approach in our Equation 6.

Assuming the presence of weatherable silicates on a planetary surface, the global weathering flux of divalent cations into the ocean, which is equal to the global CO₂ consumption flux at equilibrium and in the absence of CO₂ surface condensation, is dependent on both background CO₂ and T_{surf} . An increase to $p\text{CO}_2$ or T_{surf} leads to an increase in the weathering rate, with the change mediated by either changes to the kinetics of silicate dissolution or changes to global runoff flux, depending on which term in Equation 6 is dominant. This climate-dependence of CO₂ consumption means that silicate weathering can act as a stabilizing negative feedback on planetary climate (Walker et al., 1981). If the CO₂ outgassing flux from volcanoes and other sources (V) is greater than the CO₂ consumption flux from weathering, that is, if $V > W$, and if there is no CO₂ surface condensation, CO₂ will accumulate in the atmosphere, which, under Earth-like circumstances, tends to warm the planet. Higher $p\text{CO}_2$ and higher T_{surf} both lead to larger W , driving the CO₂ consumption rate closer and closer to V until CO₂ consumption is equal to CO₂ production and the atmospheric CO₂ inventory stabilizes. The same process in reverse acts to cool the planet and equilibrate the carbon cycle in cases where $W > V$. So, at least in the cases just discussed, climate on planets with silicate weathering will tend to find an equilibrium T_{surf} and $p\text{CO}_2$ determined by the balance between silicate weathering and CO₂ outgassing, which can be stated simply as

$$V = W, \quad (7)$$

where V (mol m⁻² yr⁻¹) is an assumed CO₂ outgassing flux and W is the weathering flux as defined in Equation 6. The intersection points in $T_{\text{surf}}-p\text{CO}_2$ space of the nullclines given by Equations 4 and 7 are climate states in equilibrium with respect to both energy and carbon fluxes.

The weathering parameterization represented by Equations 6 and 7 implies the assumption of an Earth-like tectonic regime where the resurfacing of fresh silicates occurs rapidly enough to maintain a weathering flux in balance with CO₂ outgassing. This need not be the case: for example, if a planet is in a “sluggish lid”, “episodic lid”, or “stagnant lid” tectonic regime (e.g., Foley, 2015; Kite et al., 2009; Korenaga, 2010; Lenardic, 2018; Lichtenberg et al., 2022; Valencia & O’Connell, 2007), resurfacing may not be fast enough for weathering to keep up with the outgassing rate, leading to a global depletion of weatherable materials called “supply limitation” (West et al., 2005) or “transport limitation” (Kump et al., 2000), which in turn allows for volcanic CO₂ accumulation. Further, even with rapid tectonic resurfacing, the particular climate and/or arrangement of land on a given planet may not allow for high enough weathering fluxes to match outgassing rates, as we will go on to demonstrate. For these cases, it is important to note that surface CO₂ condensation can act as another major sink of atmospheric CO₂ (Kasting, 1991; Wordsworth, Forget, Selsis, et al., 2010; von Paris et al., 2013; Turbet et al., 2017; Kadoya & Tajika, 2019; Bonati & Ramirez, 2021). As a result, under CO₂-condensing conditions, it is possible for the carbon cycle to reach equilibrium even when outgassing does not equal weathering, as condensation can make up the difference,

$$F_{\text{cond}} = V - W, \quad (8)$$

where F_{cond} (mol m⁻² yr⁻¹) is the flux of CO₂ condensing out onto the surface from the atmosphere.

3. Results

3.1. Climate Hysteresis From Temperature-Dependent Instellation Absorption of H₂O

In climate simulations with high $p\text{CO}_2$, cooling by Rayleigh scattering begins to outweigh greenhouse warming, such that temperature eventually begins to decrease while CO₂ increases (Figure 1): in panel A, starting from the lowest $p\text{CO}_2 = 1$ bar, each contour of S_{eff} moves to higher temperatures as CO₂ increases, until a peak T_{surf} is reached at a threshold $p\text{CO}_2$, beyond which T_{surf} for a given S_{eff} begins to decrease as CO₂ increases. For instance, for $S_{\text{eff}} = 0.4$ this peak is at ≈ 280 K and $p\text{CO}_2 = 10$ bar. This occurs because, at high $p\text{CO}_2$, the albedo (panel C) increases more rapidly with $p\text{CO}_2$ than the OLR (panel B) decreases, since the atmosphere has become optically thick at almost all IR wavelengths. In other words, for any given S_{eff} and background gas composition, there is a maximum temperature that cannot be exceeded by adding CO₂ to the atmosphere. This effect has been used to define the outer edge of the classical liquid water HZ as the lowest instellation at which atmospheres with 1 bar N₂, saturated H₂O, and variable CO₂ can maintain an Earth-like planet’s global-mean surface temperature

above freezing (Kasting et al., 1993; Kopparapu et al., 2013), that is, the S_{eff} where the peak temperature is $T_{\text{surf}} = 273.15$ K. Using this “maximum greenhouse limit,” the outer edge of the liquid water HZ has been placed at 1.67 astronomical units (au) (Kopparapu et al., 2013), implying $S_{\text{eff}} = \frac{1}{1.67^2} = 0.359$, with $p\text{CO}_2 \approx 6\text{--}7$ bar. Our fiducial simulations lack N_2 , but, in comparison with Kopparapu et al. (2013), produce a similar value of $S_{\text{eff}} = 0.373$ for the lowest instellation where T_{surf} can be maintained above freezing, with $p\text{CO}_2 \approx 5\text{--}6$ bars, demonstrating our climate model produces comparable results to previous efforts.

At low values of S_{eff} (e.g., along the white-dotted $S_{\text{eff}} = 0.40$ contour in Figure 1a), we find that the climate responds smoothly to increases in $p\text{CO}_2$, with T_{surf} first increasing and then decreasing until reaching a $p\text{CO}_2\text{--}T_{\text{surf}}$ combination that allows the CO_2 to condense at the surface. This is the point where a given S_{eff} contour intersects the black line that bounds the bottom-right gray area in Figure 1a. At this point CO_2 is saturated and our simulations assume that any CO_2 added to the atmosphere simply condenses out onto the surface.

As S_{eff} increases, the maximum temperature climates can reach becomes higher and higher. H_2O is saturated in our simulations and, as such, $p\text{H}_2\text{O}$ increases exponentially with temperature. Therefore, as S_{eff} increases, water’s impact on the climate also becomes more and more prominent. In addition to its well-known greenhouse effect, water can impact planetary albedo via several mechanisms, for instance via cloud and sea ice formation. Less obvious impacts of water on planetary albedo come from its contribution to Rayleigh scattering and its competing contribution to shortwave and near-IR absorption.

At low $p\text{CO}_2$ ($\lesssim 1.5$ bar) and high temperatures where $p\text{H}_2\text{O}$ is comparable to $p\text{CO}_2$, our simulations indicate that the Rayleigh scattering effect of water starts to become important, which is why the albedo contours bend leftward in the upper-left corner of Figure 1c, indicating an increase in albedo from enhanced H_2O Rayleigh scattering as temperature increases. However, at high enough CO_2 ($\gtrsim 1.5$ bar), $p\text{CO}_2$ remains much larger than $p\text{H}_2\text{O}$ even at the highest temperatures we simulated, and hence the Rayleigh scattering effect of H_2O is outweighed by that of CO_2 . This suggests that increases to surface temperature up to 360 K stop significantly increasing a planet’s Rayleigh scattering albedo via H_2O accumulation at $p\text{CO}_2$ above a few bar.

Although H_2O ’s Rayleigh scattering ceases to be important at high $p\text{CO}_2$, H_2O ’s shortwave and near-IR absorption remain important, such that the elevated water content caused by increased temperature leads to increased absorption of instellation: albedo and equilibrium S_{eff} decrease with increased temperature at high $p\text{CO}_2$, as indicated by the rightward tilt of S_{eff} and albedo contours in the upper-right quadrants of Figures 1a and 1c. In other words, at high $p\text{CO}_2$, H_2O saturation leads to temperature-dependent planetary albedo similar to that caused by the ice-albedo feedback, though occurring at temperatures higher than those where the ice-albedo feedback is relevant. This temperature-dependent instellation absorption by H_2O introduces a form of hysteresis into the climate system that is analogous to the hysteresis caused by the ice-albedo feedback (Abbot et al., 2018), the consequences of which we explore here.

3.2. CO_2 Ocean Oscillations on Temperate Exoplanets

Climate limit cycling usually refers to the potential for climates to oscillate back and forth between snowball and temperate states (Abbot, 2016; Haqq-Misra et al., 2016; Menou, 2015; Paradise & Menou, 2017). That occurs when a temperate planet on which weathering dominates outgassing has its CO_2 drawn down until the ice-albedo feedback triggers global glaciation. At this point weathering slows below the rate of outgassing and allows CO_2 to accumulate and eventual deglaciate the planet, which restarts the cycle. Our simulations indicate the existence of a distinct limit cycle that can emerge when outgassing dominates weathering: the oscillation between a CO_2 surface-condensing state and a non- CO_2 -condensing state on planets with $\text{CO}_2\text{--H}_2\text{O}$ atmospheres. This variety of limit cycling is a consequence of the temperature-dependent planetary albedo that arises from H_2O ’s shortwave and near-IR absorption, in combination with CO_2 ’s Rayleigh scattering effect.

In Equation 4, energetic equilibrium between global-mean absorbed instellation and OLR is assumed, and each S_{eff} contour in Figure 1a is a set of $T_{\text{surf}}\text{--}p\text{CO}_2$ pairs where Equation 4 holds for that particular S_{eff} value. However, at high $p\text{CO}_2$, when S_{eff} is large enough to permit the high temperatures that raise water’s vapor pressure enough to lower the planet’s albedo substantially, the right-hand branches of the S_{eff} contours become energetically unstable to perturbations in temperature and $p\text{CO}_2$. The consequences of this phenomenon for planetary climate evolution are illustrated in Figure 2. In this figure, we plot the right-hand branch of the set of T_{surf} and $p\text{CO}_2$ values that produce energetic equilibrium with $S_{\text{eff}} = 0.46$. Any combination of T_{surf} and $p\text{CO}_2$ not falling on the dark red line




Figure 2. Climatic limit cycling driven by CO₂ outgassing, Rayleigh scattering, temperature-dependent near-IR stellar absorption by H₂O, and surface CO₂ condensation. In step (I), outgassing allows CO₂ to accumulate. In step (II), pCO₂ exceeds the value that triggers a positive feedback loop between cooling surface temperature and increasing planetary albedo. In step (III), the planetary surface temperature becomes low enough to trigger surface CO₂ condensation, so continued cooling now also decreases atmospheric pCO₂. In step (IV), a positive feedback loop between warming and albedo reduction from H₂O accumulation leads to runaway warming.

in Figure 2 leads to energetic disequilibrium under an installation of $S_{\text{eff}} = 0.46$. This results in either cooling in the case where the planetary OLR is higher than the absorbed stellar radiation ($OLR > ASR = (1 - \alpha)S/4$) in the light blue region, or warming in the light red region, where $ASR > OLR$.

When the climate resides on the upper, solid portion of the red curve, the temperature responds to energetic disequilibrium as follows: at a given pCO₂, a perturbation in temperature upward from equilibrium is met with a cooling response and a perturbation in temperature downward is met with a warming response, until energetic equilibrium is re-established. Similarly, an increase in CO₂ from equilibrium increases albedo, which causes the planet to cool to maintain equilibrium. Thus small changes to T_{surf} or pCO₂ in this region of parameter space near the solid red curve are met with a negative feedback that tends to maintain energetic equilibrium and move the climate back to the red curve. Starting from somewhere on the red curve near the area labeled (I), if CO₂ in the atmosphere is increased by outgassing, moving rightward along the equilibrium curve, the climate will eventually reach the curve's rightmost point.

If pCO₂ is increased beyond the value at the rightmost point of the red curve, the increase in albedo is enough to push the climate system into a state of energetic disequilibrium where $OLR > ASR$ (the light blue region in Figure 2), and the planet begins to cool. In this region, OLR responds only weakly to changes in temperature because pCO₂ is so high that the atmosphere is mostly opaque in the IR (see Figure 1b), but the albedo responds substantially, increasing as temperature decreases, since H₂O in the atmosphere falls exponentially with temperature, reducing the atmosphere's ability to absorb instellation. This produces a positive feedback with runaway cooling (stage [III] in Figure 2), where a reduction in temperature dries the atmosphere, which decreases ASR and thus pushes the system even further out of energetic equilibrium. This accelerates the cooling and reduces the ASR further. Any plausible rate of cooling vastly exceeds plausible rates of CO₂ accumulation from outgassing (compare thermal equilibration timescale of 1,000 years (Pierrehumbert, 2010) to a carbon cycle timescale of 10⁶ years or more (Colbourn et al., 2015)), so the cooling trajectory is effectively straight down in $T_{\text{surf}}-p\text{CO}_2$ space.

After a temperature reduction of approximately 50 K, the conditions at the surface have cooled enough for CO₂ to begin to condense out onto the planetary surface, which is in this case covered by a liquid H₂O ocean. In stage (III) of the climate cycle, pCO₂ is directly dictated by surface temperature via CO₂'s Clausius Clapeyron relation.

Since the climate is still in the light blue region where $OLR > ASR$, continued cooling drives a rapid decrease in CO_2 partial pressure, resulting in decreasing planetary albedo and increasing ASR .

Eventually, the reduction in CO_2 partial pressure increases ASR enough to re-equilibrate with the OLR at the point where arrow (III) meets arrow (IV) in Figure 2. However, this equilibrium point is unstable to further reductions in CO_2 or increases to temperature, which would shift the climate into the light red region of Figure 2, where $OLR < ASR$ and warming is self-reinforcing due to the accumulation of atmospheric water vapor and resultant reduction in planetary albedo. As a result, any internal climate variability that acted to transiently warm the climate away from this unstable equilibrium would trigger the positive warming feedback loop represented by arrow (IV) in Figure 2, analogous to the cooling feedback loop represented by arrow (II). This warming feedback loop would finally carry the climate back to its initial energetically-stable state, at about 35.6 bar of CO_2 and $T_{surf} \approx 340$ K. From here, assuming outgassing continues, the planet would begin another iteration of this cycle of atmospheric CO_2 accumulation, runaway cooling, CO_2 rain-out, and runaway warming.

For this example of a CO_2 ocean cycle, we chose an instellation that kept the surface temperature at each pCO_2 within the range of temperatures we simulated (≤ 365 K). With higher S_{eff} , the maximum attainable temperature increases, and the unstable righthand branches of the S_{eff} contours shift rightward to higher pCO_2 (see Figure 1). Both of those responses to higher S_{eff} would increase the size of temperature jumps over the course of a limit cycle. Therefore, planets that start off in a limit cycling state at low S_{eff} will undergo cycles of greater and greater amplitude as their star brightens and incident instellation increases.

Up to this point, we have discussed the evolution of climates with CO_2 outgassing but without a complementary weathering feedback. This can correspond to a scenario in which weathering is “supply-limited,” that is, the supply of weatherable minerals to the planetary surface is too low for weathering to keep up with the rate of CO_2 outgassing, or a scenario where liquid CO_2 or CO_2 clathrate hydrate blankets the ocean floor, suppressing weathering reactions in seafloor basalts (a scenario discussed further in Section 4). In the next subsection, we present calculations that include a simple weathering feedback.

3.3. Bistability From the Interaction of Rayleigh Scattering, Weathering, and CO_2 Condensation

With weathering included (Equation 6) and assuming an outgassing rate of 15.8×10^{12} mol yr^{-1} ($2.1 \times$ an Earth-like rate of 7.5×10^{12} mol yr^{-1} drawn from Haqq-Misra et al. (2016)), our model produces a set of weathering-outgassing equilibria corresponding to the solid purple line in Figure 3. The curve has a negative slope because of the pCO_2 -dependence of weathering, with lower T_{surf} required for weathering/outgassing equilibrium at higher values of pCO_2 . A larger outgassing rate would result in the purple line residing at higher temperatures for a given pCO_2 , changing the locations of stable and unstable equilibria, and a smaller outgassing rate would have the opposite effect. With a large enough increase in outgassing (just an increase to $2.2 \times$ the Earth-like rate, in this case), the low- CO_2 solution becomes inaccessible, and with a large enough decrease in outgassing (a reduction to below $1.0 \times$ the Earth-like rate, in this case), the high- CO_2 solution similarly disappears. Changes to the parameters in the weathering model (e.g., changing the assumed global-mean soil thickness) would have analogous effects on the locations and presence of the equilibria. With the formulation of weathering applied here (from Graham & Pierrehumbert, 2020; Maher & Chamberlain, 2014; Winnick & Maher, 2018), the behavior of the system is quite sensitive to changes in outgassing, land fraction, or weathering parameters, while the more traditional kinetically-limited formulation introduced in Walker et al. (1981) would result in less sensitivity, as demonstrated in Graham and Pierrehumbert (2020).

Figure 3 indicates the existence of two equilibrium points where outgassing (V) is balanced by weathering (W), $V = W$, and $OLR = ASR$ for planets at a given S_{eff} , with one equilibrium climate having a higher T_{surf} and a lower pCO_2 (red and green circles a and b in Figure 3) than the other (red and green diamonds c and d). The physical reason for pairs of equilibria at each S_{eff} is CO_2 's cooling effect at high partial pressures. For both S_{eff} values (0.42 and $0.47 S_{Earth}$, green line and red line) plotted in Figure 3, the warmer, lower- pCO_2 climate equilibrium (a and b) is stable with respect to both its energy fluxes and its carbon fluxes, meaning a planet will return to that climate equilibrium if perturbed away from it. These stable climates are the equilibria that are typically explored in studies of silicate weathering on terrestrial planets.

Unlike its counterpart, the second equilibrium climate state where $V = W$ on each S_{eff} curve (c and d) is unstable to climatic perturbations. This is best illustrated by examining unstable equilibrium c with $S_{eff} = 0.42 S_{Earth}$ in




Figure 3. Comparison of stable (circles with black outline: a, b, e) and unstable (diamonds with white outline: c, d, f) climate equilibria at different instillations. The red line represents climate states with $S_{\text{eff}} = 0.47 S_{\text{Earth}}$, with the solid part of the red line representing energetically stable climates and the dashed part of the red line representing energetically unstable climates. The green line represents climates with $S_{\text{eff}} = 0.42 S_{\text{Earth}}$, with the possibility of bistability between an Earth-like scenario and a CO_2 condensing state. The purple line shows where weathering is equal to outgassing; the light red region below is where outgassing outpaces weathering, and the light blue region above is where weathering outpaces outgassing.

Figure 3. As noted earlier, compared to the timescale of carbon cycle response, the thermal equilibration timescale is instantaneous, so for this discussion we can assume that the climate is constrained to move along the green curve at all times. If a planet begins with a climate at point *c*, and its $p\text{CO}_2$ is perturbed downward (leftward on the plot), it warms up because of a reduction in CO_2 Rayleigh scattering and moves upward along the green S_{eff} isoline. This moves the planet into the blue zone of Figure 3 where $W > V$, which means that CO_2 is now being consumed by weathering faster than it can be supplied by outgassing. This imbalance in carbon fluxes leads to further reduction in $p\text{CO}_2$, enhancing the initial perturbation and pushing the climate deeper and deeper into the blue region. Eventually the planet reaches the peak temperature for that S_{eff} , at which point the continued reduction in CO_2 begins to cool the climate, slowing weathering until finally the planet reaches the stable equilibrium point *b*.

Conversely, if a planet begins on the unstable equilibrium *c* and $p\text{CO}_2$ is perturbed upward (to the right on the plot), the planet's surface will cool and the climate will move into the light red area of Figure 3, where outgassing is greater than weathering ($V > W$). With outgassing now outpacing weathering, $p\text{CO}_2$ will continue to grow, enhancing the initial climate perturbation until the $T_{\text{surf}}-p\text{CO}_2$ combination allows for CO_2 condensation at the planetary surface, at which point the carbon cycle has reached a new, stable equilibrium *e* governed by Equation 8, with the imbalance between outgassing and weathering being balanced by surface condensation of CO_2 . This suggests that rocky planets at low instillation can display carbon cycle bistability, where the same geologic boundary conditions (as represented by the parameters in Equation 6) and same stellar environment can drive two very different stable equilibrium climates with $p\text{CO}_2$ differing by an order of magnitude, one of which displays surface CO_2 condensation and one of which does not.

The carbon cycling behavior of a planet irradiated by relative instillation of $S_{\text{eff}} = 0.47 S_{\text{Earth}}$ (the green curve in Figure 3) is considerably different than that of the previous example because of the H_2O absorption-based radiative feedbacks that cause the CO_2 ocean limit cycling discussed in Section 3.2. In this case, even though the intersection point *f* (red-white diamond) between the $S_{\text{eff}} = 0.47 S_{\text{Earth}}$ curve (red, dashed curve) and the CO_2 saturation vapor pressure curve (black) is stable with respect to the carbon cycle, it is unstable with respect to energy fluxes: a small perturbation in temperature upward or CO_2 downward from that point would trigger self-reinforcing warming like that exemplified by arrow (IV) in Figure 2. This would warm the planet to $T_{\text{surf}} \approx 350$ K. This is in




Figure 4. Energetic properties of terrestrial planetary climates irradiated by Sigma Boötis, an F2V star, at low instellation as a function of surface temperature T_{surf} (y-axes) and CO_2 partial pressure (x-axes). Panels and color coding are the same as in Figure 1.

the blue region of Figure 3, where weathering outpaces outgassing, so CO_2 would subsequently be consumed by weathering until the carbon cycle reached the stable equilibrium a , where $V = W$ for $S_{\text{eff}} = 0.47 S_{\text{Earth}}$.

In summary, within the modeling framework applied in this article, a climate transition occurs between $S_{\text{eff}} = 0.42 S_{\text{Earth}}$ and $S_{\text{eff}} = 0.47 S_{\text{Earth}}$ where the climate configurations that allow for surface CO_2 condensation become energetically unstable by the same mechanism that allows for the CO_2 ocean limit cycles described in Section 3.2. However, in this case, the addition of a weathering feedback terminates the cycle before its CO_2 accumulation phase (analogous to step (I) in Figure 2) can be initiated.

3.4. Variations in Stellar Type

As noted in Section 2.1, the fiducial case studied in this article is irradiated by a solar (G2V type) spectrum drawn from Kurucz (1995). Irradiation by spectra corresponding to different stellar types can result in different climate behavior compared to what we have examined so far. Here we examine the impact of an F-type (F2V) spectrum and an M-type (M3.5V) spectrum on our basic results. In all cases, the OLR remains the same, but the planetary albedo is altered by changes to the stellar spectrum impinging on the planet. The general trend is simple to state: planets orbiting hotter, bluer stars can support the bistability between climates with and without CO_2 oceans at higher instillations than cooler, redder stars.

3.4.1. F-Type Stars

To examine the behavior of climates irradiated by F-type stars, we apply the spectrum of Sigma Boötis, an F2V star, drawn from Segura et al. (2003), in Figure 4. Because F-type stars display spectra that are shifted toward higher (bluer) frequencies than G- or M-type stars, the Rayleigh scattering effect of CO_2 is stronger for planets orbiting these stars, as Rayleigh scattering increases greatly in efficacy at shorter wavelengths. Thus, a given increase in $p\text{CO}_2$ leads to a larger increase in albedo for these planets, as demonstrated in Figure 4c, where the planetary albedo reaches nearly 0.68 at the highest $p\text{CO}_2$ shown (compared to a maximum albedo of 0.56 in simulations irradiated by a solar spectrum shown in Figure 1c). This higher sensitivity of albedo to $p\text{CO}_2$ leads to climate configurations where the bistability between climates with and without CO_2 oceans can persist to a higher instellation, reaching $\approx 54\%$ of that of modern-day Earth (Figure 4a), compared to $\approx 45\%$ for climates irradiated by the solar spectrum (Figure 1).




Figure 5. Energetic properties of terrestrial planetary climates irradiated by AD Leonis, an M3.5V star, at low instellation as a function of surface temperature T_{surf} (y-axes) and CO_2 partial pressure (x-axes). Panels and color coding are the same as in Figure 1.

3.4.2. M-Type Stars

To examine the behavior of planets orbiting M-type stars, we apply the spectrum of AD Leonis, an M3.5V star (Segura et al., 2005), in Figure 5. For planets orbiting M-type stars, which emit a larger proportion of their energy at lower (redder) frequencies than G- or F-type stars, the Rayleigh scattering impact of atmospheric $p\text{CO}_2$ is weaker. Even at the highest $p\text{CO}_2$ simulated in this paper (72 bar), albedo does not exceed 0.18 (Figure 5c) for planets orbiting AD Leonis, compared to maximum simulated albedo values of 0.56 and 0.68 for planets orbiting G- and F-type stars, respectively.

Interestingly, this weak dependence of planetary albedo upon CO_2 leads to a complete lack of bistable climate states in our M-star simulations. Without a substantial increase in albedo with $p\text{CO}_2$, CO_2 's reduction of OLR dominates its radiative impact, allowing the molecule to remain a net greenhouse gas across the entirety of the parameter space we studied. Thus, for planets with functional weathering feedbacks that orbit M-type stars, we expect there to be only one stable climate state, corresponding to an Earth-like equilibrium that lacks condensed CO_2 at the surface. Correspondingly, we do not expect planets orbiting M-stars to display limit cycling between CO_2 -condensing and non-condensing states, regardless of outgassing rate or TOA instellation.

4. Discussion

The interaction of liquid CO_2 and liquid H_2O at the planetary surface is fundamental to evaluating the climate state and surface geochemistry of planets with large inventories of both condensed phases. The range of temperatures over which liquid CO_2 would be stable in the presence of a liquid water ocean is relatively narrow (Marounina & Rogers, 2020). At temperatures above 304.5 K CO_2 becomes supercritical and does not condense. At temperatures below 282.91 K and pressures below 4.46 MPa (44.6 bar), mixtures of condensed CO_2 , condensed H_2O , and CO_2 -rich vapor are metastable, with an equilibrium state of CO_2 hydrate, a crystalline phase where water molecules encase CO_2 molecules (Wendland et al., 1999).

Experiments in Earth's ocean demonstrate that hydrate forms rapidly upon contact between liquid CO_2 and liquid H_2O , with visible masses forming over the course of just a few hours (Brewer et al., 1999). This suggests that with surface temperatures below the 282.91 K quadruple point of CO_2 hydrates mentioned in the previous paragraph, precipitation of condensed CO_2 from the atmosphere into a liquid water-rich ocean would result in immediate

formation of solid hydrates that would then sink through the water and settle on the seafloor. This may result in large-scale hydrate build-up on the ocean floor, which would suppress or halt seafloor silicate weathering, similar to high-pressure ice phases on waterworlds with extremely deep oceans (Journaux et al., 2020; Kite & Ford, 2018), removing an important CO₂ sink and making it even more difficult for a planet to exit a stable CO₂ condensing state. Other forms of low-temperature seafloor alteration would also be dramatically altered, with likely major consequences for ocean chemistry (Coogan & Gillis, 2018). Similar layered structures of hydrate and water have also been proposed for icy moons and dwarf planets (Boström et al., 2021), suggesting exploration of such bodies in the solar system may also provide insight into the structure of low-instellation terrestrial planets. Further, assuming a slow rate of subduction, the formation of large seafloor hydrate reservoirs could consume large fractions of the water in planets with Earth-like volatile inventories, since each CO₂ molecule in a hydrate is accompanied by 5.75 H₂O molecules in the most common hydrate structure (Brewer et al., 1999). Thus, under the conditions where CO₂ hydrates are stable (at temperatures below 282.91 K), a planet's subduction rate could exert a powerful direct control on ocean depth and salinity, both of which are first-order parameters in determining planetary climate and surface geochemistry (Olson et al., 2020). Finally, the coexistence of liquid CO₂ and H₂O in the air may lead to the formation of aerial CO₂ hydrates, altering the atmospheric lapse rate through latent heat release (Kasting, 1991), but it is unclear whether this would be an efficient process given the factor-of-a-thousand difference in vapor pressure between CO₂ and H₂O at relevant temperatures.

At temperatures above the hydrate quadruple point (282.9 K) and below CO₂'s triple point (304.5 K), liquid CO₂ and liquid H₂O can coexist. Under pressures like those at the sea surface, liquid CO₂ is less dense than liquid H₂O, so CO₂ that rains into the ocean from the atmosphere would float and form a layer on top of the water (House et al., 2006; Marounina & Rogers, 2020). However, liquid CO₂ is also more compressible than liquid water, such that CO₂ can become denser than water at high enough pressures (House et al., 2006; Marounina & Rogers, 2020). This means that if liquid CO₂ enters the water ocean at a deep enough point, then instead of floating on top of the water, it will sink to the seafloor. Therefore, submarine CO₂ degassing in such conditions would result in the formation of a liquid CO₂ layer at the lithosphere/ocean interface. It is unclear whether silicate weathering by liquid CO₂ is possible, so the impact of this seafloor layer on carbon sequestration is an open question. All in all, the coexistence of liquid CO₂ and liquid H₂O suggests the possibility of “layer cake” oceans at the surfaces of some rocky, low-instellation planets, with an H₂O layer nestled between two CO₂ layers. This analysis remains speculative without detailed thermodynamic modeling, which is beyond the scope of this article but would make for insightful work on the surface conditions of rocky exoplanets under low instellation.

The properties of the atmosphere and ocean under CO₂ condensing conditions also have implications for the viability of origin of life scenarios on prebiotic planets akin to the Hadean Earth. In one popular school of thought regarding the origin of life, surface UV fluxes are held to be important drivers of prebiotic chemistry (Liu et al., 2021; Sasselov et al., 2020). The surface UV flux has been quantified in a variety of models approximating early Earth and early Mars atmospheres (Ranjan & Sasselov, 2016; Ranjan et al., 2017; Rugheimer et al., 2015), and atmospheres with multibar CO₂ pressures demonstrated significant UV attenuation from scattering and absorption (Ranjan et al., 2017). This suggests that atmospheres with 10 s of bar of CO₂ like those expected for the CO₂ condensing cases examined in this paper would not receive enough UV light at their surfaces to drive the relevant prebiotic chemistry. With respect to submarine origin of life scenarios, the seafloor being covered in liquid CO₂ or CO₂ hydrate in conditions with surface CO₂ condensation would preclude water-rock reactions like serpentinization (Sleep et al., 2011) or aqueous organo-metal chemistry near hot vents (Sobotta et al., 2020). It seems that several major models for the origin of life require processes that would be difficult on planets displaying the CO₂ condensing conditions explored here, even with temperate and otherwise habitable surface climates.

Given the likely reduced potential for life to emerge on CO₂ condensing planets, an observational discriminant between these worlds and their non-condensing counterparts would be useful for prioritizing targets in the search for life beyond the solar system. Here, we propose a potential method for distinguishing between CO₂ condensing and non-condensing worlds and, more generally, constraining the $p\text{CO}_2$ of a given exoplanet. This proposal is based on the tendency of CO₂ to dimerize and form molecular complexes at high pressures and low temperatures like those that occur in the CO₂ condensing atmospheres explored in this paper (Asfin et al., 2015; Leckenby & Robbins, 1966; Slanina et al., 1992; Tsintsarska & Huber, 2007). Similar to a method proposed for constraining O₂ partial pressure using spectroscopic O₂ dimer features in exoplanet observations (Misra et al., 2014), we suggest that high resolution spectroscopy from astronomical surveys may be able to detect CO₂ dimer features on

high- $p\text{CO}_2$ exoplanets. Using a quadratic fit to estimate the CO_2 dimerization equilibrium constant as a function of temperature based on data from molecular dynamics simulations (Tsintsarska & Huber, 2007), we find that CO_2 dimers would make up $\approx 5\%$ of the atmosphere by molar fraction at the surface on planets with surface CO_2 condensation and surface temperatures between 273.15 K (H_2O 's freezing point temperature) and 304 K (CO_2 's critical temperature). Planets in the same temperature range but with lower, non-condensing $p\text{CO}_2$ levels have much smaller levels of dimerization; for example, an atmosphere with 6 bar of CO_2 and a surface temperature of 288 K would have a dimer fraction of less than a percent at the surface. Consequently, it may be possible to discriminate between CO_2 -condensing and non- CO_2 -condensing atmospheres using the presence or absence of CO_2 dimer features. These features have been detected in the near- and mid-infrared around $3,700\text{ cm}^{-1}$ ($2.7\text{ }\mu\text{m}$) (Jucks et al., 1987, 1988; Moazzen-Ahmadi & McKellar, 2013), $2,350\text{ cm}^{-1}$ ($4.3\text{ }\mu\text{m}$) (Dehghany et al., 2010; Walsh et al., 1987), and $1,250\text{--}1,400\text{ cm}^{-1}$ ($7.1\text{--}8.0\text{ }\mu\text{m}$) (Asfin et al., 2015; Baranov et al., 2004), placing the features within the proposed wavelength range of future telescope architectures like the Large Interferometer For Exoplanets (Dannert et al., 2022; Quanz, Absil, et al., 2021; Quanz, Ottiger, et al., 2021). Fox and Kim (1988) note the possibility of detection of CO_2 dimers in the atmospheres of Mars and Venus, where they are suggested to exist at parts-per-thousand mole fractions, much lower than the expected dimer abundance in CO_2 condensing atmospheres due to the low pressure and high temperatures on Mars and Venus respectively.

If the proposed method for constraining $p\text{CO}_2$ by dimer feature detection is borne out under more comprehensive examination, it may allow identification of statistical trends in $p\text{CO}_2$ versus instellation. Such a trend is expected if there is a population of Earth-like planets in the HZs of stars with CO_2 levels controlled by silicate weathering, which would introduce a trend of CO_2 decrease with increasing instellation (Bean et al., 2017; Checlair et al., 2019; Lehmer et al., 2020). Conversely, for the population of planets where CO_2 levels are controlled by condensation, the opposite trend would emerge: $p\text{CO}_2$ would increase with increasing instellation. For instance, this trend can be observed in Figure 1a, where CO_2 's saturation vapor pressure curve intersects increasingly large S_{eff} contours as $p\text{CO}_2$ increases. Statistical comparative planetology may thus be able to distinguish populations of terrestrial exoplanets with CO_2 levels controlled by different physical and chemical processes, even if the individual measurements are too low in precision to unambiguously place a given planet into either population. A more detailed analysis of spectral response for the different climate scenarios we outline will be beneficial to analyze the optimal observational architecture (Apai et al., 2019; Quanz, Absil, et al., 2021; Quanz, Ottiger, et al., 2021; The LUVOIR Team, 2019).

4.1. Caveats

Any proposal to analyze the atmospheres of exoplanets is hindered by the potential presence of clouds, and the above is no different: high altitude cloud decks consisting of liquid or solid CO_2 or H_2O may obscure parts of the atmosphere on these CO_2 condensing planets. Clouds might also impact the OLR and albedo of these planets, potentially changing the patterns of climate behavior as a function of $p\text{CO}_2$ and instellation. Thus, the inclusion of clouds may alter the conclusions of this clear-sky study. In particular, if it turns out that thick, global, high-altitude cloud decks obscure the bulk of the planetary atmosphere in most high- $p\text{CO}_2$ climates, the bistability we identify might be muted or eliminated due to the reduced importance of CO_2 Rayleigh scattering of visible light under such conditions. Planetary albedo would instead be determined by cloud properties, with little dependence on $p\text{CO}_2$. We do not include clouds in our simulations for a variety of reasons. Most importantly, it is simply not possible to self-consistently calculate realistic cloud distributions in a one-dimensional model, or even to theoretically estimate cloud deck locations or cloud condensation nuclei density, so any attempt at cloud inclusion would either require arbitrary choices of all of these fundamental parameters for four distinct cloud varieties (solid H_2O , liquid H_2O , solid CO_2 , liquid CO_2) or a many-dimensional parameter space sweep. Further, to our knowledge, the physical and optical properties of liquid CO_2 droplets have not been measured, making it difficult to estimate their impact without making unsupported guesses about their physical properties. The radiative impacts of CO_2 ice clouds have been examined in some detail (Forget & Pierrehumbert, 1997; Forget et al., 2013; Kitzmann, 2016, 2017), and most recent work has found that their net effect on climate is likely to be small under most parameter assumptions, though further work is warranted on this problem. The behavior of water clouds (either solid or liquid) in thick CO_2 atmospheres is relatively under-explored, though some 3D GCM studies of early Mars (Kite et al., 2021; Wordsworth et al., 2013) have examined this regime. Wordsworth et al. (2013) found a small radiative impact from water clouds in the cool, arid climates they simulated. Kite et al. (2021) found a significant climate effect from high-altitude water clouds in arid simulations and a minimal effect in simulations




Figure 6. A schematic illustrating the climate states examined in this article. Planets with high (Earth-like) CO_2 recycling driven by efficient weathering and mobile lid tectonics that receive high stellar irradiation are likely to occupy Earth-like climate states with H_2O oceans and relatively thin atmospheres. Planets with CO_2 recycling that may range from low to high, with either mobile lid or stagnant lid tectonics, and which receive low stellar irradiation, may occupy climate states with thick CO_2 atmospheres and surface CO_2 condensation, producing oceans with both liquid CO_2 and liquid H_2O . Planets with low CO_2 recycling at intermediate instillations may oscillate between a $\text{CO}_2\text{-H}_2\text{O}$ ocean state and a state with only H_2O oceans.

with a global ocean. Our simulations assumed an Earth-like global ocean. Thus, although clouds are a significant source of uncertainty in climate modeling, excluding their effects for a principal examination (similar to studies on the runaway greenhouse effect, e.g., Koppapu et al., 2013) of the phenomena we are studying is justified.

We also neglected the ice-albedo feedback in our simulations because the vast majority of climates we examined had surface temperatures above freezing. Previous studies (e.g., Kadoya & Tajika, 2019; Turbet et al., 2017) have found that planets that fall into globally glaciated states at low instillation may experience surface CO_2 condensation at drastically lower $p\text{CO}_2$ values than the planets we have examined, since the surface temperatures of glaciated planets are tens of Kelvin colder than surface temperatures on temperate planets, though a functional seafloor weathering feedback might be enough to draw down CO_2 to low levels and prevent CO_2 condensation even in a snowball state at low instillations (e.g., Chambers, 2020). The potential for a climate transition from a snowball state with surface CO_2 condensation to a temperate state with continued surface CO_2 condensation is an interesting target for further modeling, and may provide an alternative route to a bistable CO_2 -ocean-bearing climates on Earth-like planets with low instillation. In some regions of parameter space (especially low instillation and high $p\text{CO}_2$), the accumulation of CO_2 and resultant cooling from Rayleigh scattering could itself drive a planet into a snowball state as well, providing another intriguing and counter-intuitive climate scenario for follow-up.

Finally, we note that meridional surface temperature gradients on these planets could lead to CO_2 condensation at somewhat lower surface pressures for a given temperature than calculated here, as the poles tend to be cooler than the global mean surface temperature on planets with Earth-like obliquities and rotation rates, allowing CO_2 surface condensation with less atmospheric CO_2 accumulation. However, meridional surface temperature gradients are greatly reduced at high surface pressures (Chemke & Kaspi, 2017) and atmospheres that are made up mostly of condensable species also tend to have very small meridional temperature gradients due to the powerful winds that develop in response to the large pressure gradients that would be caused by temperature gradients in condensable-rich atmospheres. For example, the equator-to-pole temperature gradient for a pure H_2O atmosphere is calculated to be on the order of ~ 1 K (Ding & Pierrehumbert, 2018).

4.2. Conclusion

In summary, our simulations suggest that the interplay of the radiative properties of CO₂-rich atmospheres and the weathering of silicates leads to super-saturated and cyclic climates for planets under low irradiation for G-type and more massive stars. A qualitative sketch of the distinct climate regimes suggested by our study is shown in Figure 6. The cooling impact of Rayleigh scattering at very high CO₂ levels introduces bistability between a climate state where silicate weathering maintains CO₂ at relatively low levels and a state where CO₂ outgassing outpaces silicate weathering, maintaining CO₂ at such high levels that it condenses at the surface. At intermediate instellation, planetary temperatures can become large enough for water vapor in the atmosphere to significantly impact planetary albedo through absorption of incoming light, destabilizing surface CO₂ condensing climate states and giving rise to limit cycles between CO₂ condensing and non-condensing states. The dynamic interplay between radiation and carbon cycling profoundly impacts the climate state and surface geochemistry of otherwise Earth-like planets in the outer reaches of the liquid water HZ. These CO₂-condensing climate states are potentially distinguishable by observational characterization of CO₂ dimer features and a trend in *p*CO₂ versus instellation opposite to that anticipated from the nominal carbonate-silicate cycle feedback.

Data Availability Statement

We have archived the data necessary to reproduce the plots in this article at Graham et al. (2022).

Acknowledgments

RJG acknowledges scholarship funding from the Clarendon Fund and Jesus College, Oxford. TL was supported by a grant from the Simons Foundation (SCOL Award No. 611576). RTP is supported by European Research Council Advanced Grant EXOCONDENSE (Grant 740963). This AETHER publication is also funded in part by the Alfred P. Sloan Foundation under grant G202114194. We thank Robin Wordsworth, Jim Kasting, and an anonymous reviewer for helpful reviews that significantly strengthened the paper.

References

- Abbot, D. S. (2016). Analytical investigation of the decrease in the size of the habitable zone due to a limited CO₂ outgassing rate. *The Astrophysical Journal*, *827*, 117. <https://doi.org/10.3847/0004-637X/827/2/117>
- Abbot, D. S., Bloch-Johnson, J., Checlair, J., Farahat, N. X., Graham, R. J., Plotkin, D., et al. (2018). Decrease in hysteresis of planetary climate for planets with long solar days. *The Astrophysical Journal*, *854*(1), 3. <https://doi.org/10.3847/1538-4357/aaa70f>
- Agol, E., Dorn, C., & Grimm, S. L. (2021). Refining the transit-timing and photometric analysis of TRAPPIST-1: Masses, radii, densities, dynamics, and ephemerides. *The Planetary Science Journal*, *2*, 1. <https://doi.org/10.3847/PSJ/abd022>
- Apai, D., Milster, T. D., Kim, D. W., Bixel, A., Schneider, G., Liang, R., & Arenberg, J. (2019). A thousand earths: A very large aperture, ultra-light space telescope array for atmospheric biosignature surveys. *The Astronomical Journal*, *158*(2), 83. <https://doi.org/10.3847/1538-3881/ab2631>
- Asfin, R. E., Buldyreva, J. V., Sinyakova, T. N., Oparin, D. V., & Filippov, N. N. (2015). Communication: Evidence of stable van der Waals CO₂ clusters relevant to Venus atmosphere conditions. *The Journal of Chemical Physics*, *142*(5), 051101. <https://doi.org/10.1063/1.4906874>
- Baranov, Y. I., Lafferty, W. J., & Fraser, G. T. (2004). Infrared spectrum of the continuum and dimer absorption in the vicinity of the O₂ vibrational fundamental in O₂/CO₂ mixtures. *Journal of Molecular Spectroscopy*, *228*(2), 432–440. <https://doi.org/10.1016/j.jms.2004.04.010>
- Bean, J. L., Abbot, D. S., & Kempton, E. M.-R. (2017). A statistical comparative planetology approach to the hunt for habitable exoplanets and life beyond the solar system. *The Astrophysical Journal*, *841*(2), L24. <https://doi.org/10.3847/2041-8213/aa738a>
- Berner, R. (1994). GEOCARB II: a revised model of atmospheric CO₂ over Phanerozoic time. *American Journal of Science*, *294*(1), 56–91. <https://doi.org/10.2475/ajs.294.1.56>
- Berner, R. A., Lasaga, A. C., & Garrels, R. M. (1983). The carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years. *American Journal of Science*, *283*(7), 641–683. <https://doi.org/10.2475/ajs.283.7.641>
- Bonati, I., & Ramirez, R. (2021). Monthly notices of the royal astronomical society.
- Boström, M., Estes, V., Fiedler, J., Brevik, I., Buhmann, S. Y., Persson, C., et al. (2021). Self-preserving ice layers on CO₂ clathrate particles: Implications for Enceladus, Pluto, and similar ocean worlds. *Astronomy & Astrophysics*, *650*, A54. <https://doi.org/10.1051/0004-6361/202040181>
- Brewer, P. G., Friederich, G., Peltzer, E. T., & Orr, F. M. (1999). Direct experiments on the ocean disposal of fossil fuel CO₂. *Science*, *284*(5416), 943–945. <https://doi.org/10.1126/science.284.5416.943>
- Chambers, J. (2020). The effect of seafloor weathering on planetary habitability. *The Astrophysical Journal*, *896*(2), 96. <https://doi.org/10.3847/1538-4357/ab94a4>
- Checlair, J. H., Abbot, D. S., Webber, R. J., Feng, Y. K., Bean, J. L., Schwieterman, E. W., et al. (2019). A statistical comparative planetology approach to maximize the scientific return of future exoplanet characterization efforts. *arXiv preprint*. arXiv:1903.05211.
- Checlair, J. H., Villanueva, G. L., Hayworth, B. P. C., Olson, S. L., Komacek, T. D., Robinson, T. D., et al. (2021). Probing the capability of future direct-imaging missions to spectrally constrain the frequency of earth-like planets. *The Astronomical Journal*, *161*(3), 150. <https://doi.org/10.3847/1538-3881/abdb36>
- Chemke, R., & Kaspi, Y. (2017). Dynamics of massive atmospheres. *The Astrophysical Journal*, *845*, 1. <https://doi.org/10.3847/1538-4357/aa7742>
- Claire, M. W., Sheets, J., Cohen, M., Ribas, I., Meadows, V. S., & Catling, D. C. (2012). The evolution of solar flux from 0.1 nm to 160 μm: Quantitative estimates for planetary studies. *The Astrophysical Journal*, *757*(1), 95. <https://doi.org/10.1088/0004-637x/757/1/95>
- Colbourn, G., Ridgwell, A., & Lenton, T. (2015). The time scale of the silicate weathering negative feedback on atmospheric CO₂. *Global Biogeochemical Cycles*, *29*(5), 583–596. <https://doi.org/10.1002/2014gb005054>
- Coogan, L. A., & Gillis, K. M. (2013). Evidence that low-temperature oceanic hydrothermal systems play an important role in the silicate-carbonate weathering cycle and long-term climate regulation. *Geochemistry, Geophysics, Geosystems*, *14*(6), 1771–1786. <https://doi.org/10.1002/ggge.20113>
- Coogan, L. A., & Gillis, K. M. (2018). Low-temperature alteration of the seafloor: Impacts on ocean chemistry. *Annual Review of Earth and Planetary Sciences*, *46*(1), 21–45. <https://doi.org/10.1146/annurev-earth-082517-010027>

- Cox, A. N. (2015). *Allen's astrophysical quantities*. Springer.
- Cronin, T. W. (2014). On the choice of average solar zenith angle. *Journal of the Atmospheric Sciences*, 71(8), 2994–3003. <https://doi.org/10.1175/jas-d-13-0392.1>
- Dannert, F., Ottiger, M., Quanz, S. P., Laugier, R., Fontanet, E., Gheorghe, A., et al. (2022). Large Interferometer for Exoplanets (LIFE): II. Signal simulation, signal extraction, and fundamental exoplanet parameters. *arXiv e-prints*. arXiv:2203.00471.
- Dehghany, M., McKellar, A., Afshari, M., & Moazzen-Ahmadi, N. (2010). High-resolution infrared spectroscopy of carbon dioxide dimers, trimers, and larger clusters. *Molecular Physics*, 108(17), 2195–2205. <https://doi.org/10.1080/00268976.2010.496742>
- Ding, F., & Pierrehumbert, R. T. (2018). Global or local pure condensable atmospheres: Importance of horizontal latent heat transport. *The Astrophysical Journal*, 867(1), 54. <https://doi.org/10.3847/1538-4357/aae38c>
- Edwards, J., & Slingo, A. (1996). Studies with a flexible new radiation code. I: Choosing a configuration for a large-scale model. *Quarterly Journal of the Royal Meteorological Society*, 122(531), 689–719. <https://doi.org/10.1002/qj.49712253107>
- Fischer, R. A., Cottrell, E., Hauri, E., Lee, K. K. M., & Le Voyer, M. (2020). The carbon content of Earth and its core. *Proc. Nat. Acad. Sci. USA*, 117(16), 8743–8749. <https://doi.org/10.1073/pnas.1919930117>
- Foley, B. J. (2015). The role of plate tectonic–climate coupling and exposed land area in the development of habitable climates on rocky planets. *The Astrophysical Journal*, 812(1), 36. <https://doi.org/10.1088/0004-637x/812/1/36>
- Forget, F., & Pierrehumbert, R. T. (1997). Warming early Mars with carbon dioxide clouds that scatter infrared radiation. *Science*, 278(5341), 1273–1276. <https://doi.org/10.1126/science.278.5341.1273>
- Forget, F., Wordsworth, R., Millour, E., Madeleine, J. B., Kerber, L., Leconte, J., et al. (2013). 3D modelling of the early martian climate under a denser CO₂ atmosphere: Temperatures and CO₂ ice clouds. *Icarus*, 222(1), 81–99. <https://doi.org/10.1016/j.icarus.2012.10.019>
- Fox, K., & Kim, S. J. (1988). Spectra of van der Waals complexes (dimers) with applications to planetary atmospheres. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 40(3), 177–184. [https://doi.org/10.1016/0022-4073\(88\)90111-2](https://doi.org/10.1016/0022-4073(88)90111-2)
- Gaudi, B. S., Seager, S., Mennesson, B., Kiessling, A., Warfield, K., Cahoy, K., et al. (2020). The Habitable Exoplanet Observatory (HabEx) mission concept study final report. *arXiv e-prints*. arXiv:2001.06683.
- Gerlach, T. (2011). Volcanic versus anthropogenic carbon dioxide. *Eos, Transactions American Geophysical Union*, 92(24), 201–202. <https://doi.org/10.1029/2011eo240001>
- Gordon, I. E., Rothman, L. S., Hill, C., Kochanov, R. V., Tan, Y., Bernath, P. F., et al. (2017). The HITRAN2016 molecular spectroscopic database. *Journal of Quantitative Spectroscopy & Radiative Transfer*, 203, 3. <https://doi.org/10.1016/j.jqsrt.2017.06.038>
- Graham, R., Lichtenber, T., & Pierrehumbert, R. (2022). Data for “CO₂ ocean bistability on terrestrial exoplanets”. *Zenodo*. <https://doi.org/10.5281/zenodo.6546737>
- Graham, R., Lichtenberg, T., Boukrouche, R., & Pierrehumbert, R. (2021). A multispecies pseudoadiabatic for simulating condensable-rich exoplanet atmospheres. *Planetary Science Journal*, 2, 207. <https://doi.org/10.3847/PSJ/ac214c>
- Graham, R., & Pierrehumbert, R. (2020). Thermodynamic and energetic limits on continental silicate weathering strongly impact the climate and habitability of wet, rocky worlds. *The Astrophysical Journal*, 896(2), 115. <https://doi.org/10.3847/1538-4357/ab9362>
- Grewal, D. S., Dasgupta, R., & Farnell, A. (2020). The speciation of carbon, nitrogen, and water in magma oceans and its effect on volatile partitioning between major reservoirs of the Solar System rocky bodies. *Geochimica et Cosmochimica Acta*, 280, 281–301. <https://doi.org/10.1016/j.gca.2020.04.023>
- Gruszka, M., & Borysow, A. (1997). Roto-translational collision-induced absorption of CO₂ for the atmosphere of venus at frequencies from 0 to 250 cm⁻¹, at temperatures from 200 to 800 K. *Icarus*, 129(1), 172–177. <https://doi.org/10.1006/icar.1997.5773>
- Hakim, K., Bower, D. J., Tian, M., Deitrick, R., Auclair-Desrotour, P., Kitzmann, D., et al. (2021). Lithologic controls on silicate weathering regimes of temperate planets. *The Planetary Science Journal*, 2, 49. <https://doi.org/10.3847/psj/abe1b8>
- Halevy, I., Pierrehumbert, R. T., & Schrag, D. P. (2009). Radiative transfer in CO₂-rich paleoatmospheres. *Journal of Geophysical Research*, 114(D18), D18112. <https://doi.org/10.1029/2009jd011915>
- Haqq-Misra, J., Kopparapu, R. K., Batalha, N. E., Harman, C. E., & Kasting, J. F. (2016). *The Astrophysical Journal*, 827(2), 120. <https://doi.org/10.3847/0004-637x/827/2/120>
- Hayworth, B. P., & Foley, B. J. (2020). Waterworlds may have better climate buffering capacities than their continental counterparts. *The Astrophysical Journal Letters*, 902(1), L10. <https://doi.org/10.3847/2041-8213/abb882>
- Hirschmann, M. M., Bergin, E. A., Blake, G. A., Ciesla, F., & Li, J. (2021). Early volatile depletion on planetesimals inferred from C–S systematics of iron meteorite parent bodies. *Proceedings of the National Academy of Sciences*, 118(13), e2026779118. <https://doi.org/10.1073/pnas.2026779118>
- House, K. Z., Schrag, D. P., Harvey, C. F., & Lackner, K. S. (2006). Permanent carbon dioxide storage in deep-sea sediments. *Proceedings of the National Academy of Sciences*, 103(33), 12291–12295. <https://doi.org/10.1073/pnas.0605318103>
- Journaux, B., Kalousová, K., Sotin, C., Tobie, G., Vance, S., Saur, J., et al. (2020). Large ocean worlds with high-pressure ices. *Space Science Reviews*, 216(1), 7. <https://doi.org/10.1007/s11214-019-0633-7>
- Jucks, K., Huang, Z., Dayton, D., Miller, R., & Lafferty, W. (1987). The structure of the carbon dioxide dimer from near infrared spectroscopy. *The Journal of Chemical Physics*, 86(8), 4341–4346. <https://doi.org/10.1063/1.451895>
- Jucks, K., Huang, Z., & Miller, R. (1988). Infrared spectroscopy of the hydrogen cyanide dimer. *The Journal of Chemical Physics*, 88(10), 2185–2067. <https://doi.org/10.1063/1.454498>
- Kadoya, S., & Tajika, E. (2019). Outer limits of the habitable zones in terms of climate mode and climate evolution of earth-like planets. *The Astrophysical Journal*, 875(1), 7. <https://doi.org/10.3847/1538-4357/ab0aef>
- Karman, T., Miliordos, E., Hunt, K. L. C., Groenenboom, G. C., & van der Avoird, A. (2015). Quantum mechanical calculation of the collision-induced absorption spectra of N₂–N₂ with anisotropic interactions. *Journal of Chemical Physics*, 142(8), 084306. <https://doi.org/10.1063/1.4907917>
- Kasting, J. F. (1991). CO₂ condensation and the climate of early Mars. *Icarus*, 94, 1–13. [https://doi.org/10.1016/0019-1035\(91\)90137-i](https://doi.org/10.1016/0019-1035(91)90137-i)
- Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. (1993). Habitable zones around main sequence stars. *Icarus*, 101(1), 108–128. <https://doi.org/10.1006/icar.1993.1010>
- Kite, E., Manga, M., & Gaidos, E. (2009). Geodynamics and rate of volcanism on massive Earth-like planets. *The Astrophysical Journal*, 700(2), 1732–1749. <https://doi.org/10.1088/0004-637X/700/2/1732>
- Kite, E. S., & Ford, E. B. (2018). Habitability of exoplanet waterworlds. *The Astrophysical Journal*, 864(1), 75. <https://doi.org/10.3847/1538-4357/aad6e0>
- Kite, E. S., Steele, L. J., Mischna, M. A., & Richardson, M. I. (2021). Warm early Mars surface enabled by high-altitude water ice clouds. *Proceedings of the National Academy of Sciences*, 118(18), e2101959118. <https://doi.org/10.1073/pnas.2101959118>

- Kitzmann, D. (2016). Revisiting the scattering greenhouse effect of CO₂ ice clouds. *The Astrophysical Journal Letters*, 817(2), L18. <https://doi.org/10.3847/2041-8205/817/2/L18>
- Kitzmann, D. (2017). Clouds in the atmospheres of extrasolar planets. *Astronomy & Astrophysics*, 600, A111. <https://doi.org/10.1051/0004-6361/201630029>
- Kopparapu, R. K., Ramirez, R., Kasting, J. F., Eymet, V., Robinson, T. D., Mahadevan, S., et al. (2013). Habitable zones around main-sequence stars: New estimates. *The Astrophysical Journal*, 765(2), 131. <https://doi.org/10.1088/0004-637x/765/2/131>
- Korenaga, J. (2010). On the likelihood of plate tectonics on super-Earths: Does size matter? *The Astrophysical Journal Letters*, 725(1), L43–L46. <https://doi.org/10.1088/2041-8205/725/1/L43>
- Krijt, S., Bosman, A. D., Zhang, K., Schwarz, K. R., Ciesla, F. J., & Bergin, E. A. (2020). CO depletion in protoplanetary disks: A unified picture combining physical sequestration and chemical processing. *Acta Pathologica Japonica*, 899(2), 134. <https://doi.org/10.3847/1538-4357/aba75d>
- Kump, L. R., Brantley, S. L., & Arthur, M. A. (2000). Chemical weathering, atmospheric CO₂, and climate. *Annual Review of Earth and Planetary Sciences*, 28(1), 611–667. <https://doi.org/10.1146/annurev.earth.28.1.611>
- Kurucz, R. L. (1995). *CD-ROM 23*. Smithsonian Astrophysical Observatory.
- Leckenby, R., & Robbins, E. (1966). The observation of double molecules in gases. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 291, 389. <https://doi.org/10.1098/rspa.1966.0102>
- Lehmer, O. R., Catling, D. C., & Krissansen-Totton, J. (2020). Carbonate-silicate cycle predictions of Earth-like planetary climates and testing the habitable zone concept. *Nature Communications*, 11, 1. <https://doi.org/10.1038/s41467-020-19896-2>
- Lenardic, A. (2018). The diversity of tectonic modes and thoughts about transitions between them. *Philosophical Transactions of the Royal Society A: Mathematical, Physical & Engineering Sciences*, 376(2132), 20170416. <https://doi.org/10.1098/rsta.2017.0416>
- Li, J., Bergin, E. A., Blake, G. A., Ciesla, F. J., & Hirschmann, M. M. (2021). Earth's carbon deficit caused by early loss through irreversible sublimation. *Science Advances*, 7(14), eabd3632. <https://doi.org/10.1126/sciadv.abd3632>
- Li, J., Scinocca, J., Lazare, M., McFarlane, N., von Salzen, K., & Solheim, L. (2006). Ocean surface albedo and its impact on radiation balance in climate models. *Journal of Climate*, 19(24), 6314–6333. <https://doi.org/10.1175/jcli3973.1>
- Lichtenberg, T., Bower, D. J., Hammond, M., Boukrouche, R., Sanan, P., Tsai, S., & Pierrehumbert, R. T. (2021). Vertically resolved magma ocean–protoatmosphere evolution: H₂, H₂O, CO₂, CH₄, CO, O₂, and N₂ as primary absorbers. *Journal of Geophysical Research: Planets*, 126(2), e2020JE006711. <https://doi.org/10.1029/2020je006711>
- Lichtenberg, T., & Krijt, S. (2021). System-level fractionation of carbon from disk and planetesimal processing. *Acta Pathologica Japonica*, 913(2), L20. <https://doi.org/10.3847/2041-8213/abfdce>
- Lichtenberg, T., Schaefer, L. K., Nakajima, M., & Fischer, R. A. (2022). Geophysical evolution during rocky planet formation. *arXiv e-prints*. arXiv:2203.10023.
- Liu, Z., Wu, L.-F., Kufner, C. L., Sasselov, D. D., Fischer, W. W., & Sutherland, J. D. (2021). Prebiotic photoredox synthesis from carbon dioxide and sulfite. *Nature Chemistry*, 13(11), 1126–1132. <https://doi.org/10.1038/s41557-021-00789-w>
- Maher, K., & Chamberlain, C. (2014). Hydrologic regulation of chemical weathering and the geologic carbon cycle. *Science*, 343(6178), 1502–1504. <https://doi.org/10.1126/science.1250770>
- Marounina, N., & Rogers, L. A. (2020). Internal structure and CO₂ reservoirs of habitable water worlds. *The Astrophysical Journal*, 890(2), 107. <https://doi.org/10.3847/1538-4357/ab68e4>
- Marshall, B. R., & Smith, R. C. (1990). Raman scattering and in-water ocean optical properties. *Applied Optics*, 29(1), 71. <https://doi.org/10.1364/ao.29.000071>
- Menou, K. (2015). Climate stability of habitable Earth-like planets. *Earth and Planetary Science Letters*, 429, 20–24. <https://doi.org/10.1016/j.epsl.2015.07.046>
- Misra, A., Meadows, V., Claire, M., & Crisp, D. (2014). Using dimers to measure biosignatures and atmospheric pressure for terrestrial exoplanets. *Astrobiology*, 14(2), 67–86. <https://doi.org/10.1089/ast.2013.0990>
- Mlawer, E. J., Payne, V. H., Moncet, J. L., Delamere, J. S., Alvarado, M. J., & Tobin, D. C. (2012). Development and recent evaluation of the MT_CKD model of continuum absorption. *Philos. Trans. Royal Soc. A*, 370(1968), 2520–2556. <https://doi.org/10.1098/rsta.2011.0295>
- Moazzen-Ahmadi, N., & McKellar, A. (2013). Spectroscopy of dimers, trimers and larger clusters of linear molecules. *International Reviews in Physical Chemistry*, 32(4), 611–650. <https://doi.org/10.1080/0144235x.2013.813799>
- Öberg, K. I., & Bergin, E. A. (2021). Astrochemistry and compositions of planetary systems. *Physics Reports*, 893, 1–48. <https://doi.org/10.1016/j.physrep.2020.09.004>
- Oki, T., Agata, Y., Kanae, S., Saruhashi, T., Yang, D., & Musiak, K. (2001). Global assessment of current water resources using total runoff integrating pathways. *Hydrological Sciences Journal*, 46(6), 983–995. <https://doi.org/10.1080/02626660109492890>
- Olson, S. L., Jansen, M., & Abbot, D. S. (2020). Oceanographic considerations for exoplanet life detection. *The Astrophysical Journal*, 895(1), 19. <https://doi.org/10.3847/1538-4357/ab88c9>
- Paradise, A., & Menou, K. (2017). GCM simulations of unstable climates in the habitable zone. *The Astrophysical Journal*, 848, 1. <https://doi.org/10.3847/1538-4357/aa8b1c>
- Penman, D. E., Rugenstein, J. K. C., Ibarra, D. E., & Winnick, M. J. (2020). Silicate weathering as a feedback and forcing in Earth's climate and carbon cycle. *Earth-Science Reviews*, 209, 103298. <https://doi.org/10.1016/j.earscirev.2020.103298>
- Pierrehumbert, R. T. (2010). *Principles of planetary climate*. Cambridge University Press.
- Pluriel, W., Marcq, E., & Turbet, M. (2019). Modeling the albedo of Earth-like magma ocean planets with H₂O-CO₂ atmospheres. *Icarus*, 317, 583–590. <https://doi.org/10.1016/j.icarus.2018.08.023>
- Quanz, S. P., Absil, O., Benz, W., Bonfils, X., Berger, J. P., Defrere, D., et al. (2021). Atmospheric characterization of terrestrial exoplanets in the mid-infrared: Biosignatures, habitability, and diversity. *Experimental Astronomy*, 1. <https://doi.org/10.1007/s10686-021-09791-z>
- Quanz, S. P., Ottiger, M., Fontanet, E., Kammerer, J., Menti, F., Dannert, F., et al. (2021). Large Interferometer For Exoplanets (LIFE): I. Improved exoplanet detection yield estimates for a large mid-infrared space-interferometer mission. *arXiv e-prints*. arXiv:2101.07500.
- Ramirez, R. M., & Levi, A. (2018). The ice cap zone: A unique habitable zone for ocean worlds. *Monthly Notices of the Royal Astronomical Society*, 477, 4627. <https://doi.org/10.1093/mnras/sty761>
- Ranjan, S., & Sasselov, D. D. (2016). Influence of the UV environment on the synthesis of prebiotic molecules. *Astrobiology*, 16(1), 68–88. <https://doi.org/10.1089/ast.2015.1359>
- Ranjan, S., Wordsworth, R., & Sasselov, D. D. (2017). Atmospheric constraints on the surface UV environment of Mars at 3.9Ga relevant to prebiotic chemistry. *Astrobiology*, 17(8), 687–708. <https://doi.org/10.1089/ast.2016.1596>
- Rimstidt, J. D., Brantley, S. L., & Olsen, A. A. (2012). Systematic review of forsterite dissolution rate data. *Geochimica et Cosmochimica Acta*, 99, 159–178. <https://doi.org/10.1016/j.gca.2012.09.019>

- Rugheimer, S., Segura, A., Kaltenecker, L., & Sasselov, D. (2015). UV surface environment of Earth-like planets orbiting FGKM stars through geological evolution. *The Astrophysical Journal*, *806*(1), 137. <https://doi.org/10.1088/0004-637x/806/1/137>
- Sasselov, D. D., Grotzinger, J. P., & Sutherland, J. D. (2020). The origin of life as a planetary phenomenon. *Science Advances*, *6*, eaax3419. <https://doi.org/10.1126/sciadv.aax3419>
- Schneider, T., Kaul, C. M., & Pressel, K. G. (2019). Possible climate transitions from breakup of stratocumulus decks under greenhouse warming. *Nature Geoscience*, *12*(3), 163–167. <https://doi.org/10.1038/s41561-019-0310-1>
- Segura, A., Kasting, J., Meadows, V., Cohen, M., Scalzo, J., Crisp, D., et al. (2005). Biosignatures from Earth-like planets around M dwarfs. *Astrobiology*, *5*(6), 706–725. <https://doi.org/10.1089/ast.2005.5.706>
- Segura, A., Krelove, K., Kastings, J. F., Sommerlatt, D., Meadows, V., Crisp, D., et al. (2003). Ozone concentrations and ultraviolet fluxes on Earth-like planets around other stars. *Astrobiology*, *3*(4), 689–708. <https://doi.org/10.1089/153110703322736024>
- Slanina, Z., Fox, K., & Kim, S. J. (1992). A computational evaluation of altitude profiles of the equilibrium monomeric-dimeric CO₂ fractions in the atmosphere of Venus. *Thermochimica Acta*, *200*, 33–39. [https://doi.org/10.1016/0040-6031\(92\)85103-3](https://doi.org/10.1016/0040-6031(92)85103-3)
- Sleep, N. H., Bird, D. K., & Pope, E. C. (2011). Serpentinite and the dawn of life. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *366*(1580), 2857–2869. <https://doi.org/10.1098/rstb.2011.0129>
- Sobotta, J., Geisberger, T., Moosmann, C., Scheidler, C. M., Eisenreich, W., Wachtershauser, G., & Huber, C. (2020). A possible primordial acetyleno/carboxydrotrophic core metabolism. *Life*, *10*(4), 35. <https://doi.org/10.3390/life10040035>
- Soto, A., Mischna, M., Schneider, T., Lee, C., & Richardson, M. (2015). Martian atmospheric collapse: Idealized GCM studies. *Icarus*, *250*, 553–569. <https://doi.org/10.1016/j.icarus.2014.11.028>
- The LUVOIR Team. (2019). The LUVOIR mission concept study final report. *arXiv e-prints*. arXiv:1912.06219.
- Tsintsarska, S., & Huber, H. (2007). The equilibrium constant of carbon dioxide dimerization and the lifetime of the dimer. *Molecular Physics*, *105*(1), 25–31. <https://doi.org/10.1080/00268970601126726>
- Turbet, M., Forget, F., Leconte, J., Charnay, B., & Tobie, G. (2017). CO₂ condensation is a serious limit to the deglaciation of Earth-like planets. *Earth and Planetary Science Letters*, *11*, 11–21. <https://doi.org/10.1016/j.epsl.2017.07.050>
- Valencia, D., O'Connell, R., & Sasselov, D. D. (2007). Inevitability of plate tectonics on super-earths. *The Astrophysical Journal*, *670*(1), L45–L48. <https://doi.org/10.1086/524012>
- Vardavas, I., & Carver, J. H. (1984). Solar and terrestrial parameterizations for radiative-convective models. *Planetary and Space Science*, *32*(10), 1307–1325. [https://doi.org/10.1016/0032-0633\(84\)90074-6](https://doi.org/10.1016/0032-0633(84)90074-6)
- Venturini, J., Guilera, O. M., Haldemann, J., Ronco, M. P., & Mordasini, C. (2020). The nature of the radius valley. *Astronomy and Astrophysics*, *643*, L1. <https://doi.org/10.1051/0004-6361/202039141>
- von Paris, P., Grenfell, J. L., Hedelt, P., Rauer, H., Selsis, F., & Stracke, B. (2013). Atmospheric constraints for the CO₂ partial pressure on terrestrial planets near the outer edge of the habitable zone. *Astronomy & Astrophysics*, *549*, A94. <https://doi.org/10.1051/0004-6361/201219684>
- Walker, J. C. G., Hays, P. B., & Kasting, J. F. (1981). A negative feedback mechanism for the long-term stabilization of Earth's surface temperature. *Journal of Geophysical Research*, *86*(C10), 9776. <https://doi.org/10.1029/jc086ic10p09776>
- Walsh, M., England, T., Dyke, T., & Howard, B. (1987). Pulsed molecular beam infrared absorption spectroscopy of CO₂ dimer. *Chemical Physics Letters*, *142*(3–4), 265–270. [https://doi.org/10.1016/0009-2614\(87\)80935-1](https://doi.org/10.1016/0009-2614(87)80935-1)
- Wendland, M., Hasse, H., & Maurer, G. (1999). Experimental pressure–temperature data on three- and four-phase equilibria of fluid, hydrate, and ice phases in the system carbon dioxide–water. *Journal of Chemical & Engineering Data*, *44*(5), 901–906. <https://doi.org/10.1021/jc980208o>
- West, A. J., Galy, A., & Bickle, M. (2005). Tectonic and climatic controls on silicate weathering. *Earth and Planetary Science Letters*, *235*(1–2), 211–228. <https://doi.org/10.1016/j.epsl.2005.03.020>
- Winnick, M. J., & Maher, K. (2018). Relationships between CO₂, thermodynamic limits on silicate weathering, and the strength of the silicate weathering feedback. *Earth and Planetary Science Letters*, *485*, 111–120. <https://doi.org/10.1016/j.epsl.2018.01.005>
- Wordsworth, R., Forget, F., & Eymet, V. (2010). Infrared collision-induced and far-line absorption in dense CO₂ atmospheres. *Icarus*, *210*(2), 992–997. <https://doi.org/10.1016/j.icarus.2010.06.010>
- Wordsworth, R., Forget, F., Millour, E., Head, J., Madeleine, J. B., & Charnay, B. (2013). Global modelling of the early martian climate under a denser CO₂ atmosphere: Water cycle and ice evolution. *Icarus*, *222*, 1–19. <https://doi.org/10.1016/j.icarus.2012.09.036>
- Wordsworth, R., Forget, F., Selsis, F., Madeleine, J. B., Millour, E., & Eymet, V. (2010). Is Gliese 581d habitable? Some constraints from radiative-convective climate modeling. *Astronomy & Astrophysics*, *522*, A22. <https://doi.org/10.1051/0004-6361/201015053>
- Wordsworth, R. D., & Pierrehumbert, R. T. (2013). Water loss from terrestrial planets with CO₂-rich atmospheres. *The Astrophysical Journal*, *778*(2), 154. <https://doi.org/10.1088/0004-637x/778/2/154>
- Zeng, L., Jacobsen, S. B., Sasselov, D. D., Petaev, M. I., Vanderburg, A., Lopez-Morales, M., et al. (2019). Growth model interpretation of planet size distribution. *Proceedings of the National Academy of Sciences*, *116*(20), 9723–9728. <https://doi.org/10.1073/pnas.1812905116>