



Published in final edited form as:

*Cancer Gene Ther.* 2012 March ; 19(3): 160–170. doi:10.1038/cgt.2011.70.

## Oncolytic Herpes simplex virus expressing yeast cytosine deaminase: relationship between viral replication, transgene expression, prodrug bioactivation

Suguru Yamada<sup>1</sup>, Toshihiko Kuroda<sup>1</sup>, Bryan C. Fuchs<sup>1</sup>, Xiaoying He<sup>2</sup>, Jeffrey G. Supko<sup>2</sup>, Anthony Schmitt<sup>3</sup>, Christopher M. McGinn<sup>3</sup>, Michael Lanuti<sup>3</sup>, and Kenneth K. Tanabe<sup>1</sup>

<sup>1</sup>Division of Surgical Oncology, Massachusetts General Hospital Cancer Center and Harvard Medical School

<sup>2</sup>Division of Hematology/Oncology, Massachusetts General Hospital Cancer Center and Harvard Medical School

<sup>3</sup>Division of Thoracic Surgery, Massachusetts General Hospital Cancer Center and Harvard Medical School

### Abstract

Yeast cytosine deaminase (yCD) is a well-characterized prodrug/enzyme system that converts 5-fluorocytosine (5-FC) to 5-fluorouracil (5-FU), and has been combined with oncolytic viruses. However, *in vivo* studies of the interactions between 5-FC bioactivation and viral replication have not been previously reported, nor have the kinetics of transgene expression and the pharmacokinetics of 5-FC and 5-FU. We constructed a replication-conditional HSV-1 expressing yCD and examined cytotoxicity when 5-FC was initiated at different times after viral infection, and observed that earlier 5-FC administration led to greater cytotoxicity than later 5-FC administration *in vitro* and *in vivo*. Twelve days of 5-FC administration was superior to 6 days in animal models, but dosing beyond 12 days did not further enhance efficacy. Consistent with the dosing schedule results, both viral genomic DNA copy number and viral titers were observed to peak on Day 3 after viral injection and gradually decrease thereafter. The virus is replication-conditional and was detected in tumors for as long as 2 weeks after viral injection. The maximum relative extent of yCD conversion of 5-FC to 5-FU in tumors was observed on Day 6 after viral injection and it decreased progressively thereafter. The observation that 5-FU generation within tumors did not lead to appreciable levels of systemic 5-FU (<10 ng/ml) is important and has not been previously reported. The approaches used in these studies of the relationship between the viral replication kinetics, transgene expression, prodrug administration and anti-tumor efficacy are useful in the design of clinical trials of armed, oncolytic viruses.

Users may view, print, copy, download and text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: [http://www.nature.com/authors/editorial\\_policies/license.html#terms](http://www.nature.com/authors/editorial_policies/license.html#terms)

Mailing Address for Correspondence: Kenneth K. Tanabe, Massachusetts General Hospital, Yawkey 7.924, 55 Fruit Street, Boston, MA 02114. Phone: 617-724-3868, [ktanabe@partners.org](mailto:ktanabe@partners.org).

### CONFLICT OF INTEREST

None of the authors have any competing financial interests in relation to the work described.

**Keywords**

HSV-1; viral oncolysis; cytosine deaminase; pharmacokinetics

---

**INTRODUCTION**

Recent advances in the molecular biology of cancer have led to the development of novel therapeutic agents including viruses that destroy cancer cells by lytic viral replication. This approach has been termed viral oncolysis, and viruses studied for this purpose include adenovirus, Herpes simplex virus 1 (HSV-1), vaccinia virus, reovirus, Newcastle disease virus, Seneca Valley virus and measles virus (1, 2). Several clinical trials with oncolytic viral vectors have been reported, and overall the results demonstrate safety of administration (3, 4). Some of the clinical trials have demonstrated anti-tumor activity of oncolytic viruses (5–7). Nonetheless, efficacy of these viruses has been limited in clinical studies reported to date.

Combination of viral oncolysis with specific anti-cancer drugs may augment anti-tumor activity (8, 9). Prodrug/enzyme cancer therapy mediated by armed oncolytic HSV vectors has been actively investigated as one promising approach (10–12). In particular, yeast cytosine deaminase (yCD) is a well-characterized system that converts non-toxic anti-fungal agent 5-fluorocytosine (5-FC) to cytotoxic chemotherapeutic agent 5-fluorouracil (5-FU) (13). Additionally, converted 5-FU can freely diffuse from yCD expressing cells to neighboring cells (e.g. does not require gap junctions), thereby eliciting a so-called bystander effect (14). One limitation of 5-FU as a systemically delivered agent is the associated toxicity to normal tissues. Prodrug conversion strategies increase 5-FU concentration preferentially within tumors, thereby potentially decreasing systemic side effects and increasing therapeutic efficacy. In addition, not only is 5-FU an effective anti-cancer drug, it is also a potent radiosensitizer and may allow for more efficacious combinations that include radiotherapy. In our previous report, we demonstrated that administration of 5-FC along with this armed HSV-1 vector expressing yCD enhances viral oncolysis (13). The mechanism of action appeared to be local intratumoral generation of 5-FU; however, intratumoral and systemic 5-FU levels were not assessed.

Complex interactions between the viral oncolysis and the bioactivated chemotherapeutic agent can occur in the context of viral replication (15, 16). For example, a bioactivated prodrug which inhibits DNA synthesis may interfere with viral DNA synthesis and limit viral replication and consequently attenuate lytic viral replication and oncolysis (17). In our previous study, we reported that 5-FC bioactivation is associated with only a minimal decrease in HSV-1 replication (13). Other studies demonstrate inhibition of viral replication associated with 5-FC bioactivation (18–20), thereby suggesting that virus strain, cancer cell type and drug concentration influence the balance of the interaction. In the current study we examine the hypotheses that tumors infected by oncolytic HSV-1 expressing yCD produce intratumoral levels of 5-FU without generating appreciable systemic levels of 5-FU; and the efficacy of armed oncolytic viral therapy is dependent on the timing of administration of the 5-FC prodrug relative to the HSV-1 oncolytic virus. We constructed an HSV-1 mutant that

expresses yCD and assessed the impact of the timing of 5-FC relative to HSV-1 administration by determining the kinetics of viral replication and transgene expression. During these studies we measured intratumoral conversion of 5-FC to 5-FU and systemic levels of 5-FU.

## MATERIALS AND METHODS

### Cell lines and viruses

African Green Monkey kidney cells (Vero) and human colon carcinoma cells (HT29) were obtained from the American Type Culture Collection (Rockville, MD). Mouse colon carcinoma cells (MC26) were obtained from the National Cancer Institute Tumor Repository (Frederick, MD). All the cell lines were propagated in Dulbecco's modified Eagle's medium (DMEM) with 10% fetal bovine serum (both from MediaTech CellGro, Manassas, VA), supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin (Invitrogen, Carlsbad, CA). Early passage cells were used for all experiments and they were not reauthenticated. Parent herpes simplex virus type 1 (HSV-1) strain Patton was kindly provided by E. Antonio Chioocca (Ohio State University, Columbus, OH).

### Plasmid and recombinant virus construction

We used an HSV vector cloning system termed Flip-Flop HSV-BAC (bacterial artificial chromosome) system for the generation of new recombinant HSV vectors (21) (kindly provided by Dr. Roberta Martuza, MGH). Briefly, two site-specific recombinases, Cre and FLPe, were used sequentially to integrate desired sequences and to excise the BAC sequences, respectively, and since the size of the HSV-BAC-insert genome exceeds the packaging limit of HSV, only correctly recombined virus grow efficiently. Utilizing this system, we developed a HSV-1 mutant expressing the codon-optimized yCD-uracil phosphoribosyltransferase (UPRT) gene (Invivogen, San Diego, CA) with a myc-tag in the ICP6 locus under the control of the CMV-IE promoter in the HSV strain Patton, termed bMP6-CMVFcumB. A yCD-UPRT fusion gene was used since UPRT has been reported to enhance the yCD rate-limiting enzymatic conversion of 5-FC to 5-FU (22–24).  $\beta$ -galactosidase (LacZ) gene expression is driven by the ICP6 promoter.

### Viral cytotoxicity assay

Cells were plated in triplicate at a density of  $4 \times 10^4$ /ml in a 24-well plate. After 24h, cells were infected with virus at the indicated multiplicity of infection (MOI) and then media containing 5-FC at the indicated concentrations was added. After 120 h, the effect of virus with or without 5-FC on the growth of cell lines was measured by colorimetric 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT; Sigma, St. Louis, MO) assay as previously described (13). The experiment was repeated twice for each cell line.

### Detection of HSV-1 Gene and Transgene Expression

Detection of HSV-1 gene and transgene expression by western blot was performed as previously described (25). The following antibodies were used for the detection: 4A6 anti-Myc monoclonal antibody (Millipore), anti-HSV-ICP4 monoclonal antibody (US Biological, Swampscott, MA) and  $\beta$ -actin (Abcam, Cambridge, MA).

### In vivo studies

Six- to eight-week-old male BALB/c mice (Charles River Laboratories, Shrewsbury, MA) were maintained in accordance with the institutional guidelines of the Massachusetts General Hospital animal care facility. MC26 cells were implanted subcutaneously into the flank area of mice ( $1 \times 10^7$  cells/100 $\mu$ l of DMEM) and bMP6-CMVFcumB virus ( $1 \times 10^7$  plaque-forming units (pfu)/50 $\mu$ l of PBS with 10% glycerol) was injected intratumorally when tumor volumes reached 100 mm<sup>3</sup>. Mice were stratified by tumor volume and then randomly assigned to each treatment group (n = 5). 5-FC (750 mg/kg) was administered intraperitoneally. Tumor sizes were measured every 2 days by external caliper and the volume calculated as length  $\times$  width  $\times$  depth  $\times$  0.52. Each blood sample was drawn 24 h after last 5-FC was injected intraperitoneally. These blood samples were centrifuged at 2,000 rpm for 10 min and the plasma was used for the subsequent assay. Each tumor specimen was also collected 24 h after last 5-FC was injected. These tumor specimens were minced by sterilized scalpel and put into tubes with PBS and garnet beads (MO BIO Laboratories, Carlsbad, CA), and then vortexed for 15 min at 4°C. The samples were stored at -80°C. The experiment was repeated in duplicate to ensure reproducible results.

### Quantitative real-time PCR

Mouse flank tumor samples were harvested and homogenized at designated time points by ATL buffer (Qiagen, Germantown, MD). DNA was isolated from each sample according to the manufacturer's protocol (QIAamp DNA Mini Kit; Qiagen). HSV-1 DNA, Macintyre strain (Advanced Biotechnologies Inc., Columbia, MD) was used to create a quantitative standard control. Viral genomic DNA copy number was analyzed by quantitative real-time PCR (LightCycler; Roche Diagnostics Corporation, Indianapolis, IN). Primer sequences to ICP6 were as follows: forward TCTGGAGTCTGTGCGAATTTCA and reverse ATACATGCTGCGCTTAAAGTG. All reactions were done in duplicate and the experiment was repeated to ensure reproducible results.

### Plaque assay

Monolayer cultures of Vero cells grown in a six-well plate were infected with serial dilutions of virus from tumor homogenates as indicated. After removal of virus inoculum, the cells were incubated in DMEM supplemented with 1% inactivated fetal bovine serum and 0.1% pooled human immunoglobulin (Talecris Biotherapeutics, Research Triangle Park, NC) at 37° C for 3 days until plaques were visible. The cells were stained by 5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside (X-gal; Sigma) histochemistry and  $\beta$ -galactosidase-positive plaques were counted.

### LacZ staining

Five-micrometer-thick sections of frozen tumors in OCT mounted on microscope slides were fixed in 0.5% glutaraldehyde.  $\beta$ -Galactosidase activity was measured by incubating the slides in X-gal solution at 37° C overnight. X-gal solution was prepared as follows: X-gal at 20 mg/ml in dimethyl formamide, 0.1 mol/l potassium ferric cyanide, 0.1 mol/l potassium ferrocyanide, and 0.1 mol/l MgCl<sub>2</sub> (all from Sigma). Sections were coverslipped before examination under a light microscope.

## Determination of 5-FC and 5-FU in Plasma and Tumor Homogenate

The concentration of 5-FC and 5-FU in plasma and tumor tissue homogenate was measured by liquid chromatography-tandem mass spectrometry (LC-MS/MS). The procedure used to prepare samples for analysis, the chromatographic conditions, and tandem mass spectrometric detection were adapted from several analytical methods that have been previously reported (26, 27). Analytical reference samples of 5-FC, 5-FU, and 5-bromouracil (5-BU), the latter of which was used as the internal standard (IS), were purchased from Sigma-Aldrich (St. Louis, MO). Stock solutions with concentrations of approximately 1.0 mg/ml, based upon the actual amount weighed to an accuracy of 0.001 mg using a Cahn microbalance (Analytical Technology, Inc, Boston, MA), were made in water for 5-FC and methanol for 5-FU and 5-BU. A single analyte working solution containing both 5-FC and 5-FU at a concentration of 0.100 mg/ml was made by diluting the stock solutions with acetonitrile:water (1:1, v/v). Calibration standards were made by serially diluting the analyte stock solution with mouse plasma (Innovative Research, Novi, MI) to provide 8 solutions with concentrations ranging from 10-1000 ng/ml. Quality control samples were similarly prepared with concentrations of 30, 300, and 900 ng/ml. The IS working solution was made by quantitatively diluting the stock solution to 1.50 µg/ml with acetonitrile:water (1:1, v/v).

Frozen plasma samples and tissue homogenates were allowed to thaw over ice and mixed on a vortex stirrer. In a polypropylene microcentrifuge tube, 25 µl of each sample was vigorously mixed with 5 µl of IS working solution and 75 µl of chilled acetonitrile by vortexing for 30 sec. The mixture was centrifuged (12,000 g, 5-min) to sediment precipitated protein and 80 µl of the clear supernatant was transferred into another microcentrifuge tube and evaporated using a centrifugal vacuum concentrator at 40°C (Labconco, Kansas City, MO). The extract was reconstituted in 50 µl of an ammonium acetate buffer (10 mM, pH 3.76) assisted by vortexing and sonicating. An autosampler was used to inject 25 µl of this solution onto a Synergi 4 µm Hydro-RP 150 × 4.6 mm column (Phenomenex, Torrance, CA) preceded by a Phenomenex AQ C18 4.0 × 3.0 mm guard cartridge and 0.5 µm inline filter. The column was eluted at 0.5 ml/min with an isocratic mobile phase composed of 25% methanol and 75% ammonium acetate buffer (10 mM, pH 3.76), delivered using a quaternary pump. At the end of the 6.5 min run, the flow rate was increased to 1.0 ml/min to flush the column with 95% methanol for 3 min and reequilibrate it with the initial composition of the mobile phase for 3 min before starting the next run. Flow from the analytical column was directed into the spray chamber of an Agilent 6410 Triple Quad LC/MS system equipped with an API electrospray source (Agilent Technologies, Inc., Santa Clara, CA) from 0 - 6.5 min during the run and to waste at all other times. Nitrogen was used as the nebulizing gas at 60 p.s.i. and as the drying gas at a flow rate of 13 liters/min and a temperature of 350°C. The capillary voltage was 1,500 V during positive ion detection and -1,200 V for negative ion detection. MS/MS detection was performed using high purity nitrogen as the collision gas and operating the instrument in the multiple reaction monitoring mode. The polarity of the mass spectrometer was positive for monitoring 5-FC and negative for monitoring 5-FU and 5-BU. The individualized operating parameters that provided maximum response for each compound were as follows. 5-FC: transition,  $m/z$  130.0 → 113.0; fragmentor voltage, 110 V; collision energy, 20 V. 5-FU:

transition,  $m/z$  129.0→ 42.0; fragmentor voltage, -80 V; collision energy, -16 V. 5-BU: transition,  $m/z$  189.0→ 42.0; fragmentor voltage, -85 V; collision energy, -16 V. The dwell time was 70 ms for each transition and the quadrupoles were maintained at unit resolution. Quantitation was based upon integrating the extracted ion chromatograms for each transition.

The analytical method was thoroughly validated according to current recommendations (28). Average  $\pm$  SD retention times were  $3.67 \pm 0.02$  min for 5-FC,  $4.03 \pm 0.01$  min for 5-FU, and  $5.69 \pm 0.02$  min for 5-BU. Peaks that interfered with detection of either compound were not evident in chromatograms of pretreatment mouse plasma or tumor homogenate. Study samples were prepared for analysis together with a complete set of calibration standards, drug-free plasma with and without addition of the IS, a set of quality control samples and run in a computer-generated randomized order. The relationship between the 5-FC/IS and 5-FU/IS peak area ratios ( $y$ ) and known concentration of the analyte in each calibration standard ( $x$ ) were fit to the equation for a straight line ( $y = ax + b$ ) or an exponential equation ( $y = a + b \cdot x^c$ ), respectively, using WinNonlin Professional 5.0 software (Pharsight Corp., Cary, NC). Values of the parameters for the best-fit equations were used to calculate the analyte concentrations in study samples. Concentrations of 5-FC and 5-FU in tumor tissue, expressed as  $\mu\text{g/g}$  tissue weight, was calculated by multiplying their measured concentrations in the homogenate by the combined volume of tumor tissue and PBS added to prepare the homogenate and dividing the product by the tumor tissue volume, assuming that the density of 1 g/ml for tumor tissue. The geometric mean concentration of 5-FC and 5-FU was calculated from the observed concentrations for the group of animals sampled at each time point.

The average correlation coefficient for the calibration curves was 0.999 for both 5-FC and 5-FU. Interday accuracy for assaying both compounds in the quality control samples ranged from 95.0-103.9% of the known concentrations with a precision of 6.5-10.0%. 5-FC and 5-FU were measured at their 10 ng/ml lower limits of quantitation with an accuracy <108% and precision <8%. The limits of detection were 2 ng/ml for 5-FC and 1 ng/ml for 5-FU.

### Statistical analysis

Statistical analyses were performed with a Student's  $t$  test.  $p < 0.05$  was considered to be significant.

## RESULTS

### Construction of a replication-conditional HSV-1 expressing the yCD-UPRT gene

HSV-1 mutant expressing the myc-tagged yCD-UPRT gene in the ICP6 locus under the control of the CMV-IE promoter was constructed from HSV-1 strain Patton, using the FlipFlop HSV-BAC system (Fig. 1A) (21). After three rounds of limiting dilution for purification, viral isolates were selected with the correct genetic structure for each confirmed. One viral isolate termed BMP6-CMVFcumB was selected for subsequent experiments.



To confirm the protein expression of the yCD-UPRT gene, HT29 cells were infected with BMP6-CMVFcumB at an MOI of 2. Cell extracts after virus infection for 20 h showed protein expression of yCD-UPRT as assessed by western blot for the myc-tag (Fig. 1B).

### Addition of 5-FC enhances cytotoxicity against cancer cell line

We evaluated cytotoxicity in HT29 and MC26 cells infected with BMP6-CMVFcumB at a constant MOI of 0.01 or 0.185, respectively, in the presence of increasing concentrations of the prodrug 5-FC. Cell killing 5 days after viral infection was determined by MTT assay. Compared to the control virus BMP6-empty, BMP6-CMVFcumB produced increased cytotoxicity with increasing 5-FC concentrations, with a 50% cytotoxicity observed at approximately 50 µg/mL for HT-29 and 10 µg/ml for MC26 (Fig. 2A,B). We then explored the cytotoxicity against HT29 and MC26 cells infected with BMP6-CMVFcumB at increasing MOI ranging from 0.0001 to 1.6 in the presence or absence of constant concentration of 5-FC (50 µg/ml). 5-FC conferred increased cytotoxicity to HT29 cells at MOI values ranging from 0.01 to 0.1 (Fig. 2C) and to MC26 cells at MOI values ranging from 0.02 to 1.6 (Fig. 2D). When both of these cell lines were infected with BMP6-empty virus at same MOI in the presence or absence of 5-FC, as expected the addition of 5-FC did not enhance cell killing (Fig. 2E,F).

This strategy of yCD-UPRT expression in combination with 5-FC has the theoretical potential to interfere with viral replication and thereby reduce overall anti-tumor efficacy (13, 29). Accordingly, we examined cytotoxicity when 5-FC initiation was varied after BMP6-CMVFcumB infection of HT29 cells, as well as infection of murine MC26 cells. Bioactivation of 5-FC enhances cytotoxicity beyond that seen by lytic replication alone. Early 5-FC initiation was more effective than late initiation in HT29 cells, and this difference was more apparent at relatively high MOI (Fig. 3A). Similar results were seen with MC26 cells (Fig. 3B).

### Optimal dosing schedule of 5-FC in mouse xenograft model

In order to examine the influence of timing of 5-FC initiation *in vivo*, BALB/c mice with MC26 flank tumors were treated by intratumoral injection of BMP6-CMVFcumB and 5-FC was administered intraperitoneally. The initiation date for 5-FC was varied, while the duration of 5-FC was held constant (6 days). As shown in Fig. 4A, Group 4 received 5-FC on days 0 through 6; Group 5 received 5-FC on days 3 through 9; and Group 6 received 5-FC on days 6 through 12. A single intratumoral injection of BMP6-CMVFcumB virus alone (no-5-FC administered) resulted in inhibition of tumor growth compared with controls (no treatment or 5-FC alone) (Group 1 vs. Group 3;  $p=0.0022$ , Group 2 vs. Group 3;  $p=0.0062$ , respectively). Also, the addition of 5-FC on Day 0 or Day 3 enhanced the anti-tumor activity of BMP6-CMVFcumB significantly (Group 3 vs. Group 4;  $p=0.011$ ; Group 3 vs. Group 5;  $p<0.001$ ). Importantly, 5-FC initiation on Day 3 showed greater inhibition of tumor growth compared with late 5-FC initiation (Day 6) (Group 5 vs. Group 6;  $p=0.036$ ), and there was also a trend whereby initiation on Day 3 was more effective than initiation on Day 0 but this did not reach significance. The difference between initiation on Day 0 and initiation on Day 6 was not statistically significant (data not shown).

The optimal duration of 5-FC administration was assessed in the same mouse model. We investigated different durations of 5-FC administration after initiation on Day 3 after viral injection. Virus monotherapy resulted in inhibition of tumor growth compared to 5-FC alone (Group 1 vs. Group 2;  $p=0.004$ ), and treatment with virus and 5-FC administration also resulted in significant inhibition compared with virus monotherapy (Group 1 vs. Group 3;  $p=0.022$ ). Importantly, 5-FC administration for 12 days showed further inhibition of tumor growth than 5-FC administration for 6 days, however, 5-FC beyond 12 days provided no additional benefit (Fig. 4B). These results indicated that optimal duration of 5-FC administration in this model was for 12 days and dosing for more than 12 days does not enhance tumor inhibition.

### Viral replication and transgene expression in tumor cells

In order to better understand these results, we explored the kinetics of viral replication and transgene expression in this flank tumor model. bMP6-CMVFcumB virus was administered intratumorally into MC26 tumors on flanks of BALB/c mice, and the tumors were subsequently harvested at different time points between Day 0 and Day 21 following virus injection. Tumor homogenates were analyzed by quantitative real-time PCR to assess genomic HSV-DNA, viral plaque assay to assess replicative virions and LacZ staining to assess transgene expression. Both viral genomic DNA copy number and viral titers peaked on Day 3 after viral injection and the values decreased gradually (Fig. 5A, B). Importantly, these assays indicated that virus persisted at least 2 weeks after viral injection. Similarly, LacZ staining showed that transgene expression was evident in tumor homogenates between Day 3 and Day 15 after viral injection, however, no LacZ staining was observed in Day 21 tumor samples. These experimental results on viral titer and transgene expression kinetics provide support for the observations that 5-FC administration starting on Day 3 after viral injection was superior to earlier or later initiation, and that 5-FC administration for a longer duration (up to 12 days) enhanced tumor destruction.

### Mouse Pharmacokinetic Studies

A highly sensitive and specific LC-MS/MS assay was used to measure the concentrations of 5-FC and 5-FU in tumor tissue and plasma samples obtained after 3 to 18 days of daily treatment with 15 mg 5-FC, given by intraperitoneal injection, beginning 3 days after injecting the bMP6-CMVFcumB virus into subcutaneous MC26 flank tumors. The mean plasma concentration of 5-FC, which depends upon the time interval between administration of the prior dose and collection of the sample, ranged from  $0.051 \pm 0.020$   $\mu\text{g/ml}$  (Day 9) to  $2.14 \pm 8.10$   $\mu\text{g/ml}$  (Day 6) on the four days during which samples were obtained (Fig. 6). Intratumoral concentrations of 5-FC were similar to its concentration in plasma and ranged from  $0.080 \pm 0.017$   $\mu\text{g/g}$  (day 9) to  $1.46 \pm 5.41$   $\mu\text{g/g}$  (day 6). Intratumoral concentrations of 5-FC were similar to its concentration in plasma and ranged from  $0.080 \pm 0.017$   $\mu\text{g/g}$  (day 6) to  $1.46 \pm 5.41$   $\mu\text{g/g}$  (day 3). 5-FU was present at measurable concentrations in tumor samples from at least 4 of the 5 mice evaluated at each time point. The mean intratumoral concentration of 5-FU relative to the corresponding intratumoral concentration of 5-FC decreased progressively from  $0.49 \pm 1.32$  after 3 days of dosing to  $0.038 \pm 0.059$  after 18 days of dosing. This finding substantiates that the relative extent of 5-FC conversion to 5-FU decreases progressively over time following viral injection, which indeed corresponds with



the kinetics of yCD expression. The concentration of 5-FU in plasma was below the 10 ng/ml lower limit of quantitation of the LC-MS/MS assay in all samples obtained at each time interval and undetectable (<1 ng/ml) in 70% of the samples. These results demonstrate that combination therapy with a cytosine deaminase expressing virus and the prodrug 5-FC achieves therapeutic intratumoral concentrations of 5-FU without significant systemic exposure to the cytotoxic anticancer drug.

## DISCUSSION

Results of several recent clinical trials of HSV-1 mutants for viral oncolysis have demonstrated safety of these viruses for cancer therapy. In each of these clinical trials, the HSV-1 mutant tested had been genetically altered to attenuate the pathological virulence. This strategy enhances safety for clinical application, but also attenuates the oncolytic efficiency compared with wild-type HSV-1. Therefore, we and others have explored strategies to enhance HSV-1 viral oncolysis by arming HSV-1 mutants with genes to enhance prodrug activation. Both *in vitro* and *in vivo* studies have demonstrated this approach to be more efficacious than treatment with the virus alone (30–32). The yCD gene as a prodrug activation gene has several advantages in this application (13). Specifically, 5-FU has been shown to increase viral replication *in vitro* and extend animal survival *in vivo* (33). In addition the additive and synergistic effect of combining HSV-1 oncolysis with 5-FU (rather than bioactivation of 5-FC) has been reported as well (34). Several studies have suggested that enhancement of viral replication by chemotherapeutics leading to increased cell death could be a possible explanation for the detected synergistic effects (35–37). Inhibition of viral replication by chemotherapy or activated prodrugs may also be observed, and is caused either by direct interference with viral DNA synthesis or by killing surrounding cells prior to their infection. In anticipation of designing clinical trials with yCD-expressing HSV-1 mutants, we examined the relationship between the kinetics of viral replication, timing of 5-FC administration, and anti-tumor efficacy.

With HSV-1 administered via direct intratumor injection, viral DNA is immediately measurable, and following replication viral titers peak 3 days later. Transgene expression is observed in the infected tissue on day 3 and persists through day 15, with onset a bit delayed compared to onset of viral replication. Based on these kinetics, it is not a surprising observation that making 5-FC available during the height of infection (Day 3) leads to more effective tumor kill than initiating 5-FC before peak viral replication and transgene expression (Day 0) or waiting until Day 6 to introduce the prodrug. And declining concentrations of 5-FU in the tumors beyond day 6 are an expected observation and correlate with declining intratumoral transgene expression. Nonetheless, it is notable that virus and transgene expression are detectable in the tumors for at least 15 days following a *single* intratumoral injection, and bioactivated 5-FC (e.g. 5-FU) is measurable in the tissue for up to 21 days. HSV-1 replication was not greatly affected by 5-FC at a concentration of 50 µg/mL, which is consistent with previous findings (13). These data suggests that the added cytotoxicity achieved by combining bMP6-CMVFcumB with 5-FC exceeds any inhibition of viral replication by 5-FU metabolites. We did not repeat the experiments in a model of multiple viral administrations, but it is reasonable to assume that the principles elucidated in the current experiments would continue to apply.

The inhibition of tumor growth in our mouse model was modest even with prodrug administration. These results may be in part due to the single virus injection used in our study as discussed above. It is more likely though explained by our use of a syngeneic mouse model. Mouse cells are typically less infectable with HSV-1 than human cells, and therefore cell killing would be expected to be more dramatic in human tumors.

At present, detection of viral replication and transgene expression requires acquisition of tissue followed by analysis using molecular techniques such as immunohistochemical staining and PCR amplification. These assays are accurate and sensitive, but they require sacrifice of the mouse at each time point, thereby limiting comparison of measurements over time in the same animal. Recently, in order to measure the magnitude and anatomic sites of viral infection or transgene activity *in vivo*, some techniques such as magnetic resonance imaging (MRI) and positron emission tomography (PET) have been developed (38–41). However, to our knowledge, there is no study employing these techniques to measure the kinetics of viral replication and transgene expression in combination with prodrug over a prolonged period of time in a mouse model.

In this prodrug/enzyme system, the levels of 5-FU in plasma and tumors are a matter of concern. It has been suggested that  $\gamma$ CD delivered by an oncolytic HSV-1 vector could be used to reduce the side effects of 5-FU by promoting a localized tumor conversion of 5-FC to 5-FU without raising systemic 5-FU levels (42). But while previous studies in preclinical models have analyzed intratumoral 5-FU concentration in tumors treated with viruses that express  $\gamma$ CD and 5-FC (43–45), the relationship between intratumoral 5-FU concentration and plasma 5-FU concentration over time has never been explored. This study is the first to measure the systemic levels in such model, and confirms that 5-FU levels were below the level of detection in almost all of the plasma samples. By assessing viral replication and transgene expression kinetics, we were able to establish an optimal dosing schedule for this BMP6-CMVFcumb virus in combination with 5-FC based on the kinetics of viral replication and transgene expression. The approaches used in these studies are useful in the design of clinical trials of armed, oncolytic viruses.

## Acknowledgments

This work was supported by NIH grants R01CA76183 (KKT) and 5K01CA140861 (BCF)


## References

1. Parato KA, Senger D, Forsyth PA, Bell JC. Recent progress in the battle between oncolytic viruses and tumours. *Nat Rev Cancer*. 2005; 5:965–76. [PubMed: 16294217]
2. Cattaneo R, Miest T, Shashkova EV, Barry MA. Reprogrammed viruses as cancer therapeutics: targeted, armed and shielded. *Nat Rev Microbiol*. 2008; 6:529–40. [PubMed: 18552863]
3. Reid T, Galanis E, Abbruzzese J, et al. Hepatic arterial infusion of a replication-selective oncolytic adenovirus (dl1520): phase II viral, immunologic, and clinical endpoints. *Cancer Res*. 2002; 62:6070–9. [PubMed: 12414631]
4. Geevarghese SK, Geller DA, de Haan HA, et al. Phase I/II study of oncolytic herpes simplex virus NV1020 in patients with extensively pretreated refractory colorectal cancer metastatic to the liver. *Hum Gene Ther*. 2010; 21:1119–28. [PubMed: 20486770]

5. Park BH, Hwang T, Liu TC, et al. Use of a targeted oncolytic poxvirus, JX-594, in patients with refractory primary or metastatic liver cancer: a phase I trial. *Lancet Oncol.* 2008; 9:533–42. [PubMed: 18495536]
6. Senzer NN, Kaufman HL, Amatruda T, et al. Phase II clinical trial of a granulocyte-macrophage colony-stimulating factor-encoding, second-generation oncolytic herpesvirus in patients with unresectable metastatic melanoma. *J Clin Oncol.* 2009; 27:5763–71. [PubMed: 19884534]
7. Fong Y, Kim T, Bhargava A, et al. A herpes oncolytic virus can be delivered via the vasculature to produce biologic changes in human colorectal cancer. *Mol Ther.* 2009; 17:389–94. [PubMed: 19018254]
8. Heise C, Sampson-Johannes A, Williams A, McCormick F, Von Hoff DD, Kirn DH. ONYX-015, an E1B gene-attenuated adenovirus, causes tumor-specific cytolysis and antitumoral efficacy that can be augmented by standard chemotherapeutic agents. *Nat Med.* 1997; 3:639–45. [PubMed: 9176490]
9. Khuri FR, Nemunaitis J, Ganly I, et al. A controlled trial of intratumoral ONYX-015, a selectively-replicating adenovirus, in combination with cisplatin and 5-fluorouracil in patients with recurrent head and neck cancer. *Nat Med.* 2000; 6:879–85. [PubMed: 10932224]
10. Chase M, Chung R, Chiocca EA. An oncolytic viral mutant that delivers the CYP2B1 transgene and augments cyclophosphamide chemotherapy. *Nat Biotech.* 1998; 16:444–8.
11. Pawlik TM, Nakamura H, Mullen JT, et al. Prodrug bioactivation and oncolysis of diffuse liver metastases by a herpes simplex virus 1 mutant that expresses the CYP2B1 transgene. *Cancer.* 2002; 95:1171–81. [PubMed: 12209705]
12. Tyminski E, Leroy S, Terada K, et al. Brain tumor oncolysis with replication-conditional herpes simplex virus type 1 expressing the prodrug-activating genes, CYP2B1 and secreted human intestinal carboxylesterase, in combination with cyclophosphamide and irinotecan. *Cancer Res.* 2005; 65:6850–7. [PubMed: 16061668]
13. Nakamura H, Mullen JT, Chandrasekhar S, Pawlik TM, Yoon SS, Tanabe KK. Multimodality therapy with a replication-conditional herpes simplex virus 1 mutant that expresses yeast cytosine deaminase for intratumoral conversion of 5-fluorocytosine to 5-fluorouracil. *Cancer Res.* 2001; 61:5447–52. [PubMed: 11454690]
14. Freeman SM, Abboud CN, Whartenby KA, et al. The "bystander effect": tumor regression when a fraction of the tumor mass is genetically modified. *Cancer Res.* 1993; 53:5274–83. [PubMed: 8221662]
15. McCart JA, Puhlmann M, Lee J, et al. Complex interactions between the replicating oncolytic effect and the enzyme/prodrug effect of vaccinia-mediated tumor regression. *Gene Ther.* 2000; 7:1217–23. [PubMed: 10918490]
16. Aghi M, Rabkin S, Martuza RL. Effect of chemotherapy-induced DNA repair on oncolytic herpes simplex viral replication. *J Natl Cancer Inst.* 2006; 98:38–50. [PubMed: 16391370]
17. Carroll NM, Chiocca EA, Tanabe KK. The effect of ganciclovir on herpes simplex virus-mediated oncolysis. *J Surg Res.* 1997; 69:413–7. [PubMed: 9224416]
18. Simpson GR, Han Z, Liu B, Wang Y, Campbell G, Coffin RS. Combination of a fusogenic glycoprotein, prodrug activation, and oncolytic herpes simplex virus for enhanced local tumor control. *Cancer Res.* 2006; 66:4835–42. [PubMed: 16651439]
19. Guffey MB, Parker JN, Lockett WS Jr, et al. Engineered herpes simplex virus expressing bacterial cytosine deaminase for experimental therapy of brain tumors. *Cancer Gene Ther.* 2007; 14:45–56. [PubMed: 16990846]
20. Foloppe J, Kintz J, Futin N, et al. Targeted delivery of a suicide gene to human colorectal tumors by a conditionally replicating vaccinia virus. *Gene Ther.* 2008; 15:1361–71. [PubMed: 18480846]
21. Kuroda T, Martuza RL, Todo T, Rabkin SD. Flip-Flop HSV-BAC: bacterial artificial chromosome based system for rapid generation of recombinant herpes simplex virus vectors using two independent site-specific recombinases. *BMC Biotechnol.* 2006; 6:40. [PubMed: 16995942]
22. Erbs P, Regulier E, Kintz J, et al. In vivo cancer gene therapy by adenovirus-mediated transfer of a bifunctional yeast cytosine deaminase/uracil phosphoribosyltransferase fusion gene. *Cancer Res.* 2000; 60:3813–22. [PubMed: 10919655]

23. Koyama F, Sawada H, Hirao T, Fujii H, Hamada H, Nakano H. Combined suicide gene therapy for human colon cancer cells using adenovirus-mediated transfer of escherichia coli cytosine deaminase gene and Escherichia coli uracil phosphoribosyltransferase gene with 5-fluorocytosine. *Cancer Gene Ther.* 2000; 7:1015–22. [PubMed: 10917204]
24. Chung-Faye GA, Chen MJ, Green NK, et al. In vivo gene therapy for colon cancer using adenovirus-mediated, transfer of the fusion gene cytosine deaminase and uracil phosphoribosyltransferase. *Gene Ther.* 2001; 8:1547–54. [PubMed: 11704815]
25. Kuroda T, Rabkin SD, Martuza RL. Effective treatment of tumors with strong beta-catenin/T-cell factor activity by transcriptionally targeted oncolytic herpes simplex virus vector. *Cancer Res.* 2006; 66:10127–35. [PubMed: 17047077]
26. Pisano R, Breda M, Grassi S, James CA. Hydrophilic interaction liquid chromatography-APCI-mass spectrometry determination of 5-fluorouracil in plasma and tissues. *J Pharm Biomed Anal.* 2005; 38:738–45. [PubMed: 15967302]
27. Sabatini L, Barbieri A, Tosi M, Violante FS. A new high-performance liquid chromatographic/electrospray ionization tandem mass spectrometric method for the simultaneous determination of cyclophosphamide, methotrexate and 5-fluorouracil as markers of surface contamination for occupational exposure monitoring. *J Mass Spectrom.* 2005; 40:669–74. [PubMed: 15739158]
28. Shah VP, Midha KK, Findlay JW, et al. Bioanalytical method validation--a revisit with a decade of progress. *Pharm Res.* 2000; 17:1551–7. [PubMed: 11303967]
29. Pawlik TM, Nakamura H, Yoon SS, et al. Oncolysis of diffuse hepatocellular carcinoma by intravascular administration of a replication-competent, genetically engineered herpesvirus. *Cancer Res.* 2000; 60:2790–5. [PubMed: 10850415]
30. Chahlavi A, Todo T, Martuza RL, Rabkin SD. Replication-competent herpes simplex virus vector G207 and cisplatin combination therapy for head and neck squamous cell carcinoma. *Neoplasia.* 1999; 1:162–9. [PubMed: 10933051]
31. Toyoizumi T, Mick R, Abbas AE, Kang EH, Kaiser LR, Molnar-Kimber KL. Combined therapy with chemotherapeutic agents and herpes simplex virus type 1 ICP34. 5 mutant (HSV-1716) in human non-small cell lung cancer. *Hum Gene Ther.* 1999; 10:3013–29. [PubMed: 10609661]
32. Post DE, Fulci G, Chiocca EA, Van Meir EG. Replicative oncolytic herpes simplex viruses in combination cancer therapies. *Curr Gene Ther.* 2004; 4:41–51. [PubMed: 15032613]
33. Nakano K, Todo T, Zhao G, et al. Enhanced efficacy of conditionally replicating herpes simplex virus (G207) combined with 5-fluorouracil and surgical resection in peritoneal cancer dissemination models. *J Gene Med.* 2005; 7:638–48. [PubMed: 15754306]
34. Gutermann A, Mayer E, von Dehn-Rothfelser K, et al. Efficacy of oncolytic herpesvirus NV1020 can be enhanced by combination with chemotherapeutics in colon carcinoma cells. *Hum Gene Ther.* 2006; 17:1241–53. [PubMed: 17117895]
35. Petrowsky H, Roberts GD, Kooby DA, et al. Functional interaction between fluorodeoxyuridine-induced cellular alterations and replication of a ribonucleotide reductase-negative herpes simplex virus. *J Virol.* 2001; 75:7050–8. [PubMed: 11435585]
36. Eisenberg DP, Adusumilli PS, Hendershott KJ, et al. 5-fluorouracil and gemcitabine potentiate the efficacy of oncolytic herpes viral gene therapy in the treatment of pancreatic cancer. *J Gastrointest Surg.* 2005; 9:1068–77. discussion 77–9. [PubMed: 16269377]
37. Adusumilli PS, Chan MK, Chun YS, et al. Cisplatin-induced GADD34 upregulation potentiates oncolytic viral therapy in the treatment of malignant pleural mesothelioma. *Cancer Biol Ther.* 2006; 5:48–53. [PubMed: 16294031]
38. Weissleder R, Moore A, Mahmood U, et al. In vivo magnetic resonance imaging of transgene expression. *Nature Med.* 2000; 6:351–5. [PubMed: 10700241]
39. Jacobs A, Tjuvajev JG, Dubrovin M, et al. Positron emission tomography-based imaging of transgene expression mediated by replication-conditional, oncolytic herpes simplex virus type 1 mutant vectors in vivo. *Cancer Res.* 2001; 61:2983–95. [PubMed: 11306477]
40. Kuruppu D, Brownell AL, Zhu A, et al. Positron emission tomography of herpes simplex virus 1 oncolysis. *Cancer Res.* 2007; 67:3295–300. [PubMed: 17409438]
41. Gade TP, Koutcher JA, Spees WM, et al. Imaging transgene activity in vivo. *Cancer Res.* 2008; 68:2878–84. [PubMed: 18413756]

42. Harvey BG, Maroni J, O'Donoghue KA, et al. Safety of local delivery of low- and intermediate-dose adenovirus gene transfer vectors to individuals with a spectrum of morbid conditions. *Hum Gene Ther.* 2002; 13:15–63. [PubMed: 11779412]
43. Conrad C, Miller CR, Ji Y, et al. Delta24-hyCD adenovirus suppresses glioma growth in vivo by combining oncolysis and chemosensitization. *Cancer Gene Ther.* 2005; 12:284–94. [PubMed: 15650766]
44. Chang DY, Yoo SW, Hong Y, et al. The growth of brain tumors can be suppressed by multiple transplantation of mesenchymal stem cells expressing cytosine deaminase. *Int J Cancer.* 2010
45. Dias JD, Liikanen I, Guse K, et al. Targeted chemotherapy for head and neck cancer with a chimeric oncolytic adenovirus coding for bifunctional suicide protein FCU1. *Clin Cancer Res.* 2010; 16:2540–9. [PubMed: 20388844]




**Figure 1. Construction of an armed HSV-1 expressing the yCD-UPRT gene**

A) Schematic diagram of HSV-1 expressing yCD-UPRT gene. RL: repeat long, RS: repeat short, UL: unique long, US: unique short, Fcum: Codon-optimized myc-tagged yCD-UPRT, Nter: N-terminal, LacZ: B-galactosidase, SV40: Simian virus 40, pA: Polyadenylation, BGH pA: Bovine Growth Hormone Polyadenylation, CMV-IE: Cytomegalovirus-immediate early


B) HT 29 cells were infected with bMP6-CMV FcumB virus at MOI = 2 for 20 h, and western blot was used to detect the expression of yCD-UPRT by the myc-tag. Lane #1 and lane #2 represent two different viral isolates. ICP4: infected cell protein 4, which is expressed by both bMP6-CMV FcumB and bMP6-empty.






**Figure 2. 5-FC administration enhances the efficacy of the armed oncolytic virus, bMP6-CMVFcumB**

**A)** HT-29 and **B)** MC26 cells were infected with bMP6-CMVFcumB at a MOI of 0.01 or 0.185, respectively, in the presence or absence of increasing concentrations of 5-FC and cytotoxicity was determined by MTT assay. **C)** HT-29 and **D)** MC26 cells were infected with bMP6-CMVFcumB at increasing MOIs in the presence or absence of 5-FC at a concentration of 50μg/ml and cytotoxicity was determined by MTT assay. **E)** HT-29 and **F)** MC26 cells were infected with the control virus bMP6-empty at increasing MOIs in the presence or absence of 5-FC at a concentration of 50μg/ml and cytotoxicity was determined by MTT assay.




**Figure 3. Early 5-FC initiation after viral infection is more effective than late administration**  
**A)** HT-29 cells and **B)** MC26 cells were infected with BMP6-CMVFcumB at increasing MOIs and cytotoxicity was determined by MTT assay after 5-FC at a concentration of 50  $\mu\text{g/ml}$  was initiated at several time points after viral infection.



**Figure 4. Early 5-FC initiation after viral infection and 5-FC administration for 12 days is the most effective dosing schedule in a mouse model**


**A)** MC26 flank tumors were treated by intratumoral injections of bMP6-CMVFcumB in combination with intraperitoneal injections of 5-FC at different initiation times over 6 days.

**B)** MC26 flank tumors were treated by intratumoral injections of bMP6-CMVFcumB in combination with intraperitoneal injections of 5-FC for different durations times over 18 days.



**Figure 5. Viral replication and transgene expression are detected up to Day 15 after viral infection**

MC26 flank tumors were treated with intratumoral injections of bMP6-CMVFcumB, and tumors were harvested at the indicated time points and analyzed. **A)** Viral genomic DNA copy number was assessed by quantitative real-time PCR. **B)** Viral titers were assessed by plaque assay. **C)** Transgene expression was assessed by LacZ staining (Magnification  $\times 200$ ).



**Figure 6. Mouse Pharmacokinetic Studies**

Concentrations of 5-FC and 5-FU in tumor tissue and plasma samples obtained from mice after receiving 15 mg of 5-FC by intraperitoneal injection once a day for 3, 6, 12 or 18 days, beginning 3 days after intratumoral injection of the cytosine deaminase expressing virus. Samples were obtained from groups of 5 mice at each sample time. Concentrations determined in samples from individual animals are indicated as follows: (○) 5-FC in plasma, (●) 5-FC in tumor, (■) 5-FU in tumor, (□) 5-FU in plasma. Symbols drawn through the  $x$ -axis denote samples with concentrations below the 10 ng/ml lower limit of quantitation. Horizontal bars represent mean concentrations.