

SPSS (v.29) were used to perform independent sample t-tests to test for effects and Hierarchical block-wise regressions were done to check for confounding variables.

Results: Lifetime cannabis use was associated with lower global functioning ($p=0.000$), increased risk of suicidal ideation ($p=0.007$), more suicide attempts ($p=0.049$), more self-reported symptoms ($p=0.001$), more school drop-out ($p=0.000$) and with psychotic features ($p=0.024$). Even after testing for known confounders such as female sex and age LUC explained significant variance.

Conclusions: LUC is associated with increased functional and clinical characteristics. The findings are discussed in relationship with clinical practice and limitations of the study.

Disclosure: No significant relationships.

Keywords: Cannabis; out-patient; ADHD

00016

Children and adolescents exposed to maltreatment already exhibit epigenetic patterns suggestive of heightened low-grade inflammation

H. Palma-Gudiel¹, L. Marques Feixa^{1,2*}, S. Romero^{1,3}, M. Rapado-Castro^{1,4}, H. Blasco-Fontecilla^{1,5}, I. Zorrilla^{1,6}, M. Martín⁷, Á. Castro Quintas^{1,2}, J.L. Monteserin-García^{1,2}, E. Font^{1,3}, M. Ramirez^{1,8}, D. Moreno^{1,4}, M. Marín-Vila^{1,5}, N. Moreno^{1,2}, E. Binder⁹ and L. Fañanas^{1,2}

¹Centre for Biomedical Research Network on Mental Health (CIBERSAM), Instituto De Salud Carlos III, Madrid, Spain; ²University of Barcelona, Evolutionary Biology, Ecology And Environmental Sciences, Barcelona, Spain; ³Hospital Clínic, Department Of Child And Adolescent Psychiatry And Psychology, Barcelona, Spain; ⁴Hospital General Universitario Gregorio Marañón, Department Of Child And Adolescent Psychiatry, Madrid, Spain; ⁵Hospital Puerta de Hierro, Department Of Child And Adolescent Psychiatry, Majadahona, Spain; ⁶Hospital Santiago Apostol, Department Of Psychiatry, Vitoria-Gasteiz, Spain; ⁷Hospital Benito Menni, Unitat De Crisis D'adolescents, Sant Boi, Spain; ⁸Galdakao Mental Health Services, Child And Adolescent Mental Health, Galdakao, Spain and ⁹Max-Planck Institute of Psychiatry, Translational Research In Psychiatry, Munich, Germany
*Corresponding author.

doi: 10.1192/j.eurpsy.2022.223

Introduction: Childhood maltreatment (CM) is one of the best described environmental risk factors for developing any psychiatric disorder, while it also confers increased odds for obesity, cardiometabolic disorders and all-cause mortality. Inflammation has been suggested to mediate the widespread clinical effects of CM. Previously, Lighthart et al. (2016) identified a polyepigenetic signature of circulating CRP levels, a measure of chronic low-grade inflammation, that has been reliably associated with a wide array of complex disorders. The study of this biomarker could elucidate the mechanistic relationship between CM and psychiatric outcomes.

Objectives: Thus, CRP-associated epigenetic modifications were explored regarding proximal exposure to CM.

Methods: Genomic DNA was extracted from peripheral blood mononuclear cells of 157 children and adolescents (7 to 17 years old). Exposure to CM was assessed following the TASSCV criteria. Genome-wide DNA methylation was assessed by means of the

EPIC array. Fifty-two out of the 58 original CRP-associated CpG sites surpassed quality control and were included in the analysis. Age, sex, psychopathological status and cell type proportions were included as covariates.

Results: DNA methylation at 12 out of 52 CpG sites (23%) was significantly associated with exposure to CM ($p < .05$); 8 of these associations survived correction for multiple testing ($q < .05$).

Conclusions: This is the first study to date to explore the relationship between childhood maltreatment and an epigenetic signature of chronic low-grade inflammation. Our findings underscore the presence of immune dysregulation early after exposure to CM; further studies are needed to assess the long-term clinical implications of this signature in psychiatric patients.

Disclosure: No significant relationships.

Keywords: childhood maltreatment; chronic low-grade inflammation; epigenetics; DNA methylation

Psychotherapy

00017

Recovery of healthy sexuality in patients with Anorexia Nervosa treated with Enhanced Cognitive Behaviour Therapy (CBT-E): results from a two-year follow-up study highlighting the role of avoidant attachment style

E. Cassioli¹, E. Rossi¹, C. Vizzotto^{1*}, V. Malinconci¹, L. Vignozzi², V. Ricca¹ and G. Castellini¹

¹University of Florence, Psychiatry Unit, Department Of Health Sciences, Florence, Italy and ²University of Florence, Sexual Medicine And Andrology Unit, Department Of Experimental, Clinical, And Biomedical Sciences "mario Serio", Florence, Italy

*Corresponding author.


doi: 10.1192/j.eurpsy.2022.224

Introduction: There is a known association between the core psychopathological features of anorexia nervosa (AN) and sexual dysfunctions, to the point that the recovery of healthy sexuality could be considered a marker of recovery. However, no studies have evaluated the role of insecure attachment in moderating this recovery during treatment.

Objectives: To evaluate the role of insecure attachment as a possible moderator of the recovery of healthy sexuality in patients with AN treated with Enhanced Cognitive Behaviour Therapy (CBT-E).

Methods: A total of 65 patients with anorexia nervosa were treated with CBT-E in a multidisciplinary environment, after filling out self-administered questionnaires for the evaluation of general (SCL-90-R) and ED-specific psychopathology (EDE-Q), female sexuality (FSFI) and adult attachment style (ECR). The assessment was repeated after one (T1) and two years (T2).

Results: At baseline, all domains of sexual dysfunction were significantly predicted by avoidant attachment. A significant amelioration of both general and eating disorder-specific psychopathology and sexual dysfunctions was observed at all follow-up evaluations with respect to baseline levels. However, only 45% of remitted patients also showed a complete recovery of healthy sexuality: this subgroup reported significantly lower avoidance scores when compared to patients who only recovered from AN. Moderation analysis indicated that sexual desire did not increase in participants with higher levels of avoidant attachment.



Conclusions: This study highlighted the crucial role of avoidant attachment in the relationship between AN and sexual dysfunctions, underlining the importance of assessing adult attachment for a better characterization and treatment. Attachment-focused interventions may be beneficial for a full recovery.

Disclosure: No significant relationships.

Keywords: Cognitive Behaviour Therapy; Anorexia nervosa; female sexuality; adult attachment style

O0018

Mental health treatments in an Italian prison: the Parma integrated approach.

L. Pelizza*, D. Maestri, G. Paulillo and P. Pellegrini

AUSL di Parma, Department Of Mental Health, Parma, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.225

Introduction: Mental health interventions for Italian (and European) prisoners with mental disorders remain a problematic issue, despite radical changes in general psychiatric care and a 2008 major government reform transferring mental health care in prison to the National Health Service. Indeed, according to the American Psychological Association, 64% of incarcerated individuals report mental health concerns.

Objectives: The aim of this study is to describe the mental health intervention model implemented since January 2020 for prisoners allocated in the Parma Penitentiary Institutes (PPI). This approach is specifically based on specialized, “person-centered” and “person-tailored” therapeutic-rehabilitation plans in line with psychiatric treatments usually provided in community mental health-care centers of the Parma Department of Mental Health.

Methods: All the processes and procedures included in the PPI intervention model were first carefully described, paying special attention to the service for newly admitted prisoners and each typology of specialized therapeutic-rehabilitation treatment potentially provided. Additionally, a preliminary descriptive process analysis of the first six months of clinical activity was also performed.

Results: Since January 2020, 178 individuals entered the PPI service for newly admitted prisoners. In total, 83 (46.7%) of them were engaged in the services of the PPI mental health-care team (35 with pathological addiction and 48 with mental disorders):

56 prisoners were offered an integrated mental health intervention and 27 exclusively an individual psychological or psychiatric treatment.

Conclusions: The results support the potential applicability of an integrated mental health intervention in prison, planning a person-tailored rehabilitation in close collaboration with the prisoners, their families and the local mental health/social services.

Disclosure: No significant relationships.

Keywords: Psychiatric services; rehabilitation; Prison; mental health care

O0019

The effect of childhood trauma and trauma-focused psychotherapy on blood expression of MED22 in patients with major depressive disorder

A. Minelli^{1*}, E. Maffioletti², R. Carvalho Silva², A. Cattaneo³, G. Perusi⁴, M. Bortolomasi⁴ and M. Gennarelli²

¹University of Brescia, Department Of Molecular And Translational Medicine, Brescia, Italy; ²University of Brescia, Department Of Molecular And Translational Medicine, Verona, Italy; ³IRCCS Fatebenefratelli San Giovanni di Dio, Biological Psychiatry Unit, Brescia, Italy and ⁴Psychiatric Hospital, Villa Santa Chiara, Verona, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.226

Introduction: The only available genome-wide study (Minelli et al., 2018) indicated an association between the neglect CT and MED22, a transcriptional factor gene.

Objectives: To verify how the dysregulation of MED22 could be affected by environmental and genetic factors, we carried out an analysis on these components and a longitudinal study concerning the effect of trauma-focused psychotherapy in MDD patients that experienced CT.

Methods: On a large mRNA sequencing dataset including 368 MDD patients we computed the genetic (GReX) and the environmental (EReX) components affecting gene expression in relation to CT. Furthermore, we measured the expression of MED22 in 22 MDD patients treated with trauma-focused psychotherapy.

Results: The dissection of MED22 expression profiles revealed an association of neglect with environmental and genetic components ($p=6 \times 10^{-3}$ $p=2.6 \times 10^{-4}$). Furthermore, in an independent cohort of 177 controls, we also observed a significant association between cis-eSNPs of MED22 and higher neuroticism scores (best p-value: 0.00848) that are usually associated with a decreased amount of resilience to stress events. Finally, the results of psychotherapy revealed a reduction of depressive symptomatology ($p < 0.001$) and 73% of patients resulted responders at the follow-up visit. MED22 expression during psychotherapy showed a change trend ($p=0.057$) with an interaction effect with response ($p=0.035$). Responder and non-responder patients showed MED22 expression differences at different trauma-focused psychotherapy timepoints ($p=0.15$; $p=0.012$) and at the follow-up ($p=0.021$).

Conclusions: Our results provide insights suggesting that some biological and clinical consequences of CT depend on genetic background and environmental factors that could induce vulnerability or resilience to stressful life events.

Disclosure: No significant relationships.

Keywords: major depressive disorder; Childhood Trauma; Psychotherapy; Blood Expression