

2,4-Dinitro-1-naphthol

Abdul Rauf Raza,^a Aeysha Sultan,^a M. Nawaz Tahir^{b*} and Amra Rubab^a

^aDepartment of Chemistry, University of Sargodha, Sargodha, Pakistan, and

^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 17 October 2010; accepted 3 November 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.042; wR factor = 0.119; data-to-parameter ratio = 10.9.

In the title compound, $\text{C}_{10}\text{H}_6\text{N}_2\text{O}_5$, the two fused rings are almost co-planar, with an r.m.s. deviation of 0.0163 \AA . The nitro groups are oriented at dihedral angles of $2.62(11)$ and $44.69(11)^\circ$ with respect to the plane of the parent fused rings. Intramolecular $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds complete $S(6)$ ring motifs. In the crystal, molecules are linked into chains along [101] by intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. $\pi-\pi$ interactions [centroid–centroid distances = $3.6296(15)$, $3.8104(15)$ and $3.6513(14)\text{ \AA}$] might play a role in stabilizing the structure.

Related literature

For background to estrogens, see: Schwartz *et al.* (1995); O'Donnell *et al.* (2001). For related structures, see: Filipenko *et al.* (2001); Rozycka-Sokolowska *et al.* (2004). For graph-set notation, see: Bernstein *et al.* (1995). For $\pi-\pi$ interactions, see: Janiak (2000).

**Experimental***Crystal data*

$\text{C}_{10}\text{H}_6\text{N}_2\text{O}_5$	$V = 944.4(2)\text{ \AA}^3$
$M_r = 234.17$	$Z = 4$
Monoclinic, $P2_1/n$	$\text{Mo K}\alpha$ radiation
$a = 7.0512(10)\text{ \AA}$	$\mu = 0.14\text{ mm}^{-1}$
$b = 16.3541(19)\text{ \AA}$	$T = 296\text{ K}$
$c = 8.7988(10)\text{ \AA}$	$0.32 \times 0.14 \times 0.12\text{ mm}$
$\beta = 111.452(6)^\circ$	

Data collection

Bruker Kappa APEXII CCD diffractometer	6868 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	1684 independent reflections
$T_{\min} = 0.978$, $T_{\max} = 0.982$	1058 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.052$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$	155 parameters
$wR(F^2) = 0.119$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$
1684 reflections	$\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O5—H5A \cdots O1 ⁱ	0.82	2.53	3.006 (3)	118
O5—H5A \cdots O4	0.82	1.87	2.573 (2)	142
O5—H5A \cdots N2	0.82	2.47	2.892 (3)	113
C5—H5 \cdots O1	0.93	2.35	2.902 (3)	118

Symmetry code: (i) $x + 1, y, z + 1$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the provision of funds for the purchase of the diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan. ARR also acknowledges the Higher Education Commission, Government of Pakistan, for generous support of the research project (20-819).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2615).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Filipenko, O. S., Chuev, I. I., Leonova, L. S., Shilov, G. V. & Aldoshin, S. M. (2001). *Dokl. Phys. Chem.* **376**, 27–30.
- Janiak, C. (2000). *J. Chem. Soc. Dalton Trans.* pp. 3885–3896.
- O'Donnell, L., Kristen, M. R., Margaret, E. J. & Evan, R. S. (2001). *Endocr. Rev.* **22**, 289–318.
- Rozycka-Sokolowska, E., Marciniak, B. & Pavlyuk, V. (2004). *Acta Cryst. E60*, o884–o885.
- Schwartz, J., Freeman, R. & Frishman, W. (1995). *J. Clin. Pharmacol.* **35**, 1–16.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o3101 [doi:10.1107/S1600536810045034]

2,4-Dinitro-1-naphthol

A. R. Raza, A. Sultan, M. N. Tahir and A. Rubab

Comment

Estrogens have been found to have maintenance effects on bone, brain, skin, cardiovascular system (Schwartz *et al.*, 1995). In spite of their beneficial biological effects, estrogens are suspected to have relationship with the risk of cancer and thromboembolic diseases (O'Donnell *et al.*, 2001). The title compound was obtained as an interesting side product while synthesizing substituted 1-tetralone as AB ring of the estrogen skeleton which will bear substituent at positions not found in nature.

The title compound (**I**) is related to the published crystal structures of 1-naphthalenol (Rozicka-Sokolowska *et al.*, 2004) and hydroxonium 2,4-dinitro-1-hydroxy-7-sulfonatonaphthalene monohydrate (Filipenko *et al.*, 2001).

In (**I**), the two fused rings (C1—C10) are nearly planar with the largest deviation being 0.033 (2) Å at C10 (Fig. 1). The hydroxyl O5 atom is only deviating by 0.129 (2) Å from the mean plane. Owing to the intramolecular O5—H···O4 hydrogen bond (Table 1), the O3 and O4 atoms are only slightly displaced from the mean plane of the fused rings by -0.074 (2) Å and 0.023 (2) Å, respectively. The other nitro group is twisted by 44.72 (12)° with respect to the fused rings.

Strong intramolecular H-bondings of O—H···O and C—H···O types (Table 1, Fig. 1) complete S(6) ring motifs (Bernstein *et al.*, 1995). The molecules are stabilized in the form of polymeric chains due to intermolecular O—H···O hydrogen bonds (Table 1, Fig. 2). Slippest weak π—π interactions (Table 2) might play a role in stabilizing the packing.

Experimental

The 1-tetralone (1 ml, 1.1 g, 1 eq) was added to a chilled and well stirred nitrating mixture containing H₂SO₄ (2 ml, 0.57 g, 2 eq) and HNO₃ (0.5 ml, 0.7 g, 1.5 eq). The reaction mixture was then neutralized and extracted with EtOAc (3 × 25 ml) and the combined organic extract was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography and the title compound (**I**) was obtained as dull brown crystalline solid in 61–79th fraction (20 ml each) using 2.5% CHCl₃ (3 × 500 ml) as mobile phase. Yield: 15%.

Refinement

The H-atoms were positioned geometrically with (O—H = 0.82, C—H = 0.93 Å) and treated as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{O})$.

supplementary materials

Figures




Fig. 1. View of the title compound with the atom numbering scheme. The thermal ellipsoids are drawn at the 50% probability level. H-atoms are shown by small circles of arbitrary radii. The dotted line indicate the intramolecular H-bond.




Fig. 2. The partial packing (*PLATON*; Spek, 2009) which shows that molecules form polymeric chains.

2,4-Dinitro-1-naphthol

Crystal data

C ₁₀ H ₆ N ₂ O ₅	F(000) = 480
M _r = 234.17	D _x = 1.647 Mg m ⁻³
Monoclinic, P2 ₁ /n	Mo K α radiation, λ = 0.71073 Å
Hall symbol: -P 2yn	Cell parameters from 1058 reflections
a = 7.0512 (10) Å	θ = 2.5–25.1°
b = 16.3541 (19) Å	μ = 0.14 mm ⁻¹
c = 8.7988 (10) Å	T = 296 K
β = 111.452 (6)°	Needle, brown
V = 944.4 (2) Å ³	0.32 × 0.14 × 0.12 mm
Z = 4	

Data collection

Bruker Kappa APEXII CCD diffractometer	1684 independent reflections
Radiation source: fine-focus sealed tube graphite	1058 reflections with $I > 2\sigma(I)$
Detector resolution: 8.20 pixels mm ⁻¹	R_{int} = 0.052
ω scans	$\theta_{\text{max}} = 25.1^\circ$, $\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -7 \rightarrow 8$
$T_{\text{min}} = 0.978$, $T_{\text{max}} = 0.982$	$k = -19 \rightarrow 19$
6868 measured reflections	$l = -10 \rightarrow 10$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.042$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.119$	H-atom parameters constrained
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.059P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
1684 reflections	$(\Delta/\sigma)_{\max} < 0.001$
155 parameters	$\Delta\rho_{\max} = 0.18 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.3884 (3)	0.45828 (13)	0.10925 (19)	0.0656 (6)
O2	0.6120 (3)	0.36519 (11)	0.1294 (2)	0.0663 (6)
O3	0.8511 (3)	0.26242 (11)	0.6881 (2)	0.0762 (7)
O4	1.0038 (3)	0.34485 (11)	0.8845 (2)	0.0735 (6)
O5	0.9984 (3)	0.49989 (10)	0.83341 (16)	0.0500 (5)
H5A	1.0356	0.4587	0.8896	0.060*
N1	0.5444 (3)	0.42063 (12)	0.1860 (2)	0.0441 (5)
N2	0.8995 (3)	0.33110 (13)	0.7401 (3)	0.0497 (6)
C1	0.8052 (3)	0.54453 (12)	0.5693 (2)	0.0313 (5)
C2	0.8467 (4)	0.62585 (14)	0.6216 (3)	0.0415 (6)
H2	0.9197	0.6370	0.7312	0.050*
C3	0.7809 (4)	0.68880 (14)	0.5131 (3)	0.0465 (7)
H3	0.8077	0.7426	0.5489	0.056*
C4	0.6744 (4)	0.67246 (15)	0.3500 (3)	0.0487 (7)
H4	0.6329	0.7157	0.2766	0.058*
C5	0.6283 (4)	0.59409 (15)	0.2936 (3)	0.0419 (6)
H5	0.5557	0.5848	0.1832	0.050*
C6	0.6905 (3)	0.52753 (13)	0.4024 (2)	0.0313 (5)

supplementary materials

C7	0.6553 (3)	0.44389 (13)	0.3569 (2)	0.0334 (5)
C8	0.7224 (3)	0.38182 (13)	0.4646 (2)	0.0373 (6)
H8	0.6948	0.3279	0.4299	0.045*
C9	0.8335 (3)	0.39941 (13)	0.6284 (2)	0.0354 (6)
C10	0.8816 (3)	0.47868 (13)	0.6824 (2)	0.0345 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0468 (12)	0.0892 (14)	0.0412 (10)	0.0043 (11)	-0.0071 (9)	-0.0067 (9)
O2	0.0875 (16)	0.0570 (12)	0.0489 (11)	0.0026 (11)	0.0184 (10)	-0.0192 (9)
O3	0.110 (2)	0.0391 (11)	0.0731 (13)	-0.0007 (11)	0.0256 (12)	0.0117 (9)
O4	0.0881 (16)	0.0695 (13)	0.0441 (11)	-0.0059 (11)	0.0017 (11)	0.0225 (9)
O5	0.0560 (12)	0.0565 (10)	0.0260 (9)	-0.0001 (9)	0.0012 (8)	0.0046 (7)
N1	0.0459 (14)	0.0488 (12)	0.0320 (11)	-0.0085 (11)	0.0075 (10)	-0.0052 (9)
N2	0.0521 (15)	0.0483 (14)	0.0490 (13)	0.0019 (11)	0.0190 (11)	0.0158 (10)
C1	0.0258 (13)	0.0386 (12)	0.0291 (11)	0.0020 (10)	0.0096 (9)	-0.0002 (9)
C2	0.0405 (15)	0.0446 (14)	0.0385 (13)	-0.0023 (11)	0.0132 (11)	-0.0055 (11)
C3	0.0493 (17)	0.0376 (14)	0.0519 (15)	0.0043 (12)	0.0175 (13)	0.0013 (11)
C4	0.0490 (17)	0.0448 (15)	0.0489 (15)	0.0094 (12)	0.0139 (13)	0.0142 (12)
C5	0.0368 (15)	0.0526 (15)	0.0317 (12)	0.0054 (12)	0.0070 (10)	0.0065 (11)
C6	0.0235 (12)	0.0401 (12)	0.0298 (11)	0.0026 (10)	0.0093 (9)	0.0016 (9)
C7	0.0267 (13)	0.0435 (13)	0.0272 (11)	-0.0026 (10)	0.0066 (9)	-0.0030 (10)
C8	0.0347 (14)	0.0384 (13)	0.0388 (13)	-0.0021 (11)	0.0137 (11)	-0.0017 (10)
C9	0.0337 (14)	0.0404 (13)	0.0321 (12)	0.0021 (10)	0.0120 (10)	0.0083 (10)
C10	0.0307 (13)	0.0474 (14)	0.0247 (11)	0.0008 (11)	0.0093 (10)	0.0021 (10)

Geometric parameters (\AA , $^\circ$)

O1—N1	1.225 (2)	C2—H2	0.9300
O2—N1	1.213 (2)	C3—C4	1.381 (3)
O3—N2	1.214 (3)	C3—H3	0.9300
O4—N2	1.234 (3)	C4—C5	1.370 (3)
O5—C10	1.328 (2)	C4—H4	0.9300
O5—H5A	0.8200	C5—C6	1.409 (3)
N1—C7	1.469 (3)	C5—H5	0.9300
N2—C9	1.448 (3)	C6—C7	1.421 (3)
C1—C2	1.403 (3)	C7—C8	1.351 (3)
C1—C6	1.421 (3)	C8—C9	1.395 (3)
C1—C10	1.431 (3)	C8—H8	0.9300
C2—C3	1.365 (3)	C9—C10	1.380 (3)
C10—O5—H5A	109.5	C4—C5—C6	120.2 (2)
O2—N1—O1	123.9 (2)	C4—C5—H5	119.9
O2—N1—C7	118.2 (2)	C6—C5—H5	119.9
O1—N1—C7	117.9 (2)	C5—C6—C7	124.98 (19)
O3—N2—O4	122.4 (2)	C5—C6—C1	118.0 (2)
O3—N2—C9	118.8 (2)	C7—C6—C1	117.00 (19)
O4—N2—C9	118.8 (2)	C8—C7—C6	123.06 (19)

C2—C1—C6	119.75 (19)	C8—C7—N1	116.25 (19)
C2—C1—C10	120.33 (19)	C6—C7—N1	120.68 (18)
C6—C1—C10	119.90 (19)	C7—C8—C9	119.3 (2)
C3—C2—C1	120.6 (2)	C7—C8—H8	120.3
C3—C2—H2	119.7	C9—C8—H8	120.3
C1—C2—H2	119.7	C10—C9—C8	121.60 (19)
C2—C3—C4	119.8 (2)	C10—C9—N2	120.89 (19)
C2—C3—H3	120.1	C8—C9—N2	117.5 (2)
C4—C3—H3	120.1	O5—C10—C9	125.07 (19)
C5—C4—C3	121.6 (2)	O5—C10—C1	115.92 (19)
C5—C4—H4	119.2	C9—C10—C1	119.00 (18)
C3—C4—H4	119.2		
C6—C1—C2—C3	-1.1 (3)	O1—N1—C7—C6	-44.9 (3)
C10—C1—C2—C3	177.5 (2)	C6—C7—C8—C9	-0.7 (3)
C1—C2—C3—C4	-0.7 (4)	N1—C7—C8—C9	178.4 (2)
C2—C3—C4—C5	1.5 (4)	C7—C8—C9—C10	-1.9 (3)
C3—C4—C5—C6	-0.4 (4)	C7—C8—C9—N2	179.0 (2)
C4—C5—C6—C7	-178.5 (2)	O3—N2—C9—C10	178.8 (2)
C4—C5—C6—C1	-1.4 (3)	O4—N2—C9—C10	-1.5 (3)
C2—C1—C6—C5	2.1 (3)	O3—N2—C9—C8	-2.1 (3)
C10—C1—C6—C5	-176.42 (19)	O4—N2—C9—C8	177.6 (2)
C2—C1—C6—C7	179.44 (19)	C8—C9—C10—O5	-175.0 (2)
C10—C1—C6—C7	0.9 (3)	N2—C9—C10—O5	4.1 (3)
C5—C6—C7—C8	178.2 (2)	C8—C9—C10—C1	3.8 (3)
C1—C6—C7—C8	1.1 (3)	N2—C9—C10—C1	-177.11 (19)
C5—C6—C7—N1	-0.8 (3)	C2—C1—C10—O5	-2.9 (3)
C1—C6—C7—N1	-177.9 (2)	C6—C1—C10—O5	175.7 (2)
O2—N1—C7—C8	-43.7 (3)	C2—C1—C10—C9	178.2 (2)
O1—N1—C7—C8	136.1 (2)	C6—C1—C10—C9	-3.2 (3)
O2—N1—C7—C6	135.4 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O5—H5A···O1 ⁱ	0.82	2.53	3.006 (3)	118
O5—H5A···O4	0.82	1.87	2.573 (2)	142
O5—H5A···N2	0.82	2.47	2.892 (3)	113
C5—H5···O1	0.93	2.35	2.902 (3)	118

Symmetry codes: (i) $x+1, y, z+1$.

Table 2
 π – π interactions (Å, °)

Cg1 and Cg2 are the centroids of the C1–C6 and C1/C6–C10 rings, respectively. ipd is the mean interplanar distance (distance from one plane to the neighbouring centroid) and sa is the slippage angle (angle subtended by the intercentroid vector to the plane normal). For details, see Janiak (2000)

	Cg···Cg	ipd	sa
Cg1···Cg2 ⁱ	3.6296 (15)	3.365	1.305
Cg1···Cg2 ⁱⁱ	3.8104 (15)	3.552	1.323

supplementary materials

Cg2...Cg2ⁱ

3.6513 (14)

3.378

1.386

Symmetry codes: (i) 1-x, 1-y, 1-z, (ii) 2-x, 1-y, 1-z.

Fig. 1




Fig. 2

