

POSTER PRESENTATION

Open Access

Canine osteosarcoma cells exhibit resistance to aurora kinase inhibitors

Claire M Cannon^{1*}, John Pozniak^{1,2}, Milcah C Scott^{1,2}, Daisuke Ito^{1,2}, Brandi H Gorden^{1,2}, Ashley J Graef^{1,2}, Jaime F Modiano^{1,2}

From São Paulo Advanced School of Comparative Oncology
Águas de São Pedro, Brazil. 30 September - 6 October 2012

Background

Aurora kinases are overexpressed and associated with poor prognosis in several human tumours. Recently, over-expression and correlation with poor prognosis was found in canine osteosarcoma. We examined the effects of Aurora kinase inhibition in canine osteosarcoma cell lines. This was the first step to explore development of Aurora kinase inhibitors (AKIs) as novel therapeutic agents for canine osteosarcoma.

Materials and methods

We evaluated the effects of two AKIs, AZD1152 and VX680, on four canine osteosarcoma cell lines. Viability was assessed using the MTS assay, apoptosis using caspase activation and flow cytometry for Annexin V binding and 7-AAD uptake, and target modulation using Western blotting. We also evaluated gene and protein expression of Aurora kinase B, ABCB1 and ABG2.

Results

Expression of Aurora kinase B protein and mRNA was seen in all cell lines. Although Aurora kinase inhibitors induced cytotoxicity, half-maximal inhibitory concentrations were significantly higher than seen in other cancer types and induction of apoptosis was minimal at concentrations close to IC50s. AZD1152 reduced Aurora kinase B phosphorylation, indicating effective target modulation. ABCB1 and ABCG2 transporter-mediated efflux is one known mechanism of resistance against these drugs. Verapamil modestly enhanced AZD1152-mediated apoptosis and ABC transporters were expressed by a small percentage of cells,

suggesting that other mechanisms may contribute to resistance.

Conclusions

Canine osteosarcoma cells are resistant to AKIs and we suggest that these compounds are unlikely to be useful as monotherapy for this disease. Further investigation of resistance mechanisms and the potential utility of AKIs in combination therapy is warranted.

Acknowledgements

A University of Minnesota College of Veterinary Medicine Small Companion Animal Grant and an American Kennel Club Canine Health Foundation Acorn Grant (01503-A) funded this research. Claire Cannon is supported by Morris Animal Foundation Grant D08CA-501. Daisuke Ito is supported by Morris Animal Foundation First Award Grant D12CA-302.

Author details

¹Department of Veterinary Clinical Sciences, College of Veterinary Medicine, University of Minnesota, St Paul, MN, USA. ²Masonic Cancer Center, University of Minnesota, MN, USA.

Published: 4 April 2013

doi:10.1186/1753-6561-7-S2-P15

Cite this article as: Cannon et al.: Canine osteosarcoma cells exhibit resistance to aurora kinase inhibitors. *BMC Proceedings* 2013 **7**(Suppl 2): P15.

* Correspondence: cmcannon@umn.edu

¹Department of Veterinary Clinical Sciences, College of Veterinary Medicine, University of Minnesota, St Paul, MN, USA

Full list of author information is available at the end of the article