

POSTER PRESENTATION

Open Access

# The effector protein ExoY secreted by *Pseudomonas aeruginosa* is a nucleotidyl cyclase with preference for GTP

Ulrike Voigt<sup>1\*</sup>, Heike Burhenne<sup>1</sup>, Lena Sonnow<sup>1</sup>, Christine Wölfe<sup>1</sup>, Corinna Spangler<sup>1</sup>, Daniel Ladant<sup>2</sup>, Dara W Frank<sup>3</sup>, Volkhard Kaeber<sup>1</sup>, Roland Seifert<sup>1</sup>

From 5th International Conference on cGMP: Generators, Effectors and Therapeutic Implications  
Halle, Germany. 24-26 June 2011

## Background

*Pseudomonas aeruginosa* is an important opportunistic pathogen causing serious pulmonary, urogenital and systemic infections. *P. aeruginosa* injects four effector proteins into host cells via the type III secretion system. ExoY is one of these proteins.

ExoY was originally classified as adenylyl cyclase with homology to the typical calmodulin-stimulated adenylyl cyclase exotoxins CyaA from *Bordetella pertussis* and edema factor from *Bacillus anthracis*, but the pathophysiological function of ExoY has remained elusive [1,2]. Recently, our group showed that CyaA and edema factor possess a rather broad base specificity (ATP >> CTP > UTP) [3,4], raising the question of whether ExoY may also bind and metabolize nucleoside 5'- triphosphates other than ATP.

## Methods

We determined cyclic nucleotide concentrations in cells transfected with ExoY plasmid or infected with *P. aeruginosa* with a highly sensitive HPLC-MS/MS method. Moreover, we determined the catalytic activity of purified ExoY.

## Results

In mammalian cells transfected with ExoY plasmid and infected with ExoY-encoding *P. aeruginosa*, massive production of cGMP and cUMP was observed, with little production of cAMP. Purified ExoY was a highly effective nucleotidyl cyclase with the substrate preference GTP >> UTP ~ ATP > CTP. Fluorescence resonance energy transfer studies with methylanthraniloyl-substituted nucleotides corroborated the preference of ExoY for GTP. In contrast to ExoY, CyaA induced accumulation

**Table 1 Comparison of ExoY with CyaA and EF**

Parameter	ExoY	CyaA	EF
Bacterial source	<i>Pseudomonas aeruginosa</i>	<i>Bordetella pertussis</i>	<i>Bacillus anthracis</i>
Secretion type	III	I	II
Function as toxin	Unknown, "antitoxin?"	yes	yes
Activation mechanism	Cytosolic cofactor (mammalian cells and <i>D. discoideum</i> )	calmodulin	calmodulin
Substrate-specificity purified enzyme	GTP >> UTP ≥ ATP > CTP	ATP >> CTP > UTP > GTP	ATP >> CTP > UTP > GTP
Substrate-specificity intact cells	UTP ~ GTP > ATP > CTP (except for B103 cells)	ATP >> CTP > UTP >> GTP (ineffective)	ATP >> CTP >> UTP and GTP(ineffective)

\* Correspondence: voigt.ulrike@mh-hannover.de

<sup>1</sup>Institute of Pharmacology, Medical School of Hannover, D-30625 Hannover, Germany

Full list of author information is available at the end of the article

of cyclic nucleotides in the order cAMP > cCMP > cUMP in mammalian cells, and edema factor induced only cAMP- and cCMP accumulation.

## Conclusion

ExoY is a nucleotidyl cyclase with preference for GTP, and the substrate-specificity of ExoY is clearly different from that of CyaA and edema factor. Our data open the door for future studies aiming at the elucidation of the as yet unknown pathophysiological function of ExoY and which role cCMP, cGMP and cUMP play in this process.

### Author details

<sup>1</sup>Institute of Pharmacology, Medical School of Hannover, D-30625 Hannover, Germany. <sup>2</sup>Institut Pasteur, Department of Structural Biology and Chemistry, F-75724 Paris, France. <sup>3</sup>Department of Microbiology and Molecular Genetics, Medical College of Wisconsin, Milwaukee, WI 53326, USA.

Published: 1 August 2011

### References

1. Yahr TL, Vallis AJ, Hancock MK, Barbieri JT, Frank DW: ExoY, an adenylate cyclase secreted by the *Pseudomonas aeruginosa* type III system. *Microbiol* 1998, **95**:13899-13904.
2. Vallis AJ, Finck-Barbançon V, Jahr TL, Frank DW: Biological effects of *Pseudomonas aeruginosa* type III secreted proteins on CHO cells. *Infection and Immunity* 1999, **67**:2040-2044.
3. Göttle M, Dove S, Steindel P, Shen Y, Tang WJ, Geduhn J, König B, Seifert R: Molecular analysis of the interaction of *Bordetella pertussis* adenylyl cyclase with fluorescent nucleotides. *Mol Pharmacol* 2007, **72**:526-535.
4. Taha HM, Schmidt J, Göttle M, Suryanarayana S, Shen Y, Tang WJ, Gille A, Geduhn J, König B, Dove S, Seifert R: Molecular analysis of the interaction of anthrax adenylyl cyclase toxin, edema factor, with 2'(3')-O-(N-methyl) anthraniloyl)- substituted purine and pyrimidine nucleotides. *Mol Pharmacol* 2009, **75**:693-703.

doi:10.1186/1471-2210-11-S1-P74

Cite this article as: Voigt et al.: The effector protein ExoY secreted by *Pseudomonas aeruginosa* is a nucleotidyl cyclase with preference for GTP. *BMC Pharmacology* 2011 **11**(Suppl 1):P74.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

