



Crystal structure of tris(*N*-methylsalicylaldiminato- κ^2 N,O)vanadium(III)

Jessica Hilbert,* Sven Kabus, Christian Näther and Wolfgang Bensch

Institut für Anorganische Chemie, Christian-Albrechts-Universität zu Kiel, Max-Eyth-Strasse 2, 24118 Kiel, Germany. *Correspondence e-mail: jhilbert@ac.uni-kiel.de

Received 11 November 2015; accepted 12 November 2015

Edited by E. R. T. Tiekink, University of Malaya, Malaysia


The structure of the title complex, $[V(C_8H_8NO)_3]$, comprises neutral and discrete complexes, in which the V^{III} cation is coordinated by three anionic *N*-methylsalicylaldimate ligands within a slightly distorted *mer*- N_3O_3 octahedral geometry. In the crystal structure, the molecules are linked via $C-H \cdots O$ hydrogen bonds into supramolecular chains that extend along the *c* axis.

Keywords: crystal structure; vanadium(III); *N*-methylsalicylaldimate.

CCDC reference: 1436532

1. Related literature

For structures of discrete complexes of Mo and V with *N*-methylsalicylaldimate as the ligand, see: Davies & Gatehouse (1974); Cornman *et al.* (1997). For the synthesis of the starting material, see: Bonadies & Carrano (1986).



2. Experimental

2.1. Crystal data

$[V(C_8H_8NO)_3]$	$V = 2174.79 (13) \text{ \AA}^3$
$M_r = 453.40$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 7.7414 (3) \text{ \AA}$	$\mu = 0.49 \text{ mm}^{-1}$
$b = 26.0018 (7) \text{ \AA}$	$T = 170 \text{ K}$
$c = 11.1004 (4) \text{ \AA}$	$0.24 \times 0.14 \times 0.06 \text{ mm}$
$\beta = 103.265 (3)^\circ$	

2.2. Data collection

STOE IPDS-1 diffractometer	18648 measured reflections
Absorption correction: numerical (<i>X-SHAPE</i> and <i>X-RED32</i> ; Stoe, 2008)	4741 independent reflections
$T_{\min} = 0.919$, $T_{\max} = 0.974$	4054 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.030$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$	283 parameters
$wR(F^2) = 0.102$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$
4741 reflections	$\Delta\rho_{\text{min}} = -0.42 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry ($\text{\AA}$, $^\circ$).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$C27-H27 \cdots O21^i$	0.95	2.56	3.431 (2)	153

Symmetry code: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *X-Area* (Stoe, 2008); cell refinement: *X-Area*; data reduction: *X-Area*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008) and *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Acknowledgements

This work was supported by the State of Schleswig-Holstein.

Supporting information for this paper is available from the IUCr electronic archives (Reference: TK5408).

References

- Bonadies, J. A. & Carrano, C. J. (1986). *J. Am. Chem. Soc.* **108**, 4088–4095.
- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Cornman, C. R., Geiser-Bush, K. M., Rowley, S. P. & Boyle, P. D. (1997). *CrystEngComm*, **36**, 6401–6408.
- Davies, J. E. & Gatehouse, B. M. (1974). *J. Chem. Soc. Dalton Trans.* pp. 184–187.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **A71**, 3–8.
- Stoe (2008). *X-Area*, *X-RED32* and *X-SHAPE*. Stoe & Cie, Darmstadt, Germany.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2015). E71, m225 [doi:10.1107/S2056989015021453]

Crystal structure of tris(*N*-methyalsalicylaldiminato- κ^2N,O)vanadium(III)

Jessica Hilbert, Sven Kabus, Christian Näther and Wolfgang Bensch

S1. Synthesis and crystallization

Most of the chemicals are commercially available: Sn (Fluka, 99.9%), S (Alfa Aesar, 99.5%), and methylamine (abcr, 40% aqueous solution). (*N,N'*-Disalicylideneethylenediamine)oxovanadium(IV) was prepared following the procedure of Bonadies & Carrano (1986): (*N,N'*-disalicylideneethylenediamine)oxovanadium (IV) (83.8 mg, 0.25 mmol), Sn (29.7 mg, 0.25 mmol) and S (24.1 mg, 0.75 mmol) were reacted in a glass tube (inner volume 11 mL) with methylamine (1.5 mL) and H₂O (0.5 mL) under solvothermal conditions at 120 °C for 24 h. Afterwards, the solid residue was filtered off, washed with water and ethanol, and dried over silica gel. The product contains red blocks of the title complex and a small amount of brown blocks of bis(*N*-methyalsalicylaldiminato)oxovanadium(IV) (Cornman *et al.* (1997)). Even if Sn and S are not contained in the final product, they are needed for product formation, as otherwise only (*N,N'*-disalicylideneethylenediamine)oxovanadium(IV) is isolated.

S2. Refinement

The C—H H atoms were positioned with idealized geometry (methyl H atoms allowed to rotate but not to tip) and were refined isotropically with $U_{eq}(H) = 1.2 U_{eq}(C)$ (1.5 for methyl H atoms) using a riding model with C—H = 0.95 Å for aromatic H atoms and 0.98 Å for methyl H atoms.




Figure 1

The molecular structure of the title complex with atom labelling. Displacement ellipsoids are drawn at the 50% probability level.




Figure 2

Unit cell contents of the crystal structure of the title complex viewed in projection down the a axis with hydrogen bonds shown as dashed lines. For clarity, all H atoms except those that participates in hydrogen bonding are omitted.

Tris(*N*-methysalicylaldiminato- $\kappa^2 N,O$)vanadium(III)

Crystal data

$[\text{V}(\text{C}_8\text{H}_8\text{NO})_3]$

$M_r = 453.40$

Monoclinic, $P2_1/c$

$a = 7.7414$ (3) Å

$b = 26.0018$ (7) Å

$c = 11.1004$ (4) Å

$\beta = 103.265$ (3)°

$V = 2174.79$ (13) Å³

$Z = 4$

$F(000) = 944$

$D_x = 1.385$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 18648 reflections

$\theta = 1.6\text{--}27.0^\circ$

$\mu = 0.49$ mm⁻¹

$T = 170$ K

Block, red

$0.24 \times 0.14 \times 0.06$ mm

Data collection

STOE IPDS-1

diffractometer

Radiation source: fine-focus sealed tube

φ -scans

Absorption correction: numerical

(*X-SHAPE* and *X-RED32*; Stoe, 2008)

$T_{\min} = 0.919$, $T_{\max} = 0.974$

18648 measured reflections

4741 independent reflections

4054 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 27.0^\circ$, $\theta_{\min} = 1.6^\circ$

$h = -9 \rightarrow 9$

$k = -30 \rightarrow 33$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.102$

$S = 1.07$

4741 reflections

283 parameters

0 restraints

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0523P)^2 + 0.6659P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.30 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.42 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
V1	0.60947 (4)	0.34777 (2)	0.43424 (3)	0.03210 (10)
O1	0.58715 (17)	0.39024 (5)	0.28969 (12)	0.0400 (3)
C1	0.6728 (2)	0.39169 (7)	0.19994 (17)	0.0375 (4)
C2	0.8151 (3)	0.35824 (8)	0.19485 (18)	0.0390 (4)
C3	0.8995 (3)	0.36297 (9)	0.0951 (2)	0.0497 (5)
H3	0.9964	0.3410	0.0919	0.060*
C4	0.8448 (3)	0.39846 (10)	0.0029 (2)	0.0578 (6)
H4	0.9030	0.4010	−0.0634	0.069*
C5	0.7033 (3)	0.43068 (10)	0.0075 (2)	0.0550 (6)
H5	0.6646	0.4552	−0.0564	0.066*
C6	0.6186 (3)	0.42752 (9)	0.10360 (19)	0.0466 (5)
H6	0.5221	0.4499	0.1049	0.056*
C7	0.8789 (2)	0.31882 (8)	0.28513 (17)	0.0384 (4)
H7	0.9766	0.2989	0.2733	0.046*
N1	0.81716 (19)	0.30777 (6)	0.38049 (14)	0.0351 (3)
C8	0.9074 (3)	0.26689 (8)	0.46168 (18)	0.0423 (4)
H8A	0.9545	0.2808	0.5448	0.064*
H8B	1.0053	0.2533	0.4287	0.064*
H8C	0.8231	0.2392	0.4659	0.064*
O11	0.78967 (17)	0.38679 (5)	0.54398 (12)	0.0397 (3)
C11	0.7839 (3)	0.42940 (8)	0.60833 (18)	0.0402 (4)
C12	0.6228 (3)	0.45333 (8)	0.61828 (18)	0.0416 (4)
C13	0.6290 (3)	0.49804 (9)	0.6899 (2)	0.0510 (5)
H13	0.5211	0.5143	0.6952	0.061*
C14	0.7869 (4)	0.51892 (9)	0.7524 (2)	0.0580 (6)
H14	0.7882	0.5490	0.8012	0.070*
C15	0.9450 (4)	0.49555 (10)	0.7434 (2)	0.0587 (6)
H15	1.0549	0.5099	0.7863	0.070*
C16	0.9439 (3)	0.45159 (9)	0.6727 (2)	0.0501 (5)
H16	1.0533	0.4362	0.6676	0.060*
C17	0.4494 (3)	0.43340 (8)	0.56016 (18)	0.0430 (4)
H17	0.3499	0.4519	0.5740	0.052*
N11	0.4164 (2)	0.39324 (6)	0.49186 (15)	0.0378 (3)
C18	0.2295 (2)	0.37888 (9)	0.4467 (2)	0.0494 (5)

H18A	0.2020	0.3762	0.3562	0.074*
H18B	0.1538	0.4051	0.4715	0.074*
H18C	0.2079	0.3456	0.4822	0.074*
O21	0.42824 (16)	0.30065 (5)	0.34205 (11)	0.0370 (3)
C21	0.3734 (2)	0.25415 (7)	0.35945 (16)	0.0328 (4)
C22	0.4208 (2)	0.22876 (7)	0.47477 (17)	0.0352 (4)
C23	0.3555 (3)	0.17928 (8)	0.4881 (2)	0.0446 (5)
H23	0.3849	0.1630	0.5668	0.054*
C24	0.2496 (3)	0.15367 (9)	0.3894 (2)	0.0499 (5)
H24	0.2098	0.1197	0.3990	0.060*
C25	0.2023 (3)	0.17858 (8)	0.2756 (2)	0.0439 (5)
H25	0.1293	0.1613	0.2070	0.053*
C26	0.2596 (2)	0.22787 (8)	0.26080 (18)	0.0376 (4)
H26	0.2219	0.2445	0.1830	0.045*
C27	0.5263 (2)	0.25346 (7)	0.58385 (16)	0.0349 (4)
H27	0.5412	0.2352	0.6597	0.042*
N21	0.60191 (18)	0.29761 (6)	0.58834 (13)	0.0342 (3)
C28	0.6848 (3)	0.31610 (9)	0.71350 (17)	0.0426 (4)
H28A	0.6331	0.3493	0.7276	0.064*
H28B	0.8128	0.3201	0.7215	0.064*
H28C	0.6637	0.2912	0.7748	0.064*

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
V1	0.03170 (16)	0.03642 (17)	0.02820 (16)	−0.00124 (12)	0.00694 (11)	−0.00003 (12)
O1	0.0416 (7)	0.0421 (7)	0.0386 (7)	0.0028 (6)	0.0140 (5)	0.0045 (6)
C1	0.0413 (9)	0.0401 (10)	0.0313 (9)	−0.0078 (8)	0.0092 (7)	−0.0013 (8)
C2	0.0401 (9)	0.0438 (11)	0.0339 (9)	−0.0078 (8)	0.0101 (7)	−0.0057 (8)
C3	0.0530 (12)	0.0571 (13)	0.0436 (11)	−0.0077 (10)	0.0207 (9)	−0.0064 (10)
C4	0.0709 (15)	0.0681 (15)	0.0409 (12)	−0.0128 (12)	0.0264 (11)	−0.0010 (11)
C5	0.0711 (14)	0.0583 (14)	0.0364 (11)	−0.0081 (11)	0.0141 (10)	0.0083 (10)
C6	0.0543 (11)	0.0454 (11)	0.0398 (10)	−0.0047 (9)	0.0105 (9)	0.0034 (9)
C7	0.0324 (8)	0.0467 (11)	0.0369 (10)	−0.0013 (8)	0.0098 (7)	−0.0075 (8)
N1	0.0322 (7)	0.0392 (8)	0.0325 (8)	−0.0001 (6)	0.0045 (6)	−0.0042 (7)
C8	0.0383 (9)	0.0494 (11)	0.0373 (10)	0.0086 (8)	0.0044 (8)	0.0007 (9)
O11	0.0380 (7)	0.0424 (7)	0.0389 (7)	−0.0036 (5)	0.0095 (5)	−0.0066 (6)
C11	0.0474 (10)	0.0388 (10)	0.0341 (9)	−0.0077 (8)	0.0084 (8)	−0.0013 (8)
C12	0.0548 (11)	0.0368 (10)	0.0343 (9)	0.0009 (8)	0.0121 (8)	0.0004 (8)
C13	0.0704 (14)	0.0423 (11)	0.0411 (11)	0.0029 (10)	0.0145 (10)	−0.0016 (9)
C14	0.0837 (17)	0.0436 (12)	0.0470 (12)	−0.0106 (11)	0.0156 (11)	−0.0081 (10)
C15	0.0691 (15)	0.0558 (14)	0.0484 (12)	−0.0224 (12)	0.0075 (11)	−0.0072 (11)
C16	0.0520 (11)	0.0516 (12)	0.0459 (11)	−0.0129 (10)	0.0095 (9)	−0.0056 (10)
C17	0.0460 (10)	0.0461 (11)	0.0393 (10)	0.0074 (8)	0.0148 (8)	0.0021 (9)
N11	0.0366 (8)	0.0424 (9)	0.0356 (8)	0.0021 (6)	0.0105 (6)	0.0010 (7)
C18	0.0337 (9)	0.0605 (13)	0.0550 (13)	0.0018 (9)	0.0120 (9)	−0.0024 (11)
O21	0.0391 (6)	0.0404 (7)	0.0296 (6)	−0.0041 (5)	0.0041 (5)	0.0024 (5)
C21	0.0287 (8)	0.0365 (9)	0.0333 (9)	0.0011 (7)	0.0076 (6)	−0.0002 (7)

C22	0.0317 (8)	0.0394 (10)	0.0345 (9)	0.0025 (7)	0.0074 (7)	0.0020 (8)
C23	0.0429 (10)	0.0417 (11)	0.0477 (11)	0.0003 (8)	0.0073 (8)	0.0075 (9)
C24	0.0461 (11)	0.0408 (11)	0.0605 (13)	−0.0048 (9)	0.0074 (10)	0.0008 (10)
C25	0.0375 (9)	0.0464 (11)	0.0463 (11)	−0.0026 (8)	0.0062 (8)	−0.0075 (9)
C26	0.0325 (8)	0.0456 (11)	0.0342 (9)	0.0005 (7)	0.0066 (7)	−0.0038 (8)
C27	0.0311 (8)	0.0431 (10)	0.0305 (8)	0.0038 (7)	0.0070 (6)	0.0070 (8)
N21	0.0318 (7)	0.0427 (9)	0.0271 (7)	0.0009 (6)	0.0045 (6)	0.0015 (6)
C28	0.0430 (10)	0.0553 (12)	0.0271 (9)	−0.0048 (9)	0.0032 (7)	0.0012 (9)

Geometric parameters (Å, °)

V1—O11	1.9183 (13)	C13—H13	0.9500
V1—O1	1.9227 (14)	C14—C15	1.390 (4)
V1—O21	1.9641 (13)	C14—H14	0.9500
V1—N1	2.1126 (15)	C15—C16	1.386 (3)
V1—N11	2.1163 (16)	C15—H15	0.9500
V1—N21	2.1625 (15)	C16—H16	0.9500
O1—C1	1.318 (2)	C17—N11	1.281 (3)
C1—C6	1.408 (3)	C17—H17	0.9500
C1—C2	1.415 (3)	N11—C18	1.467 (2)
C2—C3	1.414 (3)	C18—H18A	0.9800
C2—C7	1.439 (3)	C18—H18B	0.9800
C3—C4	1.371 (3)	C18—H18C	0.9800
C3—H3	0.9500	O21—C21	1.310 (2)
C4—C5	1.389 (4)	C21—C22	1.412 (3)
C4—H4	0.9500	C21—C26	1.414 (2)
C5—C6	1.376 (3)	C22—C23	1.402 (3)
C5—H5	0.9500	C22—C27	1.447 (3)
C6—H6	0.9500	C23—C24	1.379 (3)
C7—N1	1.290 (2)	C23—H23	0.9500
C7—H7	0.9500	C24—C25	1.392 (3)
N1—C8	1.463 (2)	C24—H24	0.9500
C8—H8A	0.9800	C25—C26	1.379 (3)
C8—H8B	0.9800	C25—H25	0.9500
C8—H8C	0.9800	C26—H26	0.9500
O11—C11	1.325 (2)	C27—N21	1.284 (2)
C11—C16	1.404 (3)	C27—H27	0.9500
C11—C12	1.420 (3)	N21—C28	1.471 (2)
C12—C13	1.403 (3)	C28—H28A	0.9800
C12—C17	1.446 (3)	C28—H28B	0.9800
C13—C14	1.371 (3)	C28—H28C	0.9800
O11—V1—O1	98.00 (6)	C14—C13—H13	119.2
O11—V1—O21	171.59 (6)	C12—C13—H13	119.2
O1—V1—O21	90.39 (6)	C13—C14—C15	119.2 (2)
O11—V1—N1	87.14 (6)	C13—C14—H14	120.4
O1—V1—N1	88.60 (6)	C15—C14—H14	120.4
O21—V1—N1	92.56 (6)	C16—C15—C14	120.7 (2)

O11—V1—N11	88.49 (6)	C16—C15—H15	119.7
O1—V1—N11	89.82 (6)	C14—C15—H15	119.7
O21—V1—N11	92.09 (6)	C15—C16—C11	121.1 (2)
N1—V1—N11	175.10 (6)	C15—C16—H16	119.4
O11—V1—N21	87.96 (6)	C11—C16—H16	119.4
O1—V1—N21	173.21 (6)	N11—C17—C12	126.48 (19)
O21—V1—N21	83.70 (5)	N11—C17—H17	116.8
N1—V1—N21	94.99 (6)	C12—C17—H17	116.8
N11—V1—N21	87.06 (6)	C17—N11—C18	117.16 (17)
C1—O1—V1	133.07 (12)	C17—N11—V1	125.15 (13)
O1—C1—C6	118.64 (18)	C18—N11—V1	117.65 (13)
O1—C1—C2	122.96 (17)	N11—C18—H18A	109.5
C6—C1—C2	118.39 (18)	N11—C18—H18B	109.5
C3—C2—C1	118.79 (19)	H18A—C18—H18B	109.5
C3—C2—C7	117.42 (19)	N11—C18—H18C	109.5
C1—C2—C7	123.78 (17)	H18A—C18—H18C	109.5
C4—C3—C2	121.6 (2)	H18B—C18—H18C	109.5
C4—C3—H3	119.2	C21—O21—V1	135.69 (11)
C2—C3—H3	119.2	O21—C21—C22	122.67 (16)
C3—C4—C5	119.3 (2)	O21—C21—C26	119.81 (16)
C3—C4—H4	120.3	C22—C21—C26	117.52 (17)
C5—C4—H4	120.3	C23—C22—C21	119.98 (17)
C6—C5—C4	120.8 (2)	C23—C22—C27	117.86 (17)
C6—C5—H5	119.6	C21—C22—C27	122.03 (17)
C4—C5—H5	119.6	C24—C23—C22	121.5 (2)
C5—C6—C1	121.1 (2)	C24—C23—H23	119.3
C5—C6—H6	119.5	C22—C23—H23	119.3
C1—C6—H6	119.5	C23—C24—C25	118.7 (2)
N1—C7—C2	126.67 (18)	C23—C24—H24	120.6
N1—C7—H7	116.7	C25—C24—H24	120.6
C2—C7—H7	116.7	C26—C25—C24	121.06 (19)
C7—N1—C8	116.91 (16)	C26—C25—H25	119.5
C7—N1—V1	124.86 (13)	C24—C25—H25	119.5
C8—N1—V1	118.07 (12)	C25—C26—C21	121.15 (18)
N1—C8—H8A	109.5	C25—C26—H26	119.4
N1—C8—H8B	109.5	C21—C26—H26	119.4
H8A—C8—H8B	109.5	N21—C27—C22	126.46 (17)
N1—C8—H8C	109.5	N21—C27—H27	116.8
H8A—C8—H8C	109.5	C22—C27—H27	116.8
H8B—C8—H8C	109.5	C27—N21—C28	115.14 (16)
C11—O11—V1	132.48 (12)	C27—N21—V1	127.10 (12)
O11—C11—C16	118.88 (19)	C28—N21—V1	117.71 (12)
O11—C11—C12	123.17 (17)	N21—C28—H28A	109.5
C16—C11—C12	117.93 (19)	N21—C28—H28B	109.5
C13—C12—C11	119.37 (19)	H28A—C28—H28B	109.5
C13—C12—C17	117.2 (2)	N21—C28—H28C	109.5
C11—C12—C17	123.40 (18)	H28A—C28—H28C	109.5
C14—C13—C12	121.7 (2)	H28B—C28—H28C	109.5

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C27—H27 $\cdots$ O21 ⁱ	0.95	2.56	3.431 (2)	153

Symmetry code: (i) $x, -y+1/2, z+1/2$.