

# Taxonomic study on *Lathrobium* Gravenhorst (Coleoptera, Staphylinidae, Paederinae) from Longwangshan Mountain, East China

Zhong Peng<sup>1,†</sup>, Li-Zhen Li<sup>1,‡</sup>, Mei-Jun Zhao<sup>1,§</sup>

<sup>1</sup> Department of Biology, College of Life and Environmental Sciences, Shanghai Normal University, Shanghai, 200234, P. R. China

<sup>†</sup> [urn:lsid:zoobank.org:author:6CDD4A22-46AC-4DAF-8409-6D1922AE5D81](https://zoobank.org/urn:lsid:zoobank.org:author:6CDD4A22-46AC-4DAF-8409-6D1922AE5D81)

<sup>‡</sup> [urn:lsid:zoobank.org:author:BBACC7AE-9B70-4536-ABBE-54183D2ABD45](https://zoobank.org/urn:lsid:zoobank.org:author:BBACC7AE-9B70-4536-ABBE-54183D2ABD45)

<sup>§</sup> [urn:lsid:zoobank.org:author:FBE4FA1F-5BCF-405E-98C1-4F86894643EA](https://zoobank.org/urn:lsid:zoobank.org:author:FBE4FA1F-5BCF-405E-98C1-4F86894643EA)

Corresponding author: Mei-Jun Zhao ([mjzhao@shnu.edu.cn](mailto:mjzhao@shnu.edu.cn))

---

Academic editor: V. Assing | Received 18 November 2011 | Accepted 6 January 2012 | Published 13 January 2012

[urn:lsid:zoobank.org:pub:8FCBD1E9-A028-401D-8E0E-E78F7AF869DF](https://zoobank.org/urn:lsid:zoobank.org:pub:8FCBD1E9-A028-401D-8E0E-E78F7AF869DF)

---

**Citation:** Peng Z, Li L-Z, Zhao M-J (2012) Taxonomic study on *Lathrobium* Gravenhorst (Coleoptera, Staphylinidae, Paederinae) from Longwangshan Mountain, East China. ZooKeys 165: 21–32. doi: 10.3897/zookeys.165.2384

---

## Abstract

Species of the genus *Lathrobium* Gravenhorst from Longwangshan Mountain, Zhejiang, East China are studied. A total of five species are recognized, among which three are described here as new: *L. lingae* sp. n., *L. longwangshanense* sp. n. and *L. unicum* sp. n., one species was unidentified and the female of *L. tianmushanense* Watanabe is newly reported. All of these species are illustrated and keyed.

## Keywords

Coleoptera, Staphylinidae, taxonomy, *Lathrobium*, new species, key, Longwangshan, China

## Introduction

To the present, a total of 625 species of the genus *Lathrobium* Gravenhorst have been known worldwide, 64 of them from China (Löbl and Smetana 2004 and subsequent papers). Longwangshan Mountain (at. 30°24'N, 119°27'E) in the south of the Anji County forms the Tianmushan mountain range and Mt. West Tianmushan stand 9.5 kilometers apart. Only two species of *Lathrobium* have been recorded by Watanabe

(1999) from West Tianmushan Mountain, but no members of the genus have been reported yet from Longwangshan Mountain, Zhejiang Province, East China.

In recent years, we made several collecting trips to the Longwangshan Mountain, and obtained a lot of *Lathrobium* specimens. On the basis of the examination, four species were recognized and one species was unidentified, among which three were revealed to be new, and the female of *L. tianmushanense* Watanabe was newly discovered. The purpose of this paper is to describe and illustrate the *Lathrobium* species of Longwangshan Mountain, and to provide an identification key to *Lathrobium* species of Longwangshan Mountain and West Tianmushan Mountain.

## Material and methods

All specimens were collected from the leaf litter of the forest floor by sifting. They were killed with ethylacetate and then dried. Dissections were done in water. The genital organs and other dissected parts were mounted in Euparal (Chroma Gesellschaft Schmidt, Koengen, Germany) on plastic slides that were placed on the same pin as the specimen. Photos were taken by a Canon EOS 40D Camera with an MP-E 65 mm Macro Lens or by a Canon G9 Camera mounted on an Olympus CX31 microscope. The type specimens are deposited in the Insect Collection of Shanghai Normal University (SNUC).

The following abbreviations are used in the text, with all measurements in millimeters:

<b>BL</b>	length of the body from the labral anterior margin to the anal end
<b>HL</b>	length of the head from the clypeal anterior margin to the head base
<b>HW</b>	maximum width of the head
<b>PL</b>	length of the pronotum along the midline
<b>PW</b>	maximum width of the pronotum
<b>EL</b>	length of the elytra from the apex of the scutellum to the elytral posterior margin

## Taxonomy

### *Lathrobium lingae* sp. n.


urn:lsid:zoobank.org:act:741943E9-A379-42E0-807D-811A596FCE88

[http://species-id.net/wiki/Lathrobium\\_lingae](http://species-id.net/wiki/Lathrobium_lingae)

Figs 1A, 3

**Type locality.** Longwangshan Nature Reserve, Zhejiang Province, East China

**Type material** (1 ♂). HOLOTYPE: ♂, labeled ‘CHINA: ZHEJIANG Prov. / Anji County / Longwang Mt. / 25.iv.2006, alt. 950–1,200 m / Rui-Fen Ling leg.’.



**Figures 1.** Male habitus of *Lathrobium* spp., **A** *L. lingae* **B** *L. longwangshanense* **C** *L. tianmushanense*. Scales: 1.0 mm.

**Description.** Measurements and ratios (holotype): BL 6.88, HL 1.00, HW 1.11, PL 1.27, PW 1.15, EL 1.36, HL/HW 0.91, HW/PW 0.96, HL/PL 0.79, PL/PW 1.11, EL/PL 0.71.

Male (Fig. 1A). Body brown with paler apex, legs reddish brown, antennae reddish brown to yellowish brown.

Head quadrate; posterior angles broadly rounded; postgenae weakly convex ventrally; integument with coarse and moderately dense punctation; eyes reduced.

Pronotum slightly stocky, slightly broader than head; punctation sparser than that of head; interstices shining, lacking microsculpture.

Elytra at suture distinctly shorter than pronotum; wider than long; punctation well-defined; and hind wings completely reduced.

Abdomen with dense pubescence; sternite VII (Fig. 3A) with short dark modified setae on postero-median semicircular impression; sternite VIII (Fig. 3C) with triangular emargination and with short dark modified setae on deep impression; sternite IX (Fig. 3B) asymmetrical; aedeagus (Fig. 3D, 3E) with conspicuously long, slender ventral process and twisted dorsal sclerites.

Female. Unknown.

**Distribution.** East China (Zhejiang: Longwangshan Mountain).

**Etymology.** The species is named after Rui-Fen Ling, who collected the type specimens.

**Remarks.** The new species is close to *L. fengae* in similar general form. *Lathrobium lingae* differs especially by the male sternite VII with deeper impression, male sternite VIII with regularly triangular apico-median emargination, and aedeagus with gracile

ventral process and single dorsal sclerite. *Lathrobium fengae* has the male sternite VII with shallower impression, male sternite VIII with irregular apico-median emargination, and aedeagus with broad ventral process and two dorsal sclerites.

***Lathrobium longwangshanense* sp. n.**

urn:lsid:zoobank.org:act:FFAABDE9-CA6B-4DD8-971D-41BC9898C0E0

[http://species-id.net/wiki/Lathrobium\\_longwangshanense](http://species-id.net/wiki/Lathrobium_longwangshanense)

Figs 1B, 4

**Type locality.** Longwangshan Nature Reserve, Zhejiang Province, East China

**Type material** (1 ♂). HOLOTYPE: ♂, labeled ‘CHINA: ZHEJIANG PROV. / Anji County / Longwang Mt. / 25.iv.2006, alt. 950–1,200 m / Yong-Yin Wang leg.’

**Description.** Measurements and ratios (holotype): BL 9.56, HL 1.51, HW 1.58, PL 1.81, PW 1.59, EL 1.32, HL/HW 0.95, HW/PW 0.95, HL/PL 0.83, PL/PW 1.09, EL/PL 0.73.

Habitus as in Fig. 1B. Externally similar to *L. lingae*, except for the lighter average coloration, the somewhat larger body size, the denser punctation on the head and the pronotum.

Male. Sternite VI (Fig. 4A) with tufted pubescence same length as concavity; sternite VII (Fig. 4B) with weak emargination; sternite VIII (Fig. 4C) with darkish setae on impression and basal angle of asymmetrical triangular emargination with dense point-like seta; sternite IX (Fig. 4D) slightly acute anteriorly; aedeagus (Fig. 4E, 4F) with distinct long ventral process and twisted dorsal sclerites.

Female. Unknown.

**Distribution.** East China (Zhejiang: Longwangshan Mountain).

**Etymology.** The species is named after its type locality.

**Remarks.** The new species is similar in most respects to *L. tianmushanense*, but it differs in having relatively stout body, HL/PL being more than 0.80, male sternite VI with tufted pubescence at concavity and aedeagus with longer twisted dorsal sclerites. In *L. tianmushanense*, the body is relatively slender, HL/PL is more than 0.73, the male sternite VI has the concavity lacking pubescence and the dorsal sclerites of the aedeagus are much shorter.

***Lathrobium tianmushanense* Watanabe**

[http://species-id.net/wiki/Lathrobium\\_tianmushanense](http://species-id.net/wiki/Lathrobium_tianmushanense)

Figs 1C, 5

*Lathrobium tianmushanense* Watanabe, 1999: 249

**Type locality.** West Tianmushan Mountain, Zhejiang Province, East China

**Material studied** (3 ♂♂, 4 ♀♀). 1 ♂, 1 ♀, labelled ‘CHINA: ZHEJIANG PROV. / Anji County / Longwang Mt. / Qianmutian / 27.v.2009, alt. 1,300 m, / Yuan, Liu, Feng & Yin leg.’. 1 ♂, 3 ♀♀, same label data, but ‘29.v.2009’.

**Rescription.** For detailed male description of male see Watanabe (1999: 249).

Female. BL 8.06–8.34; Measurements and ratios: HL 1.17, HW 1.32, PL 1.55, PW 1.36, EL 1.02, HL/HW 0.89, HW/PW 0.97, HL/PL 0.76, PL/PW 1.14, EL/PL 0.88.

Slightly smaller than male; posterior margin of tergite VIII (Fig. 5G) weakly asymmetrical; sternite VIII (Fig. 5H) distinctly pointed in the middle; tergite IX (Fig. 5I) (not separated from X) with long and acute lateral processes; tergite X (Fig. 5I) slightly shorter than tergite IX.

**Distribution.** East China (Zhejiang: Longwangshan and Tianmushan Mountains).

**Remarks.** *Lathrobium tianmushanense* is closest to *L. cooteri* from Zhejiang by sharing a similar general form. It can be readily separated by the male sternite VI with modified setae at the concavity and male sternite VIII with relatively regular emargination. While *L. cooteri* has the male sternite VI lacking sexual characters and male sternite VIII possess an irregular emargination.

***Lathrobium uncum* sp. n.**

urn:lsid:zoobank.org:act:FBC72B91-DE34-49D0-9918-A486421C9F48

[http://species-id.net/wiki/Lathrobium\\_uncum](http://species-id.net/wiki/Lathrobium_uncum)

Figs 2A, 6

**Type locality.** Longwangshan Nature Reserve, Zhejiang Province, East China

**Type material** (7 ♂♂, 5 ♀♀). HOLOTYPE: ♂, labeled ‘CHINA: ZHEJIANG PROV. / Anji County / Longwang Mt. / 25.iv.2006, alt. 950–1,200 m / Tang Liang leg.’. PARATYPES: 7 ♂♂, 5 ♀♀, same label data as holotype; 1 ♂, 4 ♀♀, same, but ‘Qianmutian / 27.v.2009, alt. 1,300 m / Yuan, Liu, Feng & Yin leg.’.

**Description.** Measurements and ratios: BL 5.35–5.93. Holotype: HL 0.78, HW 0.81, PL 1.02, PW 0.82, EL 0.72, HL/HW 0.97, HW/PW 0.98, HL/PL 0.77, PL/PW 1.25, EL/PL 0.70.

Habitus as in Fig. 2A. Externally similar to *L. lingae*, except for the lighter average coloration, the somewhat smaller body size and the sparser punctuation on the head and the pronotum.

Male. Sternite VII (Fig. 6A) with a group of coarse setae at middle; sternite VIII (Fig. 6B) with half elliptical median emargination, two rows of modified setae in large but faint apical impression and another row along the posterior margin; sternite IX (Fig. 6C) with cuttle-bone-shaped impression; aedeagus (Fig. 6D, 6E) with a hook-shaped ventral process.

Female. Tergite VIII (Fig. 6F) with posterior margin saliently curved; sternite VIII (Fig. 6G) weakly convex posteriorly and with inconspicuous micropubescence; tergite IX (Fig. 6H) (not separated from X), with long lateral processes; tergite X relatively short.

**Distribution.** East China (Zhejiang: Longwangshan Mountain).

**Etymology.** The specific name ‘*uncum*’ (Latin adjective) means ‘hooked’.

**Remarks.** The new species and *L. tamurai* from Zhejiang share many features, particularly the general form. They can be readily distinguished by the male sternite VII




**Figures 2.** Male habitus of *Lathrobium* spp., **A** *L. uncum* **B** *Lathrobium* sp. indet. Scales: 1.0 mm.

lacking an impression and with several modified discal setae, male sternite VIII with two rows of modified setae in large but faint apical impression and another row along the posterior margin, and slender aedeagus with the ventral process being hook-shaped in *L. uncum*, whereas in *L. tamurai*, the male sternite VII has the impression evenly covered with modified setae on each side of the median part; the male sternite VIII has a horseshoe-shaped impression with dense modified setae; and the robust aedeagus has a straight ventral process.

***Lathrobium* sp. indet.**

Figs 2B, 7

**Material studied** (3 ♂♂, 3 ♀♀). 3 ♂♂, 3 ♀♀, labelled ‘CHINA: Zhejiang Prov / Anji County / Longwang Mt. / Qianmutian / 25.v.2009, alt. 1300 m / Yuan, Liu, Feng & Yin leg.’.




**Figures 3.** *Lathrobium lingae*. **A** male sternite VII **B** male sternite IX **C** male sternite VIII **D** aedeagus in lateral view **E** aedeagus in ventral view. Scales: 0.5 mm.

**Description.** Measurements and ratios: BL 6.43–6.65. Holotype: HL 0.78, PL 1.12, PW 0.93, EL 0.74, HL/HW 0.95, HW/PW 0.96, HL/PL 0.74, PL/PW 1.20, EL/PL 0.66.

Habitus as in Fig. 2B. Externally similar to *L. lingae*, except for the somewhat smaller body size and the more oblong pronotum.

Male. Sternite VII (Fig. 7A) with conspicuously modified setae at weak impression; sternite VIII (Fig. 7B) with approximately elliptic impression and furnished with numerous peg-setae, emargination irregularly shaped; sternite IX (Fig. 7C) anisomerous; aedeagus (Fig. 7D, 7E) with broad ventral process and two apical gracile dorsal sclerites.


Female. Posterior margins of tergite VIII (Fig. 7F) indistinctly asymmetrical and sternite VIII (Fig. 7G) obtusely produced at middle; tergite IX (Fig. 7H) not separated clearly and its lateral processes acute apically; tergite X relatively short.



**Figures 4.** *Lathrobium longwangshanense*. **A** male sternite VI **B** male sternite VII **C** male sternite VIII **D** male sternite IX **E** aedeagus in lateral view **F** aedeagus in ventral view. Scales: 0.5 mm.

**Distribution.** East China (Zhejiang: Longwangshan Mountain).

**Remarks.** The species resembles *L. rougemonti* Watanabe from Zhejiang by sharing the similar form, male sternite VII with weak impression and male sternite VIII with many dark setae in the large impression, but that of aedeagus typically with two closer dorsal sclerites. In *L. rougemonti*, aedeagus have two widely separated dorsal sclerites. The original description of *L. rougemonti* is based on the holotype from West Tianmushan Mountain. The type was not examined, but based on the description and




**Figures 5.** *Lathrobium tianmushanense*. **A** male sternite VI **B** male sternite VII **C** male sternite VIII **D** male sternite IX **E** aedeagus in lateral view **F** aedeagus in ventral view. **G** female tergite VIII **H** female sternite VIII **I** female tergite IX–X. Scales: 0.5 mm.

the illustration (habitus, male abdominal apex and aedeagus) provided by Watanabe (1999), there is still doubt whether the population from Longwangshan represents a new species. As intermediate form exists, it will be necessary to study the type material of *L. rougemonti* for clarification. (Assing pers. comm.)


**A key to the *Lathrobium* species from Longwang – West Tianmu Mountains**

- 1        Length of body larger than 9 mm ..... 2
- Length of body no more than 7 mm..... 3



**Figures 6.** *Lathrobium unicum*. **A** male sternite VII **B** male sternite VIII **C** male sternite IX **D** aedeagus in lateral view **E** aedeagus in ventral view. **F** female tergite VIII **G** female sternite VIII **H** female tergite IX–X. Scales: 0.5 mm.

- 2      Relatively slender (Fig. 1C), HL/PL no more than 0.75; male sternite VI (Fig. 5A) lacking tuft of pubescence at concavity; aedeagus (Fig. 5E, 5F) with short ventral process. Posterior margin of female tergite VIII (Fig. 5G) weakly asymmetrical; female sternite VIII (Fig. 5H) distinctly pointed in the middle ..... *L. tianmushanense* Watanabe
- Relatively stout (Fig. 1B), HL/PL more than 0.80; male sternite VI (Fig. 4A) with tuft of pubescence at concavity; aedeagus (Fig. 4E, 4F) with long ventral process. Female unknown ..... *L. longwangshanense* sp. n.
- 3      Light brown (Fig. 2A); male sternite VII (Fig. 6A) with modified discal setae; male sternite VIII (Fig. 6B) with sparse modified setae in shallow impression; aedeagus (Fig. 6D, 6E) elongate and with hook-shaped ventral process. Female sternite VIII (Fig. 6G) with inconspicuous micropubescence posteriorly ..... *L. unicum* sp. n.



**Figures 7.** *Lathrobium* sp. indet.. **A** male sternite VII **B** male sternite VIII **C** male sternite IX **D** aedeagus in lateral view **E** aedeagus in ventral view. **F** female tergite VIII **G** female sternite VIII **H** female tergite IX–X. Scales: 0.5 mm.

- Brown (Fig. 1A); male sternite VII (Fig. 3A) with modified setae at postero-medial margin; male sternite VIII (Fig. 3C) with dense modified setae in deep impression; aedeagus (Fig. 3D, 3E) robust and not as above. Female sternite VIII without micropubescence posteriorly (*L. lingae* female unknown).....4
- 4 Male sternite VII (Fig. 3A) with deep apico-medial impression; male sternite VIII (Fig. 3C) with regular triangular emargination; aedeagus (Fig. 3D, 3E) with single dorsal sclerite ..... ***L. lingae* sp. n.**
- Male sternite VII with shallow apico-medial impression; male sternite VIII with irregular emargination, aedeagus with two dorsal sclerites ..... ***L. rougemonti* Watanabe.**

**Acknowledgements**

All the collectors mentioned in the text are acknowledged for the field work. We are grateful to the two anonymous reviewers for the critical reading of the manuscript and

helpful comments. The study is supported by the National Natural Science Foundation of China (No. 31101659 and No. 31172134), Foundation of Shanghai Municipal Education Commission (No. 12YZ077) and by Shanghai Normal University (No. SK201234).

## References

- Gravenhorst JLC (1802) Coleoptera Microptera Brunsvicensia nec non exoticorum quotquot exstant in collectionibus entomologorum Brunsvicensium in genera familias et species distribuit. Brunsvigae, Carolus Richard, lxvi + 206 pp.
- Smetana A (2004) Subfamily Staphylininae Latreille, 1802: In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera, Vol. 2. Apollo Books, Stenstrup, 593–598.
- Watanabe Y (1999) Two new species of the group of *Lathrobium pollens* / *brachypterum* (Coleoptera, Staphylinidae) from Zhejiang Province, East China. *Elytra* 27(2): 573–580.
- Watanabe Y (1999) Two new subterranean staphylinids (Coleoptera) from East China. *Elytra* 27(1): 249–257.
- Watanabe Y, Luo ZY (1992) New species of genus *Lathrobium* (Coleoptera, Staphylinidae) from the Wu-yan-ling Nature Protective Area in Zhejiang. *Elytra* 20(1): 47–56.