

SCIENTIFIC REPORTS



OPEN

Trans-generational responses to low pH depend on parental gender in a calcifying tubeworm

Ackley Lane¹, Camilla Campanati¹, Sam Dupont² & Vengatesen Thiyagarajan¹

Received: 03 October 2014

Accepted: 24 April 2015

Published: 03 June 2015

The uptake of anthropogenic CO₂ emissions by oceans has started decreasing pH and carbonate ion concentrations of seawater, a process called ocean acidification (OA). Occurring over centuries and many generations, evolutionary adaptation and epigenetic transfer will change species responses to OA over time. Trans-generational responses, via genetic selection or trans-generational phenotypic plasticity, differ depending on species and exposure time as well as differences between individuals such as gender. Males and females differ in reproductive investment and egg producing females may have less energy available for OA stress responses. By crossing eggs and sperm from the calcareous tubeworm *Hydroides elegans* (Haswell, 1883) raised in ambient (8.1) and low (7.8) pH environments, we observed that paternal and maternal low pH experience had opposite and additive effects on offspring. For example, when compared to offspring with both parents from ambient pH, growth rates of offspring of fathers or mothers raised in low pH were higher or lower respectively, but there was no difference when both parents were from low pH. Gender differences may result in different selection pressures for each gender. This may result in overestimates of species tolerance and missed opportunities of potentially insightful comparisons between individuals of the same species.

As seawater takes up anthropogenic CO₂ emissions the pH and carbonate ion concentrations are reduced, a process called ocean acidification (OA)¹. For most marine invertebrates, OA will be experienced through many generations allowing for trans-generational phenotypic plasticity (TPP) and genetic selection to act^{2–4}. Trans-generational responses (i.e. TPP and/or genetic selection) appear to depend on species, environmental evolutionary history, exposure time and differences between individuals but much remains to be explored^{5,6}. Gender, for example, an easily identifiable characteristic that divides species into two groups, can determine how an individual responds to or is affected by change in pH⁷. Differential investment in reproduction may determine intrinsic tolerance levels⁸, possibly influencing an individual's fitness via the performance of their offspring. As of yet there are very few studies that explicitly examine the differences in stress tolerance between genders. Here, by crossing eggs and sperm from calcareous serpulid polychaete tubeworms raised in two pH environments, we observed that parental gender interacted with parental pH environment to affect offspring performance. The influence of males and females raised in low pH environments on offspring performance (i.e. metamorphosis and post-settlement growth rate) were opposite and additive, low pH males increased performance while low pH females decreased offspring performance. Intrinsic differences between males and females may result in different selection pressures or TPP strategies and while they may be hidden in nature, may shape species response to environmental change over time.

Working toward the ultimate goal of anticipating the effects of OA, most experimental designs use acute exposure of extant marine invertebrates to “future ocean” conditions². Despite the potentially misleading results of acute experiments, very few studies have considered adaptive plasticity or evolutionary

¹The Swire Institute of Marine Science and School of Biological Sciences, The University of Hong Kong, Hong Kong SAR. ²Department of Biological and Environmental Sciences, The Sven Lovén Centre for Marine Sciences - Kristineberg, University of Gothenburg, Fiskebäckskil, Sweden. Correspondence and requests for materials should be addressed to A.L. (email: ackleylane@gmail.com) or V.T. (email: rajan@hku.hk)




Figure 1. Experimental design. (1) Wild adults collected and spawned, and gametes were fertilized. (2) Larvae from wild adults were divided among two pH environments and raised to maturity (i.e. F₀). (3) Gametes from F₀ females (eggs) and males (sperm) of both pH environments were collected and fertilized in all possible combinations, creating 4 groups. (4) Each of the groups (F₁ generation) were then divided among high and low pH environments and larval metamorphosis, juvenile growth and juvenile survival were measured (4 F₁ groups × 2 pH environments × 6 replicates = 48 total F₁ culture tanks).

adaptation^{2,9}. A few studies have found standing genetic variation in tolerance to low pH^{10–13}. It has also been shown that low pH can be an evolutionary force by selecting certain genotypes¹⁴ and some have demonstrated that existing populations can exhibit pre-adaptation to low pH conditions^{15–18}. The influence of low pH during the parental life can also influence offspring performance and studies have found positive (e.g. oysters¹⁹ and fish^{20,21}), neutral (e.g. barnacles²²) and negative (e.g. urchins^{5,23}) trans-generational effects. The possibility of adaptation to environmental change over time is clearly demonstrated as standing genetic variation exists, rapid adaptive evolution is possible, differences in pH tolerance between populations exist and parental experience can alter an offspring's phenotype^{5,10–12,14–17,19–23}.

Rarely considered are the intrinsic differences in how males and females experience different pH environments²⁴. Ellis, *et al.*⁷ demonstrated the importance of gender in pH response as the metabolome of males and females of the mussel *Mytilus edulis* differed significantly under stress. Reproductive costs are at the root of differences between males and females in anisogamous systems, with females commonly allocating more energy into reproduction than males²⁵. For example, male and female polychaete worms of the genus *Ophryotrocha* (Dorvilleidae) can have different energy budgets with males investing less in reproduction and growing faster while females invest more in reproduction and grow less²⁶. Responding to and coping with stress is energetically expensive and may demand a reworking of the organisms' energy budget²⁷, which may already be stretched, particularly for females. Reproduction strategies in sessile, broadcast spawning invertebrates tend to maximize maternal fitness and not the fitness of any individual offspring. It is therefore likely that females will prioritize maintenance of their own body function over reproductive investment²⁸. Thus, it can be hypothesized that performance of females exposed to environmental change (e.g. OA), and ultimately their ability to reproduce and the quality of their offspring, will be more strongly affected than males. To date no study has examined the role that parental gender has in determining the subsequent generation's responses when exposed to OA conditions.

Here, trans-generational responses (i.e. TPP and/or genetic selection) to OA were examined by exposing both parents and offspring of the biofouling tube worm *Hydroides elegans* (Haswell, 1883) to two pH environments within their natural range, a high pH (8.1) environment and a low pH (7.8) environment. This species is well suited to laboratory experiments due to the ease of culture and relatively short generation times²⁹ and previous experiments have shown that tube growth and calcification can be affected by low pH conditions^{30–32}. We tested the effects of and interactions between the pH environment experienced by the parents (from hatching to sexual maturation, F₀ generation), parental gender, and the pH experienced by the offspring (F₁ generation, experimental design summarized in Fig. 1) on F₁ generation performances (i.e. metamorphosis success, juvenile growth and juvenile survival). We hypothesized that (i) offspring exposed to the same pH condition as that experienced by their parents will perform relatively better than those exposed to a different pH environment and (ii), because females may be more strongly affected by low pH conditions, possibly due to a tighter energy budget, they may experience stronger selection pressure or be less able to pass on positive trans-generational phenotypic changes.

Results

Larval metamorphosis. Metamorphosis success (%) was measured in both F_0 and F_1 generations (Fig. 2a). More than 50% of all the larvae successfully metamorphosed in the F_1 generation. In the F_0 generation this parameter was significantly affected by block ($F = 19.29$, $p < 0.0001$) but unaffected by pH ($p > 0.05$) or the pH by block interaction ($p > 0.05$). In the F_1 generation metamorphosis success was significantly impacted (3-ways GLM model, $F_{7,35} = 2.31$, $p = 0.048$) by the pH experienced of the F_0 females ($F = 4.72$, $p = 0.037$), F_1 generation pH ($F = 4.33$, $p = 0.045$) and their interactions ($F = 5.37$, $p = 0.026$). The pH experienced of the F_0 males and all other multiple factor interactions were not significant ($p > 0.05$). When the F_0 females were exposed to pH 7.8, the metamorphosis success of offspring exposed to pH 8.1 was decreased by 30% as compared to other treatments.

Post-metamorphic (juvenile) survival. Juvenile survival (in %) was assessed between day 1 and day 7 post-metamorphosis assay (Fig. 2b). In F_0 , pH had a significant effect with juvenile raised at pH 7.8 experiencing a 20% decrease in survival (2-ways GLM model, $F_{5,119} = 19.47$, $p < 0.0001$; pH, $F = 4.55$, $p = 0.035$). A significant block effect, due to multiple collections of wild adults for spawning, was also detected ($F = 44.37$, $p < 0.0001$) but with no interaction with pH ($F = 40.58$, $p = 0.56$). Similar survival around 60% was observed in F_1 but none of the tested parameters (pH treatment, and pH experienced by the parents) had a significant effect (3-ways GLM model, $F_{6,29} = 1.37$, $p = 0.26$).

Post-metamorphic (juvenile) growth. Seawater pH had no significant effect (2-ways GLM model, $F_{5,685} = 303.81$, $p < 0.0001$; pH, $F = 3.07$, $p = 0.095$; block, $F = 908.18$, $p < 0.0001$; pH \times block, $F = 0.17$, $p = 0.98$) on F_0 juvenile growth rate (Fig. 2c). For F_1 , the pH experienced by males (3-ways GLM, model, $F_{7,17} = 5.30$, $p = 0.0024$; males, $F = 6.21$, $p = 0.023$) and female parents ($F = 7.39$, $p = 0.014$) but also the interaction between males and pH ($F = 16.67$, $p = 0.0008$) had an impact on the juvenile growth rates. The F_1 pH and other interactions were not significant ($p > 0.05$). When F_0 males were exposed to pH 7.8 the F_1 juvenile growth rate is increased 2 times when raised at pH 8.1 (e.g. compare F8.1_M8.1 to F8.1_M7.8 in pH 8.1 environment). On the other hand, the F_1 growth rate is decreased by 25% when the F_0 female parents were exposed to pH 7.8 (e.g. F8.1_M8.1 compared to F7.8_M8.1).

Discussion

Both maternal and paternal low pH experience affected F_1 growth rate, and maternal low pH experience affected metamorphosis success, however all differences between groups of the F_1 generation (i.e. due to parental environment and gender) were only apparent when raised at high pH. The differences observed between the F_0 and F_1 generations are notable, and the reason behind the lower performance of the F_0 generation is unknown but may be because the wild adults' environment was unstable with an unpredictable food supply while the laboratory raised parental generation (F_0) had a constant food supply. To summarize, when the eggs came from individuals raised in low pH environments, F_1 metamorphosis and growth rate in the high pH environment were reduced by 30% and 25% respectively. On the other hand, when sperm came from individuals raised in low pH conditions, F_1 growth rate in high pH conditions was more than doubled. Survival data, however, suffered from high variability and no significant difference between groups was observed in the F_1 generation. So, parental gender and pH experience influenced the F_1 generation confirming that there was a trans-generational response, although the majority of the effects were only apparent when F_1 individuals were raised in high pH conditions. It is of particular interest that F_0 females may be more strongly affected by low pH than F_0 males, as F_0 low pH females, unlike F_0 low pH males, produced offspring whose metamorphosis success and growth rates decreased under high pH conditions. The fitness of each gender may be differently affected, and a species mean tolerance (without the consideration of gender) may actually overestimate the actual species tolerance as both genders are equally important in successful reproduction.

This experimental design highlights that parental influence, positive, negative, or neutral can result from a combination of paternal and maternal influence. If examining only F_1 groups whose parents experienced the same pH environments (i.e. F8.1_M8.1 and F7.8_M7.8) one would conclude that there were no significant differences when parents are raised in either low or high pH. However disentangling parental gender by examining F_1 groups where parents experience different pH environments (i.e. F7.8_M8.1 and F8.1_M7.8) shows that the ostensible neutrality of the F_0 exposure (i.e. no differences between groups F7.8_M7.8 and F8.1_8.1) is a product of opposite and additive influences of each parental gender. While in nature genders are unlikely to experience different pH environments, the intrinsic experiences of each gender and their effect on the next generation may provide insight into intra-specific variation and highlight particular and gender-specific mechanisms that are important when responding to pH change²⁴. Differences between genders stem from differences in reproductive investment, sperm is cheap while eggs are expensive^{25,26}. Females may, therefore, have less spare energy to invest in somatic growth or plastic responses that would allow survival and reproduction in changing environments³³. Further support for this hypothesis is given by Holcomb, *et al.*⁸ where high CO_2 impacted calcification only in females (at increased temperature). The long term and potentially evolutionary, consequences due to differences between genders, will be hidden as a population's phenotype changes over time unless specifically investigated. Nonetheless, the role gender differences play in evolution may be substantial as




Figure 2. Metamorphosis success (a, in %), juvenile survival (b, in %) and growth rate (c, in mm day⁻¹) of the second generation (F₁) of *Hydroides elegans* raised in two pH levels. F₁ groups are defined by the pH environment (8.1 vs 7.8) in which each of their parents were raised (F = maternal environment, M = paternal environment): group 1 = F8.1_M8.1; group 2 = F8.1_M7.8; group 3 = F7.8_M8.1; and group 4 = F7.8_M7.8. Mean values are expressed with their standard error of means (mean ± SEM) and the significance level applied was 5%.

a female's tighter energy budget may put them closer to the edge of their tolerance, resulting in stronger selection pressure on female characteristics as environments change.

The mechanisms responsible for the differences observed in the F_1 generation cannot be established from the data presented. Evolutionary selection or trans-generational phenotypic plasticity (TPP) may have acted individually or simultaneously. Evolutionary selection and TPP work via entirely different mechanisms, either selecting more fit phenotypes that are genetically determined (inheritable) or by the ability to pass epigenetic information, energetic advantage via differentially egg provisioning or hormonal signals to offspring³⁴. Each process could explain the environment-phenotype mismatch when F_1 low pH females produced offspring with lower performance in high pH conditions. Likewise, F_1 low pH males produced offspring with increased growth rates under high pH conditions, which could also be an environment-phenotype mismatch depending on whether faster growth is advantageous. Finally, differential maternal provisioning as a general increase or decrease in egg resources, which would exist in both F_1 environments, is unlikely because the differences among F_1 groups were only apparent in high pH treatments. However, advantageous natural selection or TPPs may have counteracted poor egg quality exclusively in the low pH environment.

Despite the ambiguity of the mechanisms behind the parental effects on the responses in the F_1 generation, gender may determine stress tolerance in an individual, possibly because of energy differences. The intrinsic differences in how each gender experiences environmental change became apparent when the effects of males and females from each environment were considered separately, and not as a single pre-exposed unit. Rarely considered are the differences between sexes and pH tolerance, but further research into the genetic, transcriptomic, proteomic or metabolomics levels may provide insight into the gender specific mechanisms that are vital to survival and reproduction as environments change⁷. The opposite and additive influences of each parental gender on offspring observed here may play an important role in evolution as environments change, but also may provide interesting comparisons between highly related, but fundamentally different individuals. Low pH may be a relatively novel stress for which this species has no adapted response, for example parental effects in low salinity environments proved to be positive for both genders in the tube worm *Hydroides diramphus*⁴. The different influence of each gender due to low pH may identify specific characteristics or adaptive strategies beneficial in low pH seawater, giving a more complete picture when predicting species response as OA shifts oceanic carbonate chemistry. A more flexible energy budget may link directly to performance under low pH conditions and translate into stronger natural selection on females than males. Finally, knowing the how each gender is affected by changes in pH will give a more detailed understanding of species tolerances to low pH.

Methods

The tube dwelling serpulid polychaete *Hydroides elegans* is a common biofouler that has been identified as a model species due to its economic and ecological importance, as well as its relatively short generation time (i.e. 2 weeks to maturity) and ease of culture²⁹. *H. elegans* is capable of inhabiting highly variable habitats, like the Eastern coastal waters of Hong Kong, that vary in pH (8.3 to 7.6), temperature (16 to 29 °C) and salinity (30–33)³⁵. For this multi-generational experiment wild adults were collected from the Eastern waters of Hong Kong, China field on 3 occasions and held in laboratory for no longer than 1 week before spawning. Wild adult worms were induced to spawn by gently breaking them from their tubes, those bearing mature gametes spawn immediately (3 males and 3 females on each spawning occasion, referred to as blocks). Within one hour of gamete collection eggs and sperm were combined at sperm concentrations of approximately 1×10^5 sperm ml^{-1} , shown to result in >90% fertilization and very little polyspermy³⁶. The fertilized embryos (F_0 generation) were then divided among replicates of two pH levels in culture tanks of 120 ml with 200–240 larvae in each (3 blocks \times 2 pH levels \times 54 replicates = 324 total replicate tanks, actual replication varied as some replicates did not meet the minimum requirements for inclusion). Filtered seawater in culture tanks was changed every other day and aliquots of concentrated algae cultures (*Isochrysis galbana*) were added to achieve approximate cell densities of 50,000 cells ml^{-1} . All F_0 generation animals used here were maintained for at least 3 months, from fertilization into maturity, in their respective treatments. The tubes of F_0 individuals were all >2 cm when gamete spawning for the F_1 generation was conducted.

For the 2nd generation exposed to 2 pH environments (F_1), 9 males and 9 females were randomly selected from all the F_0 individuals who spawned and crossed to create four groups of egg and sperm combinations: group 1 - pH 8.1 female \times pH 8.1 male (F8.1_M8.1), group 2 - pH 8.1 female \times pH 7.8 male (F8.1_M7.8), group 3 - pH 7.8 female \times pH 8.1 male (F7.8_M8.1), and group 4 - pH 7.8 female \times pH 7.8 male (F7.8_M7.8). The four groups of fertilized embryos (F_1 generation) were then each divided among high pH (8.16) and low pH (7.78) environments (Table 1) with (4 offspring groups \times 2 pH treatments \times 6 replicates = 48 total tanks) and raised from fertilization to 7 days post-metamorphosis to observe performance. Culture tanks, larval densities and food concentrations were all the same as those used for the F_0 generation. The experimental design (spawning, pH environments, eggs and sperm crosses) is summarized in Fig. 1.

Three performance endpoints were measured for the F_0 and F_1 generations: (1) metamorphosis, (2) juvenile survival and (3) juvenile growth. (1) Percent metamorphosis success was calculated as the percentage of larvae successfully metamorphosing within 3 days of attaining competency. Upon larval competency, glass microscope slides with natural biofilms developed for 1 week were introduced

	Measured (\pm S.D.)				Calculated (CO ₂ sys)				
	pH	Temp (°)	Salinity	TA (mM)	pCO ₂	CO ₃ ⁻²	CO ₂	Ω_{Ca}	Ω_{Ar}
F ₀ high pH	8.11 (\pm 0.04)	23.8 (\pm 0.4)	32.7 (\pm 0.6)	2.2865 (\pm 0.113)	493.2	172.4	14.6	4.2	2.8
F ₀ low pH	7.76 (\pm 0.04)	24.0 (\pm 0.4)	32.7 (\pm 0.6)	2.2865 (\pm 0.113)	1231.4	86.2	36.2	2.1	1.4
F ₁ high pH	8.16 (\pm 0.08)	23.7 (\pm 0.5)	33.7 (\pm 0.7)	2.224 (\pm 0.061)	412.4	187.1	12.1	4.5	3.0
F ₁ low pH	7.78 (\pm 0.04)	23.8 (\pm 0.5)	33.6 (\pm 0.7)	2.224 (\pm 0.061)	1127.4	88.8	33.1	2.2	1.4

Table 1. Carbonate chemistry measurements and calculations for the F₀ and the F₁ generations' environment. Calculated parameters were obtained using the CO₂SYS program using the default dissociation constants and the NBS pH scale⁵.

into culture tanks and cultures were maintained for 3 days as normal. After the 3 day metamorphosis period all animals settled on the tank and slide were counted and all larvae still swimming were counted and removed. (2) Percent juvenile survival was calculated as the percentage of attached and calcifying juveniles after the 3 day metamorphosis assay that remained alive for a further 7 days under the same conditions as those during the larval development and metamorphosis periods. (3) Juvenile growth rate (mm day⁻¹) was measured for the first 7 days post metamorphosis by image analysis comparing pictures taken just after the metamorphosis period to those taken 7 days later (ImageJ). Replicates with less than 10 juveniles were not considered for calculation of survival and metamorphosis, and growth rates were only calculated for replicates with more than 3 juveniles. Data were analyzed n-ways GLM models in the SAS/STAT software (SAS Institute 1990), and logarithmic transformations were used where appropriate.

To measure the experimental carbonate chemistry pH_(NBS scale), temperature and salinity were measured daily and total alkalinity (A_T, Apollo SciTech AS-ALK2) was measured in triplicate for all seawater before use. The two pCO₂ environments were created using continuous flow of ambient air with or without gaseous CO₂ enrichment into environmental chambers into which replicate tanks were placed and allowed to equilibrate. Carbonate chemistry was calculated using the program CO₂SYS (Pierrot, Lewis & Wallace 2006, doi : 10.3334/CDIAC/otg.CO2SYS_XLS_CDIAC105a) and resulted in high pH (8.11) and low pH (7.76) for the F₀ generation and high pH (8.16) and low pH (7.78) for the F₁ generation environments with temperature = 24 °C, salinity = 32 ppt and A_T = 2.28 mM and temperature = 24 °C, salinity = 33 and A_T = 2.24 mM respectively (full carbonate chemistry calculation reported in Table 1).

References

1. Sabine, C. L. *et al.* The oceanic sink for anthropogenic CO₂. *Science* **305**, 367–371, doi:10.1126/science.1097403 (2004).
2. Sunday, J. M. *et al.* Evolution in an acidifying ocean. *Trends Ecol Evol* **29**, 117–125, doi:10.1016/j.tree.2013.11.001 (2014).
3. Orr, J. C. *et al.* Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* **437**, 681–686, doi:10.1038/nature04095 (2005).
4. Jensen, N., Allen, R. M. & Marshall, D. J. Adaptive maternal and paternal effects: gamete plasticity in response to parental stress. *Funct Ecol* **28**, 724–733, doi:10.1111/1365-2435.12195 (2014).
5. Dupont, S., Dorey, N., Stumpp, M., Melzner, F. & Thorndyke, M. Long-term and trans-life-cycle effects of exposure to ocean acidification in the green sea urchin *Strongylocentrotus droebachiensis*. *Mar Biol* **160**, 1835–1843, doi:10.1007/s00227-012-1921-x (2013).
6. Burgess, S. C. & Marshall, D. J. Adaptive parental effects: the importance of estimating environmental predictability and offspring fitness appropriately. *Oikos* **123**, 769–776, doi:10.1111/oik.01235 (2014).
7. Ellis, R. P. *et al.* ¹H NMR metabolomics reveals contrasting response by male and female mussels exposed to reduced seawater pH, increased temperature, and a pathogen. *Environ Sci Technol* **48**, 7044–7052, doi:10.1021/es501601w (2014).
8. Holcomb, M., Cohen, A. L. & McCorkle, D. C. An investigation of the calcification response of the scleractinian coral *Astrangia poculata* to elevated pCO₂ and the effects of nutrients, zooxanthellae and gender. *Biogeosciences* **9**, 29–39 (2012).
9. Kroeker, K. J. *et al.* Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob Chang Biol* **19**, 1884–1896, doi:10.1111/gcb.12179 (2013).
10. Foo, S. A., Dworjanyn, S. A., Poore, A. G. & Byrne, M. Adaptive capacity of the habitat modifying sea urchin *Centrostephanus rodgersii* to ocean warming and ocean acidification: performance of early embryos. *PLoS One* **7**, e42497, doi:10.1371/journal.pone.0042497 (2012).
11. Sunday, J. M., Crim, R. N., Harley, C. D. & Hart, M. W. Quantifying rates of evolutionary adaptation in response to ocean acidification. *PLoS One* **6**, e22881, doi:10.1371/journal.pone.0022881 (2011).
12. Pisteos, J. C. A., Calosi, P., Widdicombe, S. & Bishop, J. D. D. Will variation among genetic individuals influence species responses to global climate change? *Oikos* **120**, 675–689, doi:10.1111/j.1600-0706.2010.19470.x (2011).
13. Chirgwin, E., Monro, K., Sgro, C. M. & Marshall, D. J. Revealing hidden evolutionary capacity to cope with global change. *Glob Chang Biol*, n/a–n/a, doi:10.1111/gcb.12929 (2015).
14. Lohbeck, K. T., Riebesell, U. & Reusch, T. B. H. Adaptive evolution of a key phytoplankton species to ocean acidification. *Nat Geosci* **5**, 346–351, doi:10.1038/ngeo1441 (2012).
15. Pespeni, M., Chan, F., Menge, B. & Palumbi, S. Signs of adaptation to aocal pH conditions across an environmental mosaic in the California current ecosystem. *Integr Comp Biol* **53**, 857–870 (2013).
16. Palumbi, S. R., Barshis, D. J., Traylor-Knowles, N. & Bay, R. A. Mechanisms of reef coral resistance to future climate change. *Science* **344**, 895–898, doi:10.1126/science.1251336 (2014).
17. Kelly, M. W., Padilla-Gamiño, J. L. & Hofmann, G. E. Natural variation and the capacity to adapt to ocean acidification in the keystone sea urchin *Strongylocentrotus purpuratus*. *Glob Chang Biol* **19**, 2536–2546, doi:10.1111/gcb.12251 (2013).
18. Calosi, P. *et al.* Adaptation and acclimatization to ocean acidification in marine ectotherms: an *in situ* transplant experiment with polychaetes at a shallow CO₂ vent system. *Phil Trans R Soc B* **368**, 20120444 (2013).

19. Parker, L. M. *et al.* Adult exposure influences offspring response to ocean acidification in oysters. *Glob Change Biol* **18**, 82–92, doi:10.1111/j.1365-2486.2011.02520.x (2012).
20. Miller, G. M., Watson, S.-A., Donelson, J. M., McCormick, M. I. & Munday, P. L. Parental environment mediates impacts of increased carbon dioxide on a coral reef fish. *Nature Clim Change* **2**, 858–861 (2012).
21. Allan, B. J., Miller, G. M., McCormick, M. I., Domenici, P. & Munday, P. L. Parental effects improve escape performance of juvenile reef fish in a high-CO₂ world. *Proc R Soc B* **281**, 20132179, doi:10.1098/rspb.2013.2179 (2014).
22. Pansch, C., Schaub, I., Havenhand, J. & Wahl, M. Habitat traits and food availability determine the response of marine invertebrates to ocean acidification. *Glob Change Biol* **20**, 765–777, doi:10.1111/gcb.12478 (2014).
23. Uthicke, S., Soars, N., Foo, S. & Byrne, M. Effects of elevated pCO₂ and the effect of parent acclimation on development in the tropical Pacific sea urchin *Echinometra mathaei*. *Mar Biol* **160**, 1913–1926 (2013).
24. Bijlsma, R. & Loeschcke, V. Environmental stress, adaptation and evolution: an overview. *J Evol Biol* **18**, 744–749, doi:10.1111/j.1420-9101.2005.00962.x (2005).
25. Hayward, A. & Gillooly, J. F. The cost of sex: quantifying energetic investment in gamete production by males and females. *PLoS One* **6**, e16557, doi:10.1371/journal.pone.0016557 (2011).
26. Premoli, M. & Sella, G. Sex economy in benthic polychaetes. *Ethol Ecol Evol* **7**, 27–48 (1995).
27. Kooijman, S. A. L. M. *Dynamic energy budget theory for metabolic organisation*. (Cambridge University Press, 2000).
28. Marshall, D. J. & Uller, T. When is a maternal effect adaptive? *Oikos* **116**, 1957–1963, doi:10.1111/j.2007.0030-1299.16203.x (2007).
29. Hadfield, M. G. Biofilms and marine invertebrate larvae: what bacteria produce that larvae use to choose settlement sites. *Ann Rev Mar Sci* **3**, 453–470 (2011).
30. Lane, A. C., Mukherjee, J., Chan, V. B. & Thiagarajan, V. Decreased pH does not alter metamorphosis but compromises juvenile calcification of the tube worm *Hydroides elegans*. *Mar Biol* **160**, 1983–1993 (2013).
31. Li, C. *et al.* Weakening Mechanisms of the Serpulid Tube in a High-CO₂ World. *Environ Sci Technol*, doi:10.1021/es501638h (2014).
32. Chan, V. *et al.* CO₂-driven Ocean Acidification Alters and Weakens Integrity of Calcareous Tube Produced by the Serpulid Tubeworm, *Hydroides elegans*. *PLoS One* **7**, e42718 (2012).
33. Dawson, T. E. & Ehleringer, J. R. Gender-specific physiology, carbon isotope discrimination, and habitat distribution in boxelder, *Acer negundo*. *Ecology* **74**, 798–815 (1993).
34. Giennapp, P., Teplitsky, C., Alho, J. S., Mills, J. A. & Merila, J. Climate change and evolution: disentangling environmental and genetic responses. *Mol Ecol* **17**, 167–178, doi:10.1111/j.1365-294X.2007.03413.x (2008).
35. Qiu, J. W. & Qian, P. Y. Combined effects of salinity and temperature on juvenile survival, growth and maturation in the polychaete *Hydroides elegans*. *Mar Ecol Prog Ser* **168**, 127–134, doi:10.3354/meps168127 (1998).
36. Pechenik, J. A., Pearse, J. S. & Qian, P. Y. Effects of salinity on spawning and early development of the tube-building polychaete *Hydroides elegans* in Hong Kong: not just the sperm's fault? *Biol Bull* **212**, 151–160 (2007).

Acknowledgments

The experiment was undertaken at the Swire Institute of Marine Science and funded by three GRF grants from the HKSAR-RGC (Grant Numbers: 780510M and 17304914). S.D. was financially supported by the Linnaeus Centre for Marine Evolutionary Biology at the University of Gothenburg (<http://www.cemeb.science.gu.se/>) and supported by a Linnaeus grant from the Swedish Research Councils VR, Formas.

Author Contributions

A.L. conceived and designed experiments, A.L. and C.C. carried out experimentation, V.T. provided experimental resources, S.D. analyzed data, A.L. lead the co-writing of the paper with C.C., S.D. and V.T.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Lane, A. *et al.* Trans-generational responses to low pH depend on parental gender in a calcifying tubeworm. *Sci. Rep.* **5**, 10847; doi: 10.1038/srep10847 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>