

POSTER PRESENTATION

Open Access



A computational model of Suprachiasmatic Nucleus (SCN) for circadian rhythm modulation based on ion channel conductance and calcium dynamics

Hyoungkyu Kim, Jaeseung Jeong*

From Nineteenth Annual Computational Neuroscience Meeting: CNS*2010
San Antonio, TX, USA. 24-30 July 2010

The suprachiasmatic nucleus (SCN), the master biological clock of circadian rhythms, has been highly interested for biological rhythms in animals. Physiological and molecular biological studies [1-3] and computational modeling approach [4] have investigated channels contributing to circadian rhythms of spike firing frequency. The aim of this study was to investigate possible contribution of each channel to circadian rhythms using Hodgkin-Huxley-type model containing BK channel, potassium channel, sodium channel, and calcium

concentration oscillation. We found that the change in channel conductance modulated firing rates from high regular firing to low irregular firing with high sensitivity of the channel. In agreement with experimental data, the persistent sodium currents also highly contributed to modulation in spike firing frequency. The leak potassium currents and fast delayed rectifier currents exhibited to restrict the circadian firing range and regulation of interspike intervals. Most interestingly, the sinusoidal shape of the diurnal change in the channel conductance reproduced the change in diurnal firing rates. The calcium concentration oscillation and BK channel activity were examined to determine the synchronized circadian



* Correspondence: jsjeong@kaist.ac.kr
Department of Bio and Brain Engineering, KAIST (Korea Advanced Institute of Science and Technology), Daejeon, South Korea

oscillation and phase shifting of firing rate in the neural network models structured by graph theory. We suggest that the model may be a guidance of channel properties in experimental conditions and be helpful to explain the circadian behaviors of SCN neurons.

Published: 20 July 2010

References

1. Jackson AC, Yao GL, Bean BP: **Mechanism of Spontaneous Firing in Dorsomedial Suprachiasmatic Nucleus Neurons.** *Journal of Neuroscience* 2004, **24**(37):7985-7998.
2. Pennartz CMA, de Jeu MTG, Bos NPA, Schaap J, Geurtsen AMS: **Diurnal modulation of pacemaker potentials and calcium current in the mammalian circadian clock.** *Nature* 2002, **416**:286-290.
3. Ikeda M, Sugiyama T, Wallace CS, Gompf HS, Yoshioka T, Miyawaki A, Allen CN: **Circadian Dynamics of Cytosolic and Nuclear Ca²⁺ in Single Suprachiasmatic Nucleus Neurons.** *Neuron* 2003, **38**(2):253-263.
4. Choon Kiat Sim, Daniel BForger: **Modeling the Electrophysiology of Suprachiasmatic Nucleus Neurons.** *J Biol Rhythms* 2007, **22**:445.

doi:10.1186/1471-2202-11-S1-P160

Cite this article as: Kim and Jeong: A computational model of Suprachiasmatic Nucleus (SCN) for circadian rhythm modulation based on ion channel conductance and calcium dynamics. *BMC Neuroscience* 2010 **11**(Suppl 1):P160.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

