

Assessment of Somatization and Medically Unexplained Symptoms in Later Life

Assessment
2018, Vol. 25(3) 374–393
© The Author(s) 2017
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1073191117721740
journals.sagepub.com/home/asm



T. J. W. van Driel¹, P. H. Hilderink¹, D. J. C. Hanssen²,
P. de Boer¹, J. G. M. Rosmalen³, and R. C. Oude Voshaar³

Abstract

The assessment of medically unexplained symptoms and “somatic symptom disorders” in older adults is challenging due to somatic multimorbidity, which threatens the validity of somatization questionnaires. In a systematic review study, the Patient Health Questionnaire–15 (PHQ-15) and the somatization subscale of the Symptom Checklist 90-item version (SCL-90 SOM) are recommended out of 40 questionnaires for usage in large-scale studies. While both scales measure physical symptoms which in younger persons often refer to unexplained symptoms, in older persons, these symptoms may originate from somatic diseases. Using empirical data, we show that PHQ-15 and SCL-90 SOM among older patients correlate with proxies of somatization as with somatic disease burden. Updating the previous systematic review, revealed six additional questionnaires. Cross-validation studies are needed as none of 46 identified scales met the criteria of suitability for an older population. Nonetheless, specific recommendations can be made for studying older persons, namely the SCL-90 SOM and PHQ-15 for population-based studies, the Freiburg Complaint List and somatization subscale of the Brief Symptom Inventory 53-item version for studies in primary care, and finally the Schedule for Evaluating Persistent Symptoms and Somatic Symptom Experiences Questionnaire for monitoring treatment studies.

Keywords

medically unexplained symptoms, MUS, somatic symptoms, somatic symptom disorders, somatoform disorders, aged, aged 80 years and older, instruments

Introduction

Somatization is the tendency to experience and communicate somatic distress in response to psychosocial stress and to seek medical help for it (Lipowski, 1988). The severity of somatization, however, is difficult to measure. In younger people, it is often assessed by a physical symptom count, especially by counting symptoms that often remain medically unexplained in clinical practice, like fatigue or dizziness. Medically unexplained physical symptoms (MUS) might thus be the result of a process of somatization and are a core criterion of somatoform disorders in the *Diagnostic and Statistical Manual of Mental Disorders, Fourth edition, text revised (DSM-IV-TR)*.

With the introduction of the *DSM-5*, the *DSM-IV-TR* section of somatoform disorders has been replaced by the new section of somatic symptom and related disorders (American Psychiatric Association, 2013). The most important change was the focus on so-called positive criteria for establishing a diagnosis, that is, the prominence of a physical symptom associated with significant distress and impairment. The major diagnosis in this section, somatic symptom disorder, emphasizes diagnosis made on the basis of positive symptoms and signs, that is, distressing somatic symptoms plus

abnormal thoughts, feelings, and behaviors in response to these symptoms. This contrasts with somatoform disorders in the *DSM-IV-TR* that poses the absence of a medical explanation for the somatic symptoms a key feature. Although somatic symptom disorders might be more useful for diagnosis in both primary and specialized (somatic) health care, the criteria for these disorders do not necessarily apply to all patients burdened by MUS. MUS, irrespective of the *DSM* classification rules, have consistently been associated with a lower quality of life, psychological distress, and increased medical consumption (Hanssen, Lucassen, Hilderink, Naarding, & Oude Voshaar, 2016; Weiss, Kleinstuber, & Rief, 2016).

A systematic review has identified 40 scales to assess self-report somatic symptoms (Zijlema et al., 2013). The Patient Health Questionnaire–15 (PHQ-15) and the somatization

¹SeniorBeter, Gendt, Netherlands

²Radboud University, Nijmegen, Netherlands

³University of Groningen, Groningen, Netherlands

Corresponding Author:

Richard C. Oude Voshaar, Department of Psychiatry, University Medical Center Groningen, PO Box 30.001, Groningen 9700 RB, Netherlands.
Email: r.c.oude.voshaar@umcg.nl

subscale of the Symptom Checklist 90-item version (SCL-90 SOM) were considered the best options to be used in large-scale population-based studies, based on several criteria among which type of symptoms, time frame, response scale, psychometric characteristics, and patient burden (Zijlema et al., 2013). Since older persons often suffer from physical symptoms due to one or more chronic somatic diseases (multimorbidity; Fortin, Stewart, Poitras, Almirall, & Maddocks, 2012), somatic symptom questionnaires may easily overestimate the severity of somatization in an older sample. Nonetheless, accumulating data emerge that somatization, MUS, and somatoform disorders are highly relevant in older persons, posing a significant burden on health-related quality of life, increased level of health care usage and potentially iatrogenic damage (Hanssen et al., 2016). Nonetheless, actual prevalence rates are somewhat lower among persons older than 65 years compared with younger persons (Hilderink, Collard, Rosmalen, & Oude Voshaar, 2013). In a systematic review of six cohort studies including both younger and older people, prevalence rates for *DSM-IV* defined somatoform disorders ranged from 1.5% through 13.0% (median 5.4%) among people aged 65 years and older, and from 10.7% through 26.8% (median 15.3%) in younger people (Hilderink et al., 2013). Whether these lower prevalence rates in later life are simply artefacts due to falsely attributing physical symptoms to (comorbid) chronic somatic diseases, or reflect real differences due to better coping with chronic illnesses and/or less disease benefits in later life have to be established.

The main objective of the present article is to explore which assessment instrument is optimal for assessing somatization in older adults. In our research program on MUS in later life, we have applied both the PHQ-15 as well as the SCL-90 SOM among older patients with MUS and medically explained symptoms. Therefore, we first test empirically how the PHQ-15 and the SCL-90 SOM relate to other measures of somatization, namely the presence of a somatoform disorder determined by a semistructured psychiatric interview (Lecrubier et al., 1997), and health anxiety measured with the Whitely Index (Pilowsky, 1967), as well as to medically explained somatic disease burden. Since these results were rather disappointing, we decided to update and extend the previously conducted systematic review of Zijlema et al. (2013) to examine which of the assessments of somatic symptoms might be most relevant for assessing somatization in older persons.

Method

Empirical Tests of the PHQ-15 and SCL-90 SOM

For the empirical tests, we used our data set of a pilot study on MUS in older patients as well as a data set on a larger case control study. Both samples have been described

elsewhere in detail (see Hanssen et al., 2016; Hilderink et al., 2009), but can be summarized as follows.

The pilot study was a consecutive case series of 37 older patients referred to our outpatient mental health clinic for a diagnostic procedure and treatment for MUS (Hilderink et al., 2009). In this pilot study, we administered the SCL-90 as well as the PHQ-15 (Kocalevent, Hinz, & Brähler, 2013), whereby the item on menstrual cycle was omitted (being not relevant in later life).

The case control study, acronym OPUS study (Older Persons with medically Unexplained Symptoms study) included 118 older persons suffering from MUS (cases) and 154 older patients suffering from medically explained symptoms for which they frequently attended their general practitioner (Hanssen et al., 2016). In this study, the Brief Symptom Inventory, 53 items (Derogatis & Melisaratos, 1983), an abbreviated version of the SCL-90 with similar psychometric properties, has been administered as a measure for psychopathological distress (Derogatis, 1975).

In both studies, the same two indicators of somatization were administered. First, the Mini International Neuropsychiatric Interview (Lecrubier et al., 1997), a semi-structured psychiatric interview administered by an old age psychiatrist, to diagnose psychiatric morbidity according to *DSM-IV-TR* criteria. Therefore, the presence of a somatoform disorder is the first indicator of somatization. Second, health anxiety was in both studies assessed with the Whitely Index (Pilowsky, 1967) and used as the second indicator of somatization in the present study. The somatic disease burden was assessed differently in both studies. In the pilot study, the Cumulative Illness Rating Scale for Geriatrics (CIRS-G; Miller et al., 1992) was administered by a geriatrician after a full geriatric assessment. In the OPUS study, the self-report version of the Charlson Index was applied (Charlson, Pompei, Ales, & MacKenzie, 1987; Katz, Chang, Sngha, Fossel, & Bates, 1996).

As a measure of construct validity, Pearson correlations coefficient of both the SCL-90/BSI-53 SOM subscale as well as the PHQ-15 sum score were calculated with both proxies of somatization (Whitely Index, somatoform disorder) expecting a high correlation, and with both measures of somatic disease burden (CIRS-G, Charlson Index) expecting a low correlation. The Pearson r correlation can be interpreted as no or minimal ($0.00 < r < 0.30$), low ($0.30 < r < 0.50$), moderate ($0.50 < r < 0.70$), high ($0.70 < r < 0.90$), or extremely high ($0.90 < r < 1.00$) correlation.

Update and Extension of the Systematic Review by Zijlema et al. (2013)

The systematic review by Zijlema et al. (2013) has been conducted to systematically search and judge all self-report questionnaires for common somatic symptoms, generally

used to assess somatic symptom burden and/or level of somatization. After an update of the literature search, in this article all questionnaires will be evaluated on their suitability for use in an older population (see Instrument Evaluation section).

Search Strategy

Since the literature search of the previous systematic review was conducted until October 2012, we repeated exactly the same literature search in the databases Medline, EMBASE, and PsycINFO from October 2012 until October 1, 2016, to select additional questionnaires. The search term contained a combination of somatoform disorder or synonyms and questionnaire or synonyms and symptoms. For Medline, the following search term was used: (“somatoform disorders/classification” [MeSH Major Topic] OR “somatoform disorders/diagnosis” [MeSH Major Topic] OR “somatoform disorders/epidemiology” [MeSH Major Topic] OR “functional somatic symptoms” [Title/Abstract]) AND (questionnaire [Title/Abstract] OR screen* [Title/Abstract] OR “self-report” [Title/Abstract] OR “index” [Title/Abstract]) AND symptoms. For EMBASE and PsycINFO, comparable search terms were used. The search was conducted without language restrictions.

Screening and Selection Procedures

The first two authors independently screened the retrieved articles. The articles were included if they described the development, evaluation, or review of self-report somatization questionnaires. The questionnaires selected had to include symptoms from more than one symptom cluster; not just symptoms of the gastrointestinal tract or cardiopulmonary system. When the symptom questionnaire was a subscale derived from a larger questionnaire, the symptom subscale had to have been separately validated and used. There were no criteria for the target population of the questionnaire.

Discrepancies between the two researchers were resolved by consensus. Full articles were then obtained for all potentially eligible studies. Based on the full text, articles that still fulfilled the inclusion criteria were included in the review.

Data Extraction

Name of questionnaire, number of items, domains assessed, answering scale, time frame, and target population were extracted for every questionnaire. Table 2 shows an overview of the questionnaires before and after October 2012.

Instrument Evaluation

The evaluation on the suitability of the identified questionnaires for an older population included (a) the number of

items not applicable in older adults (e.g., items on menstrual cycle) and the number of somatic symptoms included that in older persons usually reflect somatic disease burden, (b) the presence of standardized scores (normative data) for older persons, and (c) finally whether the instrument has been applied in an older sample previously.

Although each somatic symptom or sign can be due to somatic disease, some symptoms can be assumed typically for old age and/or common geriatric syndromes. To assess the most common geriatric syndromes, we used the Brief Assessment Tool (BAT), a geriatric assessment tool specifically developed for general practitioners (Senn & Monod, 2015). The BAT aims to identify the following geriatric syndromes: cognitive impairment, mood disorder, gait and balance impairment/falls, visual impairment, hearing impairment, urinary incontinence, malnutrition/loss of weight, and osteoporosis. The first and fourth author, being a clinical psychologist and old age psychiatrist, respectively, assessed which items of each somatization scale corresponded with a geriatric symptom or syndrome as defined by the BAT. In case of disagreement, the last author, an old age psychiatrist, made a decision.

For the third criterion, systematic literature searches were additionally conducted in Medline, EMBASE, and PsycINFO for each questionnaire separately. This was done by search strings combining the name of the questionnaire with the words indicative for a research sample consisting of older persons defined as an age older than 60 years, for example, “older,” “aged,” or “elderly” and in Medline also the MeSH-terms “aged” and “aged, 80 years and over.”

Results

Empirical Tests of the PHQ-15 and SCL-90 SOM

Pearson correlation coefficients of the PHQ-15 sum score and the SCL-90 SOM with the proxies for somatization as well as somatic disease burden are presented in Table 1. Collectively, these results showed that the presence of a somatoform disorder, as proxy for somatization, was neither significantly associated with the sum score of the SCL-90/BSI-53 SOM, nor with the PHQ-15 sum score. Furthermore, the second proxy for somatization, health anxiety, was moderately associated with both measures of somatization (SCL-90/BSI-53 SOM and PHQ-15) as well as moderately associated with the measures of somatic disease burden (CIRS-G and Charlson Index).

Update and Extension of the Systematic Review

The literature research of Zijlema et al. (2013) until October 2012 had resulted in 40 symptom questionnaires. In contrast to Zijlema et al. (2013), however, we considered the

Table 1. Construct Validity of the PHQ Sum Score and SCL-90/BSI-53 Somatization Scale as Measures of Somatization.

	Number of patients	SCL-90/BSI-53 SOM	PHQ-15
<i>Proxies for somatization</i>			
<i>Whitely Index (health anxiety)</i>			
• MUS patients (pilot study)	33	$r = 0.45, p = .009$	$r = 0.38, p = .019$
• MUS patients OPUS study	89	$r = 0.43, p < .001$	n.a.
• MES patients OPUS study	151	$r = 0.49, p < .001$	n.a.
<i>Presence of a somatoform disorder</i>			
• MUS patients (pilot study)	33	$r = 0.12, p = .506$	$r = 0.12, p = .495$
• MUS patients OPUS study	94	$r = 0.16, p = .133$	n.a.
<i>Somatic disease burden</i>			
• CIRS-G ^a (Pilot study)	29	$r = 0.33, p = .083$	$r = 0.28, p = .125$
• Charlson Index (OPUS study) MUS patients	87	$r = 0.44, p < .001$	n.a.
• Charlson Index (OPUS study) MES patients	152	$r = 0.34, p < .001$	n.a.

Note. PHQ-15 = Patient Health Questionnaire 15-item version; SCL-90 SOM = Somatization subscale of the Symptom Checklist 90-item version; BSI-53 SOM = somatization subscale of the Brief Symptom Inventory 53-item version; MUS = Medically Unexplained physical Symptoms; MES = Medically Explained Symptoms; OPUS = Older Persons with medically Unexplained Symptoms (acronym for a study); CIRS-G = Cumulative Illness Rating Scale for Geriatrics.

^aCIRS-G total score calculated without psychiatric disease.

two abbreviated versions of the somatization scale of the SCL-90 (SCL-90 SOM), that is, the BSI-18 SOM (six items) and the BSI-53 SOM (seven items), as separate questionnaires. Therefore, 41 symptom questionnaires were available based on Zijlema et al. (2013). The extended literature search from October 2012 until October 1, 2016, retrieved a total of 631 hits (Medline, $n = 187$, EMBASE, $n = 157$, PsycINFO, $n = 287$), including 75 duplicates. We excluded 436 studies identified on title and abstract. After full text screening, we excluded 114 studies because they were not about somatization ($n = 34$), were not a self-report questionnaire ($n = 3$), or were about questionnaires already included ($n = 77$). A total of six articles were additionally included in the review, describing five new questionnaires. This resulted in a final number of 46 questionnaires for the present review (Table 2).

Evaluation of Questionnaires

Table 3 shows all questionnaires identified. While 8 out of 46 (17%) questionnaires have normative data for older persons, only the Brief Symptom Screen (BSS) was specifically validated in an older sample. The additional questionnaire-specific searches in PubMed, EMBASE, and PsycINFO revealed that only 20 out of 46 (43%) questionnaires were ever used in an older population.

The identification of items overlapping with common geriatric syndromes revealed that all items identified by the first rater (first author) were also identified by the second rater (fourth author). Discrepancies could be clustered in three groups. The first group of items included fatigue, tiredness, loss of energy, and feeling weak; the second group of items loss of appetite and weight loss; and finally

the third group (an individual item) was insomnia (sleep onset). The third rater (last author) concluded that these symptoms could all be classified as somatic symptoms of depression and therefore overlap with depression. Taken this decision into account, the median proportion of items overlapping with common geriatric syndromes, plus the number of items not applicable for old age, was 25%. The variability between questionnaire was large, with a range from 0% for the Schedule for Evaluating Persistent Symptoms (SEPS) through as high as 87% for the Health-49. For the PHQ-15, SCL-90 SOM, and SCL-53 SOM, these percentages were 33%, 25%, and 14%, respectively.

Discussion

The PHQ-15 and the SCL-90/BSI-53 SOM both had low correlations with the severity of health anxiety among MUS patients, while neither the PHQ-15 nor the SCL-90/BSI-53 SOM correlated with the presence of a somatoform disorder according to *DSM-IV-TR* criteria. These findings did not match with our expectations of a moderate to high correlation between both symptom scales and both proxies of somatization. Even more important to note, the PHQ-15 and the SCL-90/BSI-53 SOM were both correlated with measures of somatic disease burden with an effect size comparable to their association with health anxiety. Although the findings with respect to the PHQ-15 need replication in a larger sample, collectively, these findings indicate that both scales do not perform well as indices of somatization in older persons.

An update of the literature search conducted by Zijlema et al. (2013) resulted in five additional self-report somatic screenings lists, namely the BSS (Ritchie et al., 2013), the

Table 2. Overview of the 46 Somatization Questionnaires and Their Properties.

Questionnaire	Items	Domain assessed	Scale	Time frame	Target population
4 DSQ	16	Somatization	5 Categories: no to very often or constantly	Past week	Primary care patients
ASR	11	Somatic complaints	3 Categories: not true to very true or often true	Past 6 months	Adults
BDS Checklist	25	BDS; pattern of symptoms rather than a simple symptom count (based on SCAN interview)	5 Categories: not at all to a lot	Past month	Patients
BSI-18 SOM	6	Somatization	5 Categories: not at all to a lot	Past week	Adolescents and adults
BSI-53 SOM^a	7	Somatization	5 Categories: not at all to a lot	Past week	Adolescents and adults
BSI	44	Somatic symptoms associated with anxiety and depression	3 Categories: symptoms absent to present on more than 15 days during the past month	Past month	Patients
BSS^a	10	Somatic complaints	Yes/no	Past 4 weeks	Community dwelling older adults
Cambodian SSI	23	Somatic symptoms and cultural syndromes: with a 12-item somatic subscale and an 11-item syndrome subscale	5 Categories: not at all to extremely	Past month	Traumatized Cambodian Refugees
C-PSC	12	Psychosomatic symptoms	Frequency: 5 categories: not a problem to every day. Severity: 5 categories: not a problem to very, very bad		Children
CSI	36	Intensity of somatic complaints	4 Categories: not at all to a whole lot	Past 2 weeks	Children
FBL	78	Somatic complaints	Frequency: 5 categories: almost every day to almost never. Intensity: 5 categories: very strongly to insensitive	Lately	
GBB-24	24	Physical complaints	5 Categories: never to severe		Patients and general population
GSL	37	Psychosomatic stress symptoms	4 Categories: never to constantly		
Health-49	7	Somatoform complaints	5 Categories: not at all to very much		
ICD-10 symptom list	14	Somatization disorder	Yes/no	Past 2 years	Patients
Kellner's SQ	17	Somatic symptoms	Yes/no or true/false	Past week to day	Patients and general population
Malaise Inventory	8	Psychiatric morbidity	Yes/no	No specific time frame, focus on recent state	
Manu	5	Somatization disorder	Yes/no		
MSPQ	13	Heightened somatic and autonomic awareness	4 Categories: not at all to extremely, could not have been worse	Past week	Specially for chronic backache patients
NSS	6	Nonspecific symptoms for nonpsychotic morbidity	Present/not present	At least 3 months	Patients
Othmer and DeSouza	7	Somatization disorder	Yes/no	Lifetime	General population

(continued)

Table 2. (continued)

Questionnaire	Items	Domain assessed	Scale	Time frame	Target population
PHQ	14	Somatic symptoms	Items 1-11, 7 categories: <i>not at all to all of the time</i> ; items 12-13, 7 categories: <i>0 times to 7+ times</i> ; item 14, 7 categories: <i>1 day to 7+ days</i>		Staff members of a hospital
PHQ-15	15	Probable somatoform disorders	3 Categories: <i>not at all to bothered a lot</i>	Past month	Primary care patients
PILL	54	Common physical symptoms and sensations	5 Categories: <i>never or almost never to more than once every week</i>	Lifetime	
PSC-17	17	Psychosomatic symptoms	Frequency 5 categories: <i>daily to not a problem</i> ; intensity; 5 categories: <i>extremely bothersome to not a problem</i>	Past week	Primary care patients
PSC-51	51	Somatization	4 Categories: <i>not at all to most of the time</i>	Past week	Primary care patients
PSS	35	Psychosomatic symptoms	Frequency: 4 categories: <i>never to almost every day</i> ; disturbance; 3 categories; <i>none to strong</i>	Past 3 months	Children and adolescents
PVPS	14	Somatization	3 Categories: <i>never occurred to frequently occurred</i>	Past month	People of Vietnamese origin
RPSQ	26	Somatization in IBS patients	4 Categories: <i>not at all to most of the time</i>	Past month	IBS patients
R-SOMS-2	29	Somatization	Yes/no	Past 2 years	Primary care patients
QUISS-P ^a	18	Severity of somatoform disorders	5 Categories; mixed categories	Usually	Inpatient and outpatients form psychiatric and psychosomatic hospitals
SCI	22	Various physical symptoms	Frequency, 5 categories: <i>never to daily</i> ; intensity, 5 categories: <i>no problems to extremely troublesome</i>	Past month	General population
SCL-11	11	Common somatic complaints	5 Categories: <i>almost never to quite often</i>	Past month	Children
SCL-90 SOM	12	Somatization	5 Categories: <i>not at all to extremely psychiatric</i>	Past week	Medical outpatients/ general population
SEPS ^a	Section 2, 9 items	Medically unexplained symptoms	4 Mixed categories	Lifetime	Medical patients
SHC	29	Subjective health complaints	Severity, 4 categories: <i>not at all to serious</i> ; duration: number of days	Past month	General population
SOMS-7	53	Intervention effects in somatoform disorders	5 Categories: <i>not at all to very severe</i>	Past week	Primary care patients
SSEQ ^a	15	Psychological processes in somatoform disorder	6 Categories from <i>never to always</i>	Usually	Psychosomatic inpatients
SSI	35	Somatization	Yes/no	Lifetime	Primary care patients

(continued)

Table 2. (continued)

Questionnaire	Items	Domain assessed	Scale	Time frame	Target population
SSS-8^a	8	Presence and severity of somatic symptoms	5 Categories from <i>not at all</i> to <i>very much</i>	Past 7 days	General population
SQ-48	7	Somatization	5 Categories: <i>never</i> to <i>very often</i>	Past week	Clinical and nonclinical population
Swartz	11	Symptoms that potentially predict a diagnosis of DID/DSM-3 somatization disorder	Yes/no	Lifetime	General population
Syrian Symptom Checklist	19	Psychosomatic symptoms; diagnose individuals, follow-up treatment, evaluate treatment intervention	4 Categories: <i>never</i> to <i>always</i>	Past few weeks	
WHO-SSD	12	Somatiform disorder	Yes/no	Past 6 months	General population
YSR	9	Somatic complaints	3 Categories: <i>not true</i> to <i>very true</i> or <i>often true</i>	Past 6 months	11- to 18-year-olds
Von Zerssen	24	Somatic complaints	4 Categories: <i>not at all</i> to <i>strong</i>		

Note. 4 DSQ = Four-Dimensional Symptom Questionnaire (Terluin et al., 2006); ASR = Adult Self-Report (Achenbach & Rescorla, 2003); BDS Checklist = Bodily Distress Syndrome Checklist (Budtz-Lilly, Fink, Ørnbøl, Christensen, & Rosendal, 2015); BSI = Bradford Somatic Inventory (Mumford, 1989; Mumford et al., 1991); BSI-18 SOM = Brief Symptom Inventory 18-item version somatization scale (Galdon et al., 2008); BSI-53 SOM = Brief Symptom Inventory 53-item version somatization scale (Derogatis & Melisaratos, 1983); BSS = Brief Symptom Screen (Ritchie et al., 2013); Cambodian SSI = Cambodian Somatic Symptom and Syndrome Inventory (Hinton, Kredlow, Bui, Pollack, & Hofmann, 2012); C-PSC = Children's Psychosomatic Symptom Checklist (Garber, Walker, & Zeman, 1991); Wisniewski, Naglieri, & Mulick, 1988); CSI = Children's Somatization Inventory (Walker, Garber, & Greene 1991); FBL = Freiburger Beschwerden Liste (Freiburg Complaint List, Fahrenberg, 1995); GBB-24 = Giessener Beschwerdebogen (Giessen Subjective Complaints List; Brähler, Schumaner, & Brähler, 2000); GSL = Goldberg Symptom List (Herman & Lester, 1994); Health-49 = Hamburger Module zur Erfassung allgemeiner Aspekte psychosozialer Gesundheit für die therapeutische Praxis (Hamburger modules to measure general aspects of psychosocial health for therapeutic praxis; Rabung et al., 2009); ICD-10 Symptom List = International Classification of Diseases-10 Symptom List (Khoo, Mathers, McCarthy, & Low, 2012; WHO, 1993); Kellner's SQ = Kellner's Symptom Questionnaire (Kellner, 1987); Malaise Inventory (Rodgers, Pickles, Power, Collishaw, & Maughan, 1999); Manu, Lane, Matthews, and Escobar (1989); MSPQ = Modified Somatic Perception Questionnaire (Main, 1993); NISS = Nonspecific Symptom Screen (Srinivasan & Suresh, 1991); PHQ = Physical Health Questionnaire (Schat, Kelloway, & Desmarais, 2005); Spence, Helmreich, & Pred, 1987); PHQ-15 = Patient Health Questionnaire (Kroenke, Spitzer, deGruy, & Swindle, 1998; Kroenke, Spitzer, & Williams, 2002); PILL = Pennebaker Inventory of Limbic Languidness (Pennebaker, 1982); PSC-17 = Psychosomatic Symptom Checklist (Atanasio, Andrasik, Blanchard, & Arena, 1984); PSC-51 = Physical Symptom Checklist (de Waal, Arnold, Spinhoven, Eekhof, & van Hemert, 2005); PSS = Uppitnika Pshhosomatskih Simptomov (Psycho-somatic Symptoms Questionnaire; Vulić-Prtorić, 2005); PVPS = Phan Vietnamese Psychiatric Scale (Nettleton, 2006); QUISS = the Quantification Inventory for Somatoform Syndromes (Wedekind, Bandelow, Fentzahn, Trümper, & Rütter, 2007); RPSQ = Recent Physical Symptoms Questionnaire (MacLean, Palsson, Turner, & Whitehead, 2012); R-SOMS-2 = Revised Screening for Somatoform Symptoms (Fabiao, Silva, Barbosa, Fleming, & Rief, 2010); SCI = Somatic Symptom Checklist Instrument (Bohman et al., 2012); SCL = Somatic Complaint List (Jellesma, Rief, & Terwogt, 2007); SCL-90 SOM = Symptom Checklist, 90-item version somatization scale (Derogatis, Lipman, Rickels, Uhlenhuth, & Covi, 1974); **SEPS: Schedule for Evaluating Persistent Symptoms (Tyrer et al., 2012)**; **SSEQ = Somatic Symptoms Experiences Questionnaire (Herzog et al., 2014)**; **SSS-8 = the Somatic Symptom Scale-8 (Gierk et al., 2014)**; SHC = Subjective Health Complaints Inventory (Ursin, Endresen, & Ursin, 1988); SOMS-7 = Screening for Somatoform Symptoms (Rief & Hiller, 2003, 2008); SSC = Syrian Symptom Checklist (Rudwan, 2000); SSI = Somatic Symptom Index; SQ-48 = Symptom Questionnaire 48 (Carlier, Schulte-van Maaren, Wardenaar, & Zitmann, 2012); WHO-SSD = World Health Organization (WHO)—Screening for Somatoform Disorders (Phillips, Fallon, & King, 2008); YSR = Youth Self-Report (Giedhill & Garraida, 2006); Von Zerssen (Ladwig, Marten-Mittag, Lacruz, Henningsen, & Creed, 2010); IBS = irritable bowel syndrome.

^aLists derived from literature search after October 2012 are bold.

Table 3. The 46 Somatic Screening Lists and Their Properties for the Old Aged.

Questionnaire	Overlapping items between symptom questionnaire and geriatric syndrome	Proportion, n/N (%), of items overlapping with commonly geriatric syndromes or items that are not applicable for the old aged	Normative data for older persons	Older adults included in the study
4 DSQ	Dizziness or light-headed (gait and balance) Painful muscles (osteoporosis) Neck pain (osteoporosis) Back pain (osteoporosis) Blurred vision (visual impairment) I feel dizzy or light headed (gait and balance) Problems with eyes (visual impairment)	5/16 (31)	—	Koorevaar, Terluin, van 't Riet, Madden, and Bulstra (2016); n = 200; age = 15-85, shoulder patients
ASR	I feel dizzy or light headed (gait and balance) Problems with eyes (visual impairment)	2/11 (18)	—	—
BDS Checklist	Pain in arms or legs (osteoporosis) Muscular aches or pain (osteoporosis) Pain in the joints (osteoporosis) Excessive fatigue (malnutrition/mood disorder) Impairment of memory (cognitive impairment) Dizziness (gait and balance) Faintness or dizziness (gait and balance)	6/25 (24)	—	Budtz-Lilly et al. (2015); n = 1,356; age = 18-95; primary care patients Budtz-Lilly et al. (2015); n = 2,480; age = 26-71; primary care patients
BSI-18 SOM	Faintness or dizziness (gait and balance)	1/6 (17)	—	Asner-Self, Schreiber, and Marotta (2006); n = 100; age = 18-80; American volunteers Tanji et al. (2008); n = 96; age = 57-75; patients with morbus Parkinson and their spouses Petkus, Gum, King, Kallimanis, and Wetherell (2009); n = 136; trauma exposed older adults Wetherell et al. (2010); n = 54; age = 70-78; patients with generalized anxiety disorder Campo, Agarwal, LaStrayo, O'Connor, and Pappas (2014); n = 40; age = 58-93; prostate cancer survivors Cohen (2014); n = 321; aged > 60; cancer patients Russell et al. (2015); n = 152; mean age = 64; colorectal cancer survivors Ritsner, Ponizovsky, Kurs, and Modai (2000); n = 996; age = 18-87; Jewish immigrants Pietrzak et al. (2005); n = 48; age = 60+; patients with pathological gambling Pietrzak (2006); n = 21; age = 60+; patients with pathological gambling Zweig and Turkel (2007); n = 129; age = 63-87; community dwelling elderly Klein, Lezotte, Heltsh, Fauerbach, and Holavanahalli (2011); n = 737; age 55+; patient with brain injury van Noorden et al. (2012); n = 892; MUS patients referred to geriatric outpatient psychiatry Videler, Rossi, Schoevaars, van der Feltz-Cornelis, and van Alphen (2014); n = 31; age = 60-78; patients with personality disorder Pereira, Martins, Alves, and Canavarro (2014); n = 185; aged 50+; HIV-infected patients Dijk, Voshaar, Lucassen, Comijs, and Hanssen (2015); n = 153; age = 60+; patients with MUS Andersen et al. (2015); n = 1,000; age 60+; patients with alcohol use disorder
BSI-53 SOM	Faintness or dizziness (gait and balance)	1/7 (14)	Hale, Hinz, and Brähler (1984); n = 498; mean age = 74; comparison between adults and older adults Hale and Cochran (1992); n = 220; age > 65; comparison between four age cohorts Chester (2001); n = 498; age > 65; raw score means for independent living older adults	

(continued)

Table 3. (continued)

Questionnaire	Overlapping items between symptom questionnaire and geriatric syndrome	Proportion, n/N (%), of items overlapping with commonly geriatric syndromes or items that are not applicable for the old aged	Normative data for older persons	Older adults included in the study
BSI	Have you had pain or tension in your neck and shoulder? (osteoporosis) Has there been darkness or mist in front of your eyes (visual impairment) Have you felt aches or pains all over the body? (osteoporosis) Have you been feeling tired, even if you are not working? (mood disorder) Pain in your legs (osteoporosis) Dizzy (gait and balance) Feeling tired (mood disorder) Balance dizziness (gait and balance) Daily pain (osteoporosis) Poor appetite (mood disorder) Anhedonia (mood disorder) Dizziness (gait and balance) Standing up and feeling dizzy (gait and balance)	6/44 (14)	—	Saeed, Mubbashar, Dogar, Mumford, and Mubbashar (2001): n = 664; age = 18-80; Rural community in Pakistan Kahn and Taj (2011): n = 200; age = 18-80; migrated Pakistan men and women
BSS	Blurred vision (visual impairment) Tinnitus (hearing impairment) Neck soreness (osteoporosis) Sore arms and legs (osteoporosis) Poor appetite (mood disorder) Backaches (osteoporosis) Sad (mood disorder) Feel stiff all over (osteoporosis) Feel dizzy (gait and balance) Eye pain when reading (visual impairment) Blindness (visual impairment) Fainting (gait and balance) Memory loss (cognitive impairment) Blurred vision (visual impairment) Deafness (hearing impairment) Dizziness (gait and balance) Pain in arms and legs (osteoporosis) Pain in joints (osteoporosis) Back pain (osteoporosis) Trouble walking (gait and balance) Low energy (malnutrition, mood disorder) Ermuden Sie schnell (mood disorder) Haben Sie appetitmangel (mood disorder, malnutrition)	5/10 (50)	Ritchie et al. (2013): n = 1,000; age = 65+; community dwelling older adults in Alabama	—
Cambodian SSI	Blurred vision (visual impairment) Tinnitus (hearing impairment) Neck soreness (osteoporosis) Sore arms and legs (osteoporosis) Poor appetite (mood disorder) Backaches (osteoporosis) Sad (mood disorder) Feel stiff all over (osteoporosis) Feel dizzy (gait and balance) Eye pain when reading (visual impairment) Blindness (visual impairment) Fainting (gait and balance) Memory loss (cognitive impairment) Blurred vision (visual impairment) Deafness (hearing impairment) Dizziness (gait and balance) Pain in arms and legs (osteoporosis) Pain in joints (osteoporosis) Back pain (osteoporosis) Trouble walking (gait and balance) Low energy (malnutrition, mood disorder) Ermuden Sie schnell (mood disorder) Haben Sie appetitmangel (mood disorder, malnutrition)	7/23 (30)	—	Friberg et al. (2007): n = 61,320; age = 45-74; patients with oropharyngeal carcinomas
C-PSC	Blurred vision (visual impairment) Tinnitus (hearing impairment) Neck soreness (osteoporosis) Sore arms and legs (osteoporosis) Poor appetite (mood disorder) Backaches (osteoporosis) Sad (mood disorder) Feel stiff all over (osteoporosis) Feel dizzy (gait and balance) Eye pain when reading (visual impairment) Blindness (visual impairment) Fainting (gait and balance) Memory loss (cognitive impairment) Blurred vision (visual impairment) Deafness (hearing impairment) Dizziness (gait and balance) Pain in arms and legs (osteoporosis) Pain in joints (osteoporosis) Back pain (osteoporosis) Trouble walking (gait and balance) Low energy (malnutrition, mood disorder) Ermuden Sie schnell (mood disorder) Haben Sie appetitmangel (mood disorder, malnutrition)	5/12 (42)	—	—
CSI	Blurred vision (visual impairment) Tinnitus (hearing impairment) Neck soreness (osteoporosis) Sore arms and legs (osteoporosis) Poor appetite (mood disorder) Backaches (osteoporosis) Sad (mood disorder) Feel stiff all over (osteoporosis) Feel dizzy (gait and balance) Eye pain when reading (visual impairment) Blindness (visual impairment) Fainting (gait and balance) Memory loss (cognitive impairment) Blurred vision (visual impairment) Deafness (hearing impairment) Dizziness (gait and balance) Pain in arms and legs (osteoporosis) Pain in joints (osteoporosis) Back pain (osteoporosis) Trouble walking (gait and balance) Low energy (malnutrition, mood disorder) Ermuden Sie schnell (mood disorder) Haben Sie appetitmangel (mood disorder, malnutrition)	11/36 (31)	—	—
FBL	Blurred vision (visual impairment) Tinnitus (hearing impairment) Neck soreness (osteoporosis) Sore arms and legs (osteoporosis) Poor appetite (mood disorder) Backaches (osteoporosis) Sad (mood disorder) Feel stiff all over (osteoporosis) Feel dizzy (gait and balance) Eye pain when reading (visual impairment) Blindness (visual impairment) Fainting (gait and balance) Memory loss (cognitive impairment) Blurred vision (visual impairment) Deafness (hearing impairment) Dizziness (gait and balance) Pain in arms and legs (osteoporosis) Pain in joints (osteoporosis) Back pain (osteoporosis) Trouble walking (gait and balance) Low energy (malnutrition, mood disorder) Ermuden Sie schnell (mood disorder) Haben Sie appetitmangel (mood disorder, malnutrition) Schachegefühl (malnutrition) Müdigkeit (malnutrition) Nackenschmerzen (osteoporosis) Schulterschmerzen (osteoporosis) Kreuzschmerzen (osteoporosis) Schmerzen in den Armen (osteoporosis) Schmerzen in den Beinen (osteoporosis)	9/78 (12)	Fahrenberg (1995): n = 2,070; age > 70; general population	—

(continued)

Table 3. (continued)

Questionnaire	Overlapping items between symptom questionnaire and geriatric syndrome	Proportion, n/N (%), of items overlapping with commonly geriatric syndromes or items that are not applicable for the old aged	Normative data for older persons	Older adults included in the study
GBB-24	Gliederschmerzen (osteoporosis) Ruckenschmerzen (osteoporosis) Nackenschmerzen (osteoporosis) Müdigkeit (mood disorder)	4/24 (17)	Gunzelmann, Goldstein, Sirockman, and Green (1996): n = 764; age > 60; general population	Gunzelmann, Goldstein, Sirockman, and Green (2002): n = 593; age > 60; general population Gunzelmann, Hinz, and Brähler (2006): n = 630; age = 61-95; GBB-24 used as construct validity instrument with Nottingham Health Profile Stankuniene et al. (2012): n = 624; age = 60-84; general population Csoff, Macassa, and Lindert (2010): n = 593; age = 60-84; immigrants Germany Valdearenas, Torres-Gonzalez, de Dios Luna, and Cervilla (2012): n = 562; age = 60-84; nondemented community-dwelling elderly Stankunas et al. (2013): n = 4,467; age = 60-84; association between somatic complaints and educational level
GSL	Lower back pain (osteoporosis) Fatigue (mood disorder) Angry feelings (mood disorder) Sleep onset insomnia (mood disorder) Worrisome thoughts (mood disorder) Early morning awakenings (mood disorder) Loss of appetite (mood disorder) Neck, shoulder muscle aches (osteoporosis) Periods of depression (mood disorder) Sadness (mood disorder) Back pain (osteoporosis) Thoughts that you would rather be dead (mood disorder) Lack of interest (mood disorder) Pain in muscles or joints (osteoporosis) Feeling of hopelessness (mood disorder) Pain in the limbs, extremities, or joints (osteoporosis)	9/37 (24)	—	—
Health-49		6/7 (86)	—	Rabung et al. (2009): n = 1,548; clinical samples; n = 5,630 primary care patients; all ages
ICD-10 Symptom list		1/14 (7)	—	Schafer, Hansen, Schon, Hofels, and Altiner (2012): n = 50,786; age > 70; primary care
Kellner's SQ	Poor appetite (mood disorder) Muscle pain (osteoporosis)	2/17 (12)	—	Callixte et al. (2015): n = 187; age > 60; neurological patients Riccieri, Del Basso, Tomba, Offidani, and Prignano (2014): n = 70; all ages; psoriasis patients
Malaise Inventory	Tired (mood disorder) Depressed (mood disorder) Early waking (mood disorder) Poor appetite (mood/malnutrition)	5/8 (63)	—	Quine and Charnley (1987): n = 226; age unknown, but carers for elderly > 65 (often elderly themselves) Grant, Nolan, and Ellis (1990): n = 125; age unknown, spouses caring of partner aging 65 or older
Manu	Blurred vision (visual impairment)	1/5 (20)	—	—
MSPQ	Dizziness (gait and balance) Blurred vision (visual impairment) Muscles in neck aching (osteoporosis)	3/13 (23)	—	Staerkle et al. (2004): n = 388; age = 18-87; low back pain patients Roh et al. (2008): n = 111; age = 45-83; Parkinson patients Donaldson et al. (2011): n = unknown; age = 55-75; patients with neck and low back pain Havakeshian and Mammion (2013): n = 159; mean age 65; spinal surgery patients
NSS	Forgetfulness (cognitive impairment) Giddiness/dizziness (gait and balance) General aches and pain (osteoporosis) Fatigability (malnutrition) Feeling weak (malnutrition)	5/6 (83)	—	—

(continued)

Table 3. (continued)

Questionnaire	Overlapping items between symptom questionnaire and geriatric syndrome	Proportion, n/N (%), of items overlapping with commonly geriatric syndromes or items that are not applicable for the old aged	Normative data for older persons	Older adults included in the study
Othmer and DeSouza	Amnesia (cognitive impairment) Painful extremities (osteoporosis) Blindness (visual impairment) Dysmenorrhea	4/7 (58)	—	—
PHQ	Difficulty getting to sleep (mood disorder) Woken up during the night (mood disorder) How often has your sleep been peaceful and disturbed (mood disorder)	3/14 (21)	—	—
PHQ-15	Back pain (osteoporosis) Pain in your arms, legs, or joints (osteoporosis) Dizziness (gait and balance) Feeling tired or having low energy (mood disorder/malnutrition) Menstrual cramps or other problems with your periods	5/15 (33)	Nordin, Palmquist, and Nordin (2013); Swedish population norms for ages 70 to 79	Sloane, Hartman, and Mitchell (1994): n = 65; age > 60; patients with chronic dizziness Sha et al. (2005): n = 3,498; age > 60; validity of symptoms in predicting hospitalization and mortality Montalban, Comas, and Garcia-Garcia (2010): n = 3,362; age = 18-90; outpatient psychiatric patients Jeong et al. (2014): n = 2,100; age > 60; relationship somatic symptoms—depression Qian, Rem, Yu, He, and Li (2014): n = 1,329; age = 37-71; general hospital Graham, Balard, and Pak (1997): n = 109; 52% age > 65; informal carers for dementia patients
PILL	ringing in ears (hearing impairment) Back pain (osteoporosis) Dizziness (gait and balance) Stiff joints (osteoporosis)	4/54 (7)	—	—
PSC-17	Fatigue (malnutrition, mood disorder) Backaches (osteoporosis) Depression (mood disorder) General stiffness (osteoporosis) Dizziness (gait and balance)	5/17 (29)	—	—
PSC-51	Feeling tired or having low energy (malnutrition/mood disorder) Easily fatigued without exertion (mood disorder) Dizziness (gait and balance) Forgetfulness (cognitive impairment) Muscle aches or soreness (osteoporosis) Deafness (hearing impairment) Double vision or blurred vision (visual impairment) Blindness (visual impairment) Loss of appetite (mood disorder) Weight loss (malnutrition) Joint pain (osteoporosis) Back pain (osteoporosis)	12/51 (24)	—	—

(continued)

Table 3. (continued)

Questionnaire	Overlapping items between symptom questionnaire and geriatric syndrome	Proportion, n/N (%), of items overlapping with commonly geriatric syndromes or items that are not applicable for the old aged	Normative data for older persons	Older adults included in the study
PSS	Dizziness (gait and balance) Pain in the back (osteoporosis) Lack of energy (mood disorder) Pain in joints (osteoporosis) Pain in arms and legs (osteoporosis) Loss of balance (gait and balance) Double vision (visual impairment) Blurred vision (visual impairment) Sudden loss of vision (visual impairment) Sudden loss of hearing (hearing impairment) Sudden loss of memory (cognitive impairment) Loss of appetite (mood disorder/ malnutrition)	12/35 (34)	—	—
PVPS	Dizzy spells (gait and balance) Tired eyes, sore eyes, or flashy lights (visual impairment) Worn out or low in energy (mood disorder/malnutrition) Painful joints (osteoporosis) Increasingly tired day after day (mood disorder)	5/14 (36)	—	Phan (2004): n = 180; age = 16-75; primary care
QUISS-P	Back pain (osteoporosis) Pain in arms or legs (osteoporosis) Pain in the joints (osteoporosis) Loss of memory (cognitive impairment) Disturbance in balance and coordination (gait and balance) Deafness (hearing impairment) Blindness (visual impairment) Tiredness (mood disorder) Loss of appetite (mood disorder) Dizziness (gait and balance) Back pain (osteoporosis) Muscles aches (osteoporosis) Poor appetite (mood disorders) Constant tiredness (mood disorder) Joint pain (osteoporosis) Pain in the arms/legs (osteoporosis) Impaired coordination in balance (gait and balance) Amnesia (cognitive impairment) Excessive tiredness (mood disorder) Pain in arms and legs (gait and balance) Dizziness (gait and balance) Poor appetite (mood disorder) Dizzy (gait and balance) Tired (mood disorder/malnutrition) Pain in arms and legs (osteoporosis)	9/18 (50)	—	Wedekind et al. (2007): n = 96; age = 18-75; inpatients and outpatients from psychiatric and psychosomatic hospitals
RPSQ	Dizziness (gait and balance) Back pain (osteoporosis) Muscles aches (osteoporosis) Poor appetite (mood disorder) Constant tiredness (mood disorder) Joint pain (osteoporosis)	5/26 (19)	—	—
R-SOMS-2	Pain in the arms/legs (osteoporosis) Impaired coordination in balance (gait and balance)	5/29 (17)	—	—
SCI	Excessive tiredness (mood disorder) Pain in arms and legs (gait and balance) Dizziness (gait and balance)	3/22 (14)	—	—
SCL-11	Poor appetite (mood disorder) Dizzy (gait and balance) Tired (mood disorder/malnutrition) Pain in arms and legs (osteoporosis)	3/11 (27)	—	—

(continued)

Table 3. (continued)

Questionnaire	Overlapping items between symptom questionnaire and geriatric syndrome	Proportion, n/N (%), of items overlapping with commonly geriatric syndromes or items that are not applicable for the old aged	Normative data for older persons	Older adults included in the study
SCL-90 SOM	Fatiness or dizziness (gait and balance) Pain in lower back (osteoporosis) Soreness of your muscle (osteoporosis)	3/12 (25)	Creed et al. (2011); n = 44; age = 66-96; geriatric adult population	Hassel et al. (2007); n = 125; age = 60+; the correlation between OHRQoL and somatization older patients from primary geriatric medical hospital
SEPS				
SHC	Shoulder pain (osteoporosis) Neck pain (osteoporosis) Upper back pain (osteoporosis) Arm pain (osteoporosis) Low back pain (osteoporosis) Sadness/depression (mood disorder) Tiredness (mood disorder) Dizziness (gait and balance)	0/9 (0) 8/29 (28)	Thygesen, Lindstrom, Saevareid, and Engedal (2009); n = 242; age > 75 >; older adults; community dwelling and receiving in home care	Lhiebeak, Eriksen, and Ursin (2002); n = 1,240; age = 1.5-84; general population
SOMS-7	Back pain (osteoporosis) Joint pain (osteoporosis) Pain in the legs and arms (osteoporosis) Loss of appetite (mood disorder/malnutrition) Impaired coordination of balance (gait and balance) Double vision (visual impairment) Blindness (visual impairment) Deafness (hearing impairment) Amnesia (cognitive impairment) <i>Painful menstruation</i> <i>Irregular menstruation</i> <i>Excessive menstrual bleeding</i> <i>Frequent vomiting during pregnancy</i>	13/53 (25)	—	Sack, Boroske-Leiner, and Lahmann (2010); n = 240; age = 18-74; outpatients of the department for psychosomatic medicine and psychotherapy in Germany Kliem et al. (2014); n = 2,434; age = 14-84; general population
SSEQ				
SSI	— Pain in the extremities (osteoporosis) Back pain (osteoporosis) Joint pain (osteoporosis) Amnesia (cognitive impairment) Deafness (hearing impairment) Double vision (visual impairment) Blurred vision (visual impairment) Blindness (visual impairment) Fainting of loss of consciousness (gait and balance) <i>Painful menstruation</i> <i>Irregular menstrual periods</i> <i>Excessive menstrual bleeding</i> <i>Vomiting throughout pregnancy</i>	0/15 (0) 13/35 (37)	— —	— —

(continued)

Table 3. (continued)

Questionnaire	Overlapping items between symptom questionnaire and geriatric syndrome	Proportion, n/N (%), of items overlapping with commonly geriatric syndromes or items that are not applicable for the old aged	Normative data for older persons	Older adults included in the study
SSS-8	Back pain, (osteoporosis) Pain in arms/legs or joints (osteoporosis) Dizziness (gait and balance) Feeling tired (mood disorder) or low energy (malnutrition loss of weight)	4/8 (50)	Gierk et al. (2014): n = 190; age = 14-91 (normgroup > 79); general population	—
SQ-48	I felt dizzy or lightheaded (gait and balance)	1/7 (14)	—	—
Swartz	Pain in extremities (osteoporosis)	2/11 (20)	—	Swartz et al. (1986): n = 900; age > 60; general population
Syrian Symptom Checklist	I feel dizzy (gait and balance) I suffer tiredness (mood disorder) I feel fatigued (mood disorder) I feel exhausted (mood disorder) I feel lethargic (mood disorder)	5/19 (26)	—	—
WHO-SSD	Back pain (osteoporosis) Dizziness (gait and balance) Feelings of muscles and aches (osteoporosis) Persistent fatigue after minor mental or physical effort (mood disorder) <i>Irregular menstrual periods</i> <i>Excessive menstrual bleeding</i>	6/12 (50)	—	—
YSR	I feel dizzy (gait and balance) I feel tired (mood disorder) Aches or pain (osteoporosis) Problems with eyes (visual impairment)	4/9 (44)	—	—
Von Zerssen	Kreuz oder Ruckenschmerzen (osteoporosis) Schwindelgefühl (gait and balance) Nackten oder Schulterschmerzen (osteoporosis) Gewichtsaufnahme (malnutrition)	4/24 (17)	—	—

Note. 4 DSQ = Four-Dimensional Symptom Questionnaire; ASR = Adult Self-Report; BDS Checklist = Bodily Distress Syndrome Checklist; BSI = Bradford Somatic Inventory; BSI-18 SOM = Brief Symptom Inventory 18-item version somatization scale; BSI-53 SOM = Brief Symptom Inventory 53-item version somatization scale; **BSS = Brief Symptom Screen**; Cambodian SSI = Cambodian Somatic Symptom and Syndrome Inventory; C-PSC = Children's Psychosomatic Symptom Checklist; CSI = Children's Somatization Inventory; FBL = Freiburger Beschwerden Liste (Freiburg Complaint List); GBB-24 = Giessen Beschwerdebogen (Giessen Subjective Complaints List); GSL = Goldberg Symptom List; Health-49 = Hamburger Module zur Erfassung allgemeiner Aspekte psychosozialer Gesundheit für die therapeutische Praxis (Hamburger modules to measure general aspects of psychosocial health for therapeutic praxis); ICD-10 Symptom List = International Classification of Diseases-10 Symptom List; Kellner's SQ = Kellner's Symptom Questionnaire; MSPQ = Modified Somatic Perception Questionnaire; NSS = Nonspecific Symptom Screen; PHQ = Physical Health Questionnaire; PHQ-15 = Patient Health Questionnaire; PILL = Pennebaker Inventory of Limbic Languidness; PSC-17 = Psychosomatic Symptom Checklist; PSC-51 = Physical Symptom Checklist; PSS = Uptinika Psihosomatskih Simptoma (Psychosomatic Symptom Questionnaire); PVPS = Phan Vietnamese Psychiatric Scale; **QUISS = the Quantification Inventory for Somatoform Syndromes**; RPSQ; Recent Physical Symptoms Questionnaire; R-SOMS-2 = Revised Screening for Somatoform Symptoms; SCI = Somatic Symptom Checklist; Instrument; SCL = Somatic Complaint List; SCL-90 SOM = Symptom Checklist 90-item version somatization scale; **SEPS = Schedule for Evaluating Persistent Symptoms**; **SEEQ = Somatic Symptoms Experiences Questionnaire**; **SSS-8 = the Somatic Symptom Scale-8**; SHC = Subjective Health Complaints Inventory; SOMS-7 = Screening for Somatoform Symptoms; SSC = Syrian Symptom Checklist; SSI = Somatic Symptom Index; SQ-48 = Symptom Questionnaire 48; WHO-SSD = WHO-Screener for Somatoform Disorders; YSR = Youth Self-Report. Items in the symptom list who are not applicable for the old age are in italics. Lists derived from literature search after: October 2012, are in bold.

Quantification Inventory for Somatoform Syndromes (Wedekind et al., 2007), the SEPS (Tyrrer et al., 2012), the Somatic Symptoms Experiences Questionnaire (SSEQ; Herzog et al., 2014), and the Somatic Symptom Scale–8 (Gierk et al., 2014).

Less than half (20 out of 46, 43%) of these 46 scales have been administered in studies with exclusively or a substantial number of older adults. Of these studies, the BSI-18 SOM (Galdón et al., 2008), the BSI-53 SOM (Derogatis & Melisaratos, 1983), the Giessener Beschwerdebogen (Brähler et al., 2000), the Modified Somatic Perception Questionnaire (Main, 1983), and the PHQ-15-item version (PHQ-15; Kocalevent et al., 2013) have been applied in more than two studies (see Table 3). As pointed out below, these questionnaires, however, do not have the most optimal characteristics for an older population.

Of the 46 questionnaires, only the BSS has been validated for older adults (Ritchie et al., 2013). Nonetheless, the aim of the BSS was to evaluate overall symptom load in older adult populations in order to estimate illness burden and distress, so not necessarily somatization. Since especially symptoms related to common chronic conditions are included, it may more or less result in a symptom count relevant for the A-criterion of somatic symptom disorder in *DSM-5* (American Psychiatric Association, 2013). As 50% of the items overlap with common geriatric syndromes, the BSS is not a good (severity) indicator of possible MUS or somatization in later life.

For eight questionnaires (8 out of 46, 17%), normative data for older persons are available, that is, the BSI-53 SOM, BSS, Freiburg Complaint List (FBL; Fahrenberg, 1995), Giessener Beschwerdebogen, PHQ-15, SCL-90 SOM, Subjective Health Complaints Inventory (Ursin, Endresen, & Ursin, 1988), and the Somatic Symptom Scale–8. Therefore, future results based on any of these scales can be interpreted in the context of scores derived from another geriatric population.

With respect to overlap with common geriatric syndromes, we cannot give a cutoff value above which a specific list should be discouraged to use in geriatric population. The findings of Objective 1 suggests that even a low proportion of overlap may already be too much (i.e., 14%, 25%, and 33% for the BSI-53 SOM, SCL-90 SOM, and PHQ-15, respectively). Of the eight scales with normative data of a geriatric population, only one scale, the FBL has less than 14% of their items overlapping with symptoms of common geriatric syndromes. Since all questionnaires that use symptom counts to measure the level of somatization, we advise to only use these questionnaires in older population when adjustment for the common geriatric syndromes is possible.

Our review identified two scales without any overlap with common geriatric syndromes, that is, the SEPS and the SSEQ. Both scales focus on subjective experiences related to somatic symptoms instead of the symptoms themselves.

Unfortunately, none of these scales have normative data for older age groups.

Previously, the PHQ-15 and SCL-90 SOM have been identified as most suitable self-report somatic symptom questionnaires to be used in large-scale studies, because they have been extensively validated, are relatively short, easy to use, and of little burden to participants (Zijlema et al., 2013). Our findings, however, show that the PHQ-15 and the SCL-90 SOM considerably overlap with common geriatric symptoms and probably overestimate the level of somatization in older persons. Recently, somatic symptom count (based on the PHQ-15, added with 10 items on specific neurological symptoms and 5 on mental state) hardly differentiated between patients with ($n = 1,144$) and without ($n = 2,637$) MUS referred to an outpatient neurology clinic (Carson, Stone, Hansen, Duncan, & Cavanagh, 2015). Therefore, these questionnaires should only be used when adjustment for MES or common geriatric syndromes is possible.

Some methodological comments, however, need to be made. First of all, many recommendations have been given to increase validity and reliability when designing self-report questionnaires for older adults. Examples of these recommendations include the avoidance of reverse-scored items, a short questionnaire with preferably short, easy to understand items (to avoid fatigue of the participant), a dichotomized response scale (e.g., yes/no), and short reference period (time window) to avoid recall bias (e.g., Green, Goldstein, Sirockman, & Green, 1993; Johnco, Knight, Tadic, & Wuthrich, 2015; Pachana et al., 2007; Yesavage et al., 1983). To our knowledge, none of these recommendations have been empirically tested in order to show that adapting a questionnaire will indeed increase the validity of reliability when applied in older samples. Moreover, many studies, even among frail elderly, generally use several scales together without (reporting) any problems (e.g., Collard, Comijs, Naarding, & Oude Voshaar, 2014; Hanssen et al., 2016). Also, in clinical practice, older patients often have difficulty in making a dichotomous, often black or white choice. And finally, a short reference period seems less applicable for the often, chronic somatic symptoms related to somatization. Nonetheless, although the scientific merits of these criteria have to be established, they may be relevant when choosing a list for a specific study (e.g., as short as possible being a secondary outcome measure). Therefore, these characteristics have been summarized in Table 2.

Second, we focused on self-report questionnaires being most relevant to apply in research studies, acknowledging limited resources being at odds with sample size needs. Moreover, self-report questionnaires are also relevant in later life to get unbiased information about the complaints by minimizing the influence of formal and informal caregivers, often involved with older adults with physical complaints. On the other hand, older persons are more inclined than their younger counterparts to give socially desirable answers (e.g., Saeed et al., 2001).

Third, the validity of all questionnaires can be questioned as no gold standard exists for the measurement of explained and unexplained physical symptoms (especially not in later life). As the agreement between geriatricians whether a physical symptom is explained, partly explained or fully explained is quite high (Benraad et al., 2013), this could be considered as gold standard in future cross-validation studies in old-age samples.

To conclude, our review does not allow to give a simple advice which scale is most optimal to administer in studies in old age. This implies that the field is served by more studies cross-validating these questionnaires in an older population. Depending on the exact research question, however, specific choices can be made. In case overestimation of somatization should be excluded, the SEPS or SSEQ can be used (although both questionnaires are not validated in an older sample yet). These questionnaires, however, do not simply count symptoms, but rely on subjective experiences about physical symptoms. When symptom counts are needed as indication for possible somatization, the FBL emerges as the most optimal questionnaire when simply based on our criteria of the presence of normative data and a low percentage of overlap with common geriatric syndromes. Nonetheless, to our knowledge this questionnaire is only available in the German language and has as much as 78 items. For pragmatic reasons, therefore, one has to rely on less optimal alternatives, which are all provided by the present review. We present our recommendations for most optimal choices below, given the research objective.

Recommended for population-based cohort studies, especially when a broad age range is included (i.e., younger and older persons):

- Somatization subscale Symptom Checklist 90-item version^a
- Patient Health Questionnaire^a

Recommended for screening in primary care (based on lowest level of overlap with somatic diseases and availability for older patients):

1. Freiburger Beschwerden Liste/Freiburg Complaint List
2. Somatization subscale Brief Symptom Inventory 53-item version

Recommended for treatment monitoring (emphasis on subjective experiences):

1. Schedule for Evaluating Persistent Symptoms
 2. Somatic Symptoms Experiences Questionnaire
-

^aAdjustment for the common geriatric syndromes is necessary to make age-related comparisons.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

- Achenbach, T. M., & Rescorla, L. A. (2003). *Manual for the ASEBA adult forms & profiles*. Burlington: University of Vermont, Research Center for Children, Youth, & Families.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: Author.
- Andersen, K., Bogenschütz, M., Buhlinger, G., Behrendt, S., Bilberg, R., & Braun, B. (2015). Outpatient treatment of alcohol use disorders among subjects 60+: Design of a randomized clinical Trial conducted in three countries (Elderly studies). *BMC Psychiatry, 15*, 280.
- Asner-Self, K., Schreiber, J. B., & Marotta, S. A. (2006). A cross-cultural analysis of the Brief Symptom Inventory-18. *Cultural Diversity and Ethnic Minority Psychology, 12*, 367-375.
- Attanasio, V., Andrasik, F., Blanchard, E. B., & Arena, J. G. (1984). Psychometric properties of the SUNYA revision of the Psychosomatic Symptom Checklist. *Journal of Behavioral Medicine, 7*, 247-257.
- Benraad, C. E. M., Hilderink, P. H., van Driel, T. J. W., Disselhorst, L. G., Lubberink, B., van Wolferen, L., . . . Oude Voshaar, R. C. (2013). Physical functioning in older persons with somatoform disorders: A pilot study. *Journal of the American Medical Directors Association, 14*, 9-13.
- Bohman, H., Jonsson, U., Paaren, A., von Knorring, L., Olsson, G., & von Knorring, A. L. (2012). Prognostic significance of functional somatic symptoms in adolescence: A 15-year community-based follow-up study of adolescents with depression compare with healthy peers. *BMC Psychiatry, 12*, 90.
- Brähler, E., Schumacher, J., & Brähler, C. (2000). First all-Germany standardization of the brief form of the Giessen Complaints Questionnaire GBB-24. *Psychotherapie Psychosomatik Medizinische Psychologie, 50*, 14-21.
- Buttz-Lilly, A., Fink, P., Ørnboel, V. G., Christensen, K., & Rosendal, M. (2015). A new questionnaire to identify bodily distress in primary care: The "BDS checklist." *Journal of Psychosomatic Resource, 78*, 536-545.
- Callixte, K. T., Clet, T. B., Jacques, D., Faustin, Y., Francois, D. J., & Maturin, T. T. (2015). The pattern of neurological diseases in elderly people in outpatient consultations in Sub-Saharan Africa. *BMC Research, 8*, 159.
- Campo, R., Agarwal, N., LaStayo, P. C., O'Connor, K., & Pappas, L. (2014). Levels of fatigue and distress in senior prostate cancer survivors enrolled in a 12-week randomized controlled trial of Qigong. *Journal of Cancer Survival, 8*, 60-69.
- Carlier, I., Schulte-Van Maaren, Y., Wardenaar, K., Giltay, E., Van Noorden, M., Vergeer, P., & Zitman, F. (2012). Development and validation of the 48-item Symptom Questionnaire (SQ-48) in patients with depressive, anxiety and somatoform disorders. *Psychiatry Research, 200*, 904-910.
- Carson, A. J., Stone, J., Hansen, C. H., Duncan, R., & Cavanagh, J. (2015). Somatic symptom count scores do not identify patients with symptoms unexplained by disease: A prospective cohort study of neurology patients. *Journal of Neurology, Neurosurgery, and Psychiatry, 86*, 295-301.
- Charlson, M. E., Pompei, P., Ales, K. L., & MacKenzie, C. R. (1987). A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *Journal of Chronic Diseases, 40*, 373-383.

- Chester, G. A. (2001). Normative data for the brief symptom inventory for mature and independent living adults. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 62(4-B), 2108.
- Cohen, M. (2014). Depression, anxiety, and somatic symptoms in older cancer patients: A comparison across age groups. *Psychooncology*, 23, 151-157.
- Collard, R. M., Comijs, H. C., Naarding, P., & Oude Voshaar, R. C. (2014). Physical frailty: Vulnerability of patients suffering from late-life depression. *Aging & Mental Health*, 18, 570-578.
- Csoff, R. M., Macassa, G., & Lindert, J. (2010). Somatic complaints among elderly migrants in Germany. *Zeitschrift für Psychiatrie, Psychologie und Psychotherapie*, 58, 199-206.
- de Waal, M. W. M., Arnold, I. A., Spinhoven, P., Eekhof, J. A. H., & van Hemert, A. M. (2005). The reporting of specific physical symptoms for mental distress in general practice. *Journal of Psychosomatic Research*, 59, 89-95.
- Derogatis, L. R. (1975). *The Symptom Checklist-90-R*. Baltimore, MD: Clinical Psychometric Research.
- Derogatis, L. R., Lipman, R. S., Rickels, K., Uhlenhuth, E. H., & Covi, L. (1974). The Hopkins Symptom Checklist (HSCL): A self-report symptom inventory. *Behavioral Sciences*, 19, 1-15.
- Derogatis, L. R., & Melisaratos, N. (1983). The Brief Symptom Inventory: An introductory report. *Psychological Medicine*, 13, 595-605.
- Donaldson, M. B., Learman, K., Wright, A., Brown, C., Howes, C., & Cook, C. E. (2011). Factor structure and concurrent/convergent validity of the Modified Somatic Perception Questionnaire and Pain Beliefs Instrument. *Journal of Manipulative and Physiological Therapeutics*, 34, 30-36.
- Dijk, S. V., Voshaar, R. O., Lucassen, P., Comijs, H., & Hanssen, D. (2015). Big Five personality and medical unexplained symptoms in later life. *American Journal of Geriatric Psychiatry*, 23, S76-S78.
- Fabiao, C., Silva, M.C., Barbosa, A., Fleming, M., & Rief, W. (2010). Assessing medically unexplained symptoms: evaluation of a shortened version of the SOMS for use in primary care. *BMC Psychiatry*, 10, 34.
- Fahrenberg, J. (1995). Somatic complaints in the German population. *Journal of Psychosomatic Research*, 39, 809-817.
- Fortin, M., Stewart, M., Poitras, M. E., Almirall, J., & Maddocks, H. A. (2012). Systematic review of prevalence studies on multimorbidity: Toward a more uniform methodology. *Annals of Family Medicine*, 10, 142-151.
- Friborg, J. T., Yuan, J. M., Wang, R., Koh, W. P., Lee, H. P., & Yu, M. C. (2007). A prospective study of tobacco and alcohol use as risk factors for pharyngeal carcinomas in Singapore Chinese. *Cancer*, 109, 1183-1191.
- Galdón, M. J., Durá, E., Andreu, Y., Ferrando, M., Murgui, S., Pérez, S., & Ibanez, E. (2008). Psychometric properties of the Brief Symptom Inventory-18 in a Spanish breast cancer population. *Journal of Psychosomatic Research*, 65, 533-539.
- Garber, J., Walker, L. S., & Zeman, J. (1991). Somatization symptoms in a community sample of children and adolescents: Further validation of the Children's Somatization Inventory. *Journal of Consulting and Clinical Psychology*, 3, 588-595.
- Gierk, B., Kohlmann, S., Kroenke, K., Spangenburg, L., Zenger, M., Brähler, E., & Lowe, B. (2014). The Somatic Symptom Scale-8 (SSS-8): A brief measure of somatic symptom burden. *JAMA Internal Medicine*, 174, 399-407.
- Gledhill, J., & Garralda, M. E. (2006). Functional symptoms and somatoform disorders in children and adolescents: The role of standardised measures in assessment. *Child and Adolescent Mental Health*, 11, 208-214.
- Graham, C., Balard, C., & Pak, S. (1997). Carers' knowledge of dementia, their coping strategies and morbidity. *International Journal of Geriatric Psychiatry*, 12, 931-936.
- Grant, G. A., Nolan, M., & Ellis, N. A. (1990). A reappraisal of the Malaise Inventory. *Social Psychiatry and Psychiatric Epidemiology*, 25, 170-178.
- Green, J., Goldstein, F., Sirockman, B. E., & Green, R. C. (1993). Variable awareness of deficits in Alzheimer's disease. *Neuropsychiatry, Neuropsychology, and Behavioral Neurology*, 6, 159-165.
- Gunzelmann, T., Hinz, A., & Brähler, E. (2006). Subjective health in older people. *Psycho-Social-Medicine*, 27, 3.
- Gunzelmann, T., Schumacker, J., & Brähler, E. (1996). Physical complaints in old age: Standardization of the Giessen Complaint Questionnaire GBB-24 in over 60-year-old patients. *Zeitschrift für Gerontologie und Geriatrie*, 29, 110-118.
- Gunzelmann, T., Schumacker, J., & Brähler, E. (2002). Prevalenz von Schmerzen im alter. [Prevalence of pain in late life] *Ergebnisse Schmerz*, 16, 249-254.
- Hale, W. D., & Cochran, C. D. (1992). Age differences in self-reported symptoms of psychological distress. *Journal of Clinical Psychology*, 48, 633-637.
- Hale, W. D., Cochran, C. D., & Hedgepeth, B. E. (1984). Norms for the elderly on the Brief Symptom Inventory. *Journal of Consulting and Clinical Psychology*, 52, 321-322.
- Hanssen, D. J., Lucassen, P. L., Hilderink, P. H., Naarding, P., & Oude Voshaar, R. C. (2016). Health-related quality of life in older persons with medically unexplained symptoms. *American Journal of Geriatric Psychiatry*, 24, 1117-1127.
- Hassel, A. J., Rolko, C., Leisen, J., Schmitter, M., Rexroth, W., & Leckel, M. (2007). Oral health-related quality of life and somatization in the elderly. *Quality Life Research*, 16, 253-261.
- Havakeshian, S., & Mannion, A. F. (2013). Negative beliefs and psychological disturbance in spine surgery patients: A cause or consequence of a poor treatment outcome? *European Spine Journal*, 22, 2827-2835.
- Herman, S. L., & Lester, D. (1994). Physical symptoms of stress, depression, and suicidal ideation in high school students. *Adolescence*, 29, 639-641.
- Herzog, A., Voigt, K., Meyer, B., Wollburg, E., Weinmann, N., Langs, G., & Löwe, B. (2014). Psychological and interactional characteristics of patients with somatoform disorders: Validation of the Somatic Symptoms Experiences Questionnaire (SSEQ) in a clinical psychosomatic population. *Journal of Psychosomatic Research*, 78, 553-562.
- Hilderink, P. H., Benraad, C. E. M., van Driel, D., Buitelaar, J. K., Speckens, A. E. M., Olde Rikkert, M. G. M., & Oude Voshaar, R. C. (2009). Medically unexplained physical symptoms in elderly people: A pilot study of psychiatric geriatric characteristics. *American Journal of Geriatric Psychiatry*, 17, 1085-1088.

- Hilderink, P. H., Collard, R., Rosmalen, J. G. M., & Oude Voshaar, R. C. (2013). Prevalence of somatoform disorders and medically unexplained symptoms in old age populations in comparison with younger age groups: A systematic review. *Aging Research Reviews, 12*, 151-156.
- Hinton, D. E., Kredlow, M. A., Bui, E., Pollack, M. H., & Hofmann, S. G. (2012). Treatment change of somatic symptoms and cultural syndromes among Cambodian refugees with PTSD. *Depression and Anxiety, 29*, 147-154.
- Jellesma, F. C., Rieffe, C., & Terwogt, M. M. (2007). The somatic complaint list: Validation of a self-report questionnaire assessing somatic complaints in children. *Journal of Psychosomatic Research, 63*, 399-401.
- Jeong, H., Han, C., Park, M. H., Ryu, S. H., Pae, C. U., Lee, J. Y., . . . Steffens, D. C. (2014). Influence of the number and severity of somatic symptoms on the severity of depression and suicidality in community-dwelling elders. *Asia-Pacific Psychiatry, 3*, 274-283.
- Johnco, C., Knight, A., Tadic, D., & Wuthrich, V. M. (2015). Psychometric properties of the Geriatric Anxiety Inventory (GAI) and its short-form (GAI-SF) in a clinical and non-clinical sample of older adults. *International Psychogeriatrics, 27*, 1089-1097.
- Kahn, A. M., & Taj, R. (2011). To assess the extent of psychological impact on IDP's residing in the camps of Shahmansoor, Swabi, Khyber Pukhtun Kha, Pakistan. *European Psychiatry, 26*, 1.
- Katz, J. N., Chang, L. C., Sngha, O., Fossel, A. H., & Bates, D. W. (1996). Can comorbidity be measured by questionnaire rather than medical record review? *Medical Care, 34*, 73-84.
- Kellner, R. (1987). A symptom questionnaire. *Journal of Clinical Psychiatry, 48*, 268-274.
- Khoo, E. M., Mathers, N. J., McCarthy, S. A., & Low, W. Y. (2012). Somatisation disorder and its associated factors in multi-ethnic primary care clinic attenders. *International Journal of Behavioral Medicine, 19*, 165-173.
- Klein, M. B., Lezotte, D. C., Heltshe, S., Fauerbach, J., & Holavanahalli, R. K. (2011). Functional and psychosocial outcomes of older adults after burn injury: Results from a multicenter database of severe burn injury. *Journal of Burn Care & Research, 32*, 66-78.
- Kliem, S., Beller, J., Kroger, C., Birowicz, T., Zenger, M., & Brähler, E. (2014). Dimensional latent structure of somatic symptom reporting in two representative population studies: Results from taxometric analyses. *Psychological Assessment, 26*, 484-492.
- Kocalevent, R.-D., Hinz, A., & Brähler, E. (2013). Standardization of a screening instrument (PHQ-15) for somatization syndromes in the general population. *BioMed Central Psychiatry, 13*, 91.
- Koorevaar, R. C., Terluin, B., van 't Riet, E., Madden, K., & Bulstra, S. K. (2016). Validation of the four-dimensional symptom questionnaire (4DSQ) and prevalence of psychological symptoms in orthopedic shoulder patients. *Journal of Orthopedic Resource, 34*, 683-691.
- Kroenke, K., Spitzer, R. L., deGruy, F. V., & Swindle, R. (1998). A symptom checklist to screen for somatoform disorders in primary care. *Psychosomatics, 39*, 263-272.
- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2002). The PHQ-15: Validity of a new measure for evaluating the severity of somatic symptoms. *Psychosomatic Medicine, 64*, 258-266.
- Ladwig, K. H., Marten-Mittag, B., Lacruz, M. E., Henningsen, P., & Creed, F. (2010). Screening for multiple somatic complaints in a population-based survey: Does excessive symptom reporting capture the concept of somatic symptom disorders? Findings from the MONICA-KORA Cohort Study. *Journal of Psychosomatic Research, 68*, 427-437.
- Lecrubier, Y., Sheehan, D. V., Weiller, E., Amorim, P., Bonora, I., Harnett Sheehan, K., . . . Dunbar, G. C. (1997). The Mini International Neuropsychiatric Interview (MINI). A short diagnostic structured interview: Reliability and validity according to the CIDI. *European Psychiatry, 12*, 224-231.
- Lhiebaek, C., Eriksen, H. R., & Ursin, H. (2002). Prevalence of subjective health complaints (SHC) in Norway. *Scandinavian Journal of Public Health, 30*, 20-29.
- Lipowski, Z. J. (1988). Somatization: The concept and its clinical application. *American Journal of Psychiatry, 145*, 1358-1368.
- MacLean, E.W., Palsson, O.S., Turner, M.J., & Whitehead, W.E. (2012). Development and validation of new disease-specific measures of somatization and comorbidity in IBS. *Journal of Psychosomatic Research, 73*, 351-355.
- Miller, M., Paradis, C. F., Houck, P. R., Mazumdar, S., Stack, J. A., Rifai, A. H., . . . Reynolds, C. F., III. (1992). Rating chronic medical illness burden in geropsychiatric practice and research: Application of the Cumulative Illness Rating Scale. *Psychiatric Research, 41*, 237-248.
- Main, C. J. (1983). The Modified Somatic Perception Questionnaire (MSPQ). *Journal of Psychosomatic Research, 27*, 503-514.
- Manu, P., Lane, T. J., Matthews, D. A., & Escobar, J. I. (1989). Screening for somatization disorder in patients with chronic fatigue. *General Hospital Psychiatry, 11*, 294-297.
- Montalban, R., Comas, V. A., & Garcia-Garcia, M. (2010). Validation of the Spanish version of the PHQ-15 questionnaire for the evaluation of physical symptoms in patients with depression and/or anxiety disorders: DEPRESOMA study. *Actas Espanolas de Psiquiatria, 38*, 345-357.
- Mumford, D. B. (1989). Somatic sensations and psychological distress among students in Britain and Pakistan. *Social Psychiatry and Psychiatric Epidemiology, 24*, 321-326.
- Mumford, D. B., Bavington, J. T., Bhatnagar, K. S., Hussain, Y., Mirza, S., & Naraghi, M. M. (1991). The Bradford Somatic Inventory: A multiethnic inventory of somatic symptoms reported by anxious and depressed patients in Britain and the Indo-Pakistan subcontinent. *British Journal of Psychiatry, 158*, 379-386.
- Nettleton, S. (2006). "I just want permission to be ill": Towards a sociology of medically unexplained symptoms. *Social Science & Medicine, 62*, 1167-1178.
- Nordin, S., Palmquist, E., & Nordin, M. (2013). Psychometric evaluation and normative data for a Swedish version of the Patient Health Questionnaire 15-Item Somatic Symptom Severity Scale. *Scandinavian Journal of Psychology, 54*, 112-117.
- Othmer, E., & DeSouza, C. (1985). A screening test for somatization disorder (hysteria). *American Journal of Psychiatry, 142*, 1146-1149.

- Pachana, N. A., Byrne, G. J., Siddle, H., Koloski, N., Harley, E., & Arnold, E. (2007). Development and validation of the Geriatric Anxiety Inventory. *International Psychogeriatrics, 19*, 103-114.
- Phan, T. (2004). An ethnographically derived measure of anxiety, depression and somatization: The Phan Vietnamese Psychiatric Scale. *Transcultural Psychiatry, 41*, 200-232.
- Pennebaker, J. W. (1982). *The psychology of physical symptoms*. New York, NY: Springer-Verlag.
- Pereira, M., Martins, A., Alves, S., & Canavaro, C. (2014). Assessing quality of life in middle-aged and older adults with HIV: Psychometric testing of the WHOQOL-HIV-Bref. *Quality of Life Research, 23*, 2473-2479.
- Petkus, A. J., Gum, A. M., King-Kallimanis, B., & Wetherell, J. L. (2009). Trauma history is associated with psychological distress and somatic symptoms in home bound older adults. *American Journal of Geriatric Psychiatry, 17*, 810-818.
- Phillips, K. A., Fallon, B. A., & King, S. (2008). Somatoform and factitious disorders and malingering measures. In A. J. Rush, M. B. First & D. Blacker (Eds.), *Handbook of psychiatric measures* (2nd ed., pp. 559-585). Arlington, VA: American Psychiatric Publishing.
- Pietrzak, R. H., & Petry, N. M. (2006). Severity of gambling problems and psychosocial functioning in older adults. *Journal of Geriatric Psychiatry and Neurology, 19*, 106-113.
- Pietrzak, R. H., Robert, H., Molina, C., Ladd, G. T., & Kerins, G. J. (2005). Health and psychosocial correlates of disordered gambling in older adults. *American Journal of Geriatric Psychiatry, 13*, 510-519.
- Pilowsky, I. (1967). Dimensions of hypochondriasis. *British Journal of Psychiatry, 113*, 89-93.
- Russell, L., Gough, K., Drosowsky, A., Schofield, P., Aranda, S., Butow, P. N., . . . Jefford, M. (2015). Psychological distress, quality of life, symptoms and unmet needs of colorectal cancer survivors near the end of treatment. *Journal of Cancer Survival, 9*, 462-470.
- Qian, J., Rem, Z., Yu, D., He, X., & Li, C. (2014). The value of the Patient Health Questionnaire-15 (PHQ-15) for screening somatic symptoms in general hospital. *Chinese Mental Health Journal, 28*, 173-178.
- Quine, L., & Charnley, H. (1987). The malaise inventory as a measure of stress in carers. In J. Twigg (Ed.), *Evaluating support to informal carers of mentally handicapped children and elderly people*. Research report from the University of Kent, Personal Social Services Research Unit. Presented at a conference held in York.
- Rabung, S., Harfst, T., Kowski, S., Koch, U., Wittchen, H. U., & Schultz, H. (2009). Psychometric analysis of a short form of the "Hamburg Modules for the Assessment of Psychosocial Health" (HEALTH-49). *Zeitschrift für Psychosomatische Medizin und Psychotherapie, 55*, 162-179.
- Ricceri, F., Del Basso, D., Tomba, E., Offidani, E., & Prignano, F. (2014). Clinimetric assessment in psoriasis patients with different illness severity. *Journal of the American Academy of Dermatology, 70*(Suppl. 1), AB166.
- Rief, W., & Hiller, W. (2003). A new approach to the assessment of the treatment effects of somatoform disorders. *Psychosomatics, 44*, 492-498.
- Rief, W., & Hiller, W. (2008). *Das Screening für somatoforme Störungen SOMS. Manual zum Fragebogen* [The screening for somatoform symptoms]. Bern, Switzerland: Huber-Verlag.
- Ritchie, C. S., Hearld, K. R., Gross, A., Allman, R., Sawyer, P., Sheppard, K., . . . Roth, D. L. (2013). Measuring symptoms in community-dwelling older adults: The psychometric properties of a brief symptom screen. *Medical Care, 51*, 949-955.
- Ritsner, M., Ponizovsky, A., Kurs, R., & Modai, I. (2000). Somatization in an immigrant population in Israel: A community survey of prevalence, risk factors, and help-seeking behavior. *American Journal of Psychiatry, 157*, 385-392.
- Rodgers, B., Pickles, A., Power, C., Collishaw, S., & Maughan, B. (1999). Validity of the Malaise Inventory in general population samples. *Social Psychiatry and Psychiatric Epidemiology, 34*, 333-341.
- Roh, J., Kim, B., Jang, J., Seo, W., Lee, S., Kim, J., . . . Koh, S. (2008). The relationship of pain and health-related quality of life in Korean patients with Parkinson's disease. *Acta Neurologica Scandinavica, 119*, 793-403.
- Rudwan, S. J. (2000). A checklist of the Syrian psychological psychosomatic symptoms. *Journal of Social Science, 28*, 113-138.
- Sack, M., Boroske-Leiner, K., & Lahmann, C. (2010). Association of nonsexual and sexual traumatizations with body image and psychosomatic symptoms in psychosomatic outpatients. *General Hospital Psychiatry, 32*, 315-320.
- Saeed, K., Mubbashar, S., Dogar, I., Mumford, D., & Mubbashar, M. (2001). Comparison of self-reporting Questionnaire and Bradford Somatic Inventory as screening instruments for psychiatric morbidity in community settings in Pakistan. *Medical Forum Monthly, 12*, 14-16.
- Schafer, I., Hansen, H., Schon, G., Hofels, S., & Altiner, A. (2012). The influence of age, gender and socio-economic status on multimorbidity patterns in primary care: First results from the multicare cohort study. *BMC Health Service resource, 12*, 89.
- Schat, A. C., Kelloway, E. K., & Desmarais, S. (2005). The Physical Health Questionnaire (PHQ): Construct validation of a self-report scale of somatic symptoms. *Journal of Occupational Health Psychology, 10*, 363-381.
- Senn, N., & Monod, S. (2015). Development of a comprehensive approach for the early diagnosis of geriatric syndromes in general practice. *Frontiers in Medicine, 2*, 78.
- Sha, M., Callahan, C. M., Counsell, S. R., Westmoreland, G. R., Stump, T. E., & Kroenke, K. (2005). Physical symptoms as a predictor of health care use and mortality among older adults. *American Journal of Medicine, 118*, 301-306.
- Spence, J. T., Helmreich, R. L., & Pred, R. S. (1987). Impatience versus achievement strivings in the type A pattern: Differential effects on students' health and academic achievement. *Journal of Applied Psychology, 72*, 522-528.
- Srinivasan, T. N., & Suresh, T. R. (1991). The nonspecific symptom screening method: Detection of nonpsychotic morbidity based on nonspecific symptoms. *General Hospital Psychiatry, 13*, 106-114.
- Sloane, P. D., Hartman, M., & Mitchell, C. M. (1994). Psychological factors associated with chronic dizziness in patients aged 60 and older. *Journal of the American Geriatric Society, 42*, 847-852.

- Stankunas, M., Soares, J. F. J., Stankuniene, M. G. M., Torres, G. F., Loannidi-Kapolou, E., Barros, H., & Lindert, J. (2013). Differences in reporting somatic complaints in elderly by education level. *Central European Journal of Medicine*, *8*, 125-131.
- Stankuniene, A., Stankunas, M., Soares, J. J., Avery, M., Melchiorre, G., Torres-Gonzales, F., . . . Lindert, J. (2012). Somatic complaints and refrain from buying prescribed medications: Results from a cross-sectional study on people 60 years and older living in Kaunas (Lithuania). *Daru*, *20*, 78.
- Staerkle, R., Mannion, A., Elfering, A., Junge, A., Semmer, N. K., Jacobshagen, N., . . . Boos, D. (2004). Longitudinal validation of the Fear-Avoidance Beliefs Questionnaire (FABQ) in a Swiss-German sample of low back pain patients. *European Spine Journal*, *13*, 332-340.
- Swartz, M., Hughes, D., George, L., Blazer, D., Landerman, R., & Buchholz, K. (1986). Developing a screening index for community studies of somatization disorder. *Journal of Psychiatric Research*, *20*, 335-343.
- Tanji, H., Anderson, K. E., Gruber-Baldini, A. L., Fishman, P. S., Reich, S. G., Weiner, W. J., & Shulman, L. M. (2008). Mutuality of the marital relationship in Parkinson's disease. *Movement Disorder*, *23*, 1843-1849.
- Terluin, B., van Marwijk, H. W., Ader, H. J., de Vet, H. C., Penninx, B. W., & Hermens, M. L. (2006). The Four-Dimensional Symptom Questionnaire (4DSQ): A validation study of a multidimensional self-report questionnaire to assess distress, depression, anxiety and somatization. *BMC Psychiatry*, *6*, 34.
- Thygesen, E., Lindstrom, T. C., Saevaroid, H. I., & Engedal, K. (2009). The Subjective Health Complaints Inventory: A useful instrument to identify various aspects of health and ability to cope in older people? *Scandinavian Journal of Public Health*, *37*, 690-696.
- Tyrer, H., Ali, L., Cooper, F., Selvwright, P., Bassett, P., & Tyrer, P. (2012). The Schedule for Evaluating Persistent Symptoms (SEPS): A new method of recording medically unexplained symptoms. *International Journal of Social Psychiatry*, *59*, 281-287.
- Ursin, H., Endresen, I. M., & Ursin, G. (1988). Psychological factors and self-reports of muscle pain. *European Journal of Applied Physiology and Occupational Physiology*, *57*, 282-290.
- Valdearenas, L., Torres-Gonzalez, F., de Dios Luna, J., & Cervilla, J. (2012). Prevalence and correlates of psychotic symptoms in the Spanish elderly community. *European Psychiatry*, *27*(Suppl. 1), 1.
- van Noorden, M., van Fenema, E. M., van der Wee, N. J. A., van Rood, Y. R., & Carlier, I. V. E. (2012). Predicting outcomes of mood, anxiety and somatoform disorders: The Leiden routine outcome monitoring study. *Journal of Affective Disorder*, *142*, 122-131.
- Videler, A. C., Rossi, G., Schoevaars, M., van der Feltz-Cornelis, C. M., & van Alphen, S. P. (2014). Effects of schema group therapy in older outpatients: A proof of concept study. *International Psychogeriatrics*, *26*, 1709-1717.
- Vulić-Prtorić, A. (2005). Upitnik Psihosomatskih Simptoma za Djecu i Adolescente: Procjena Valjanosti [Questionnaire for psychosomatic symptoms in children and adolescents: Evaluation of validity]. *Suvremena Psihologija*, *8*, 211-227.
- Walker, L. S., Garber, J., & Greene, J. W. (1991). Somatization symptoms in pediatric abdominal pain patients: Relation to chronicity of abdominal pain and parent somatization. *Journal of Abnormal Child Psychology*, *19*, 379-394.
- Wedekind, D., Bandelow, B., Fentzahn, E., Trümper, P., & Rütger, E. (2007). The quantification inventory for somatoform syndromes (QUISS): A novel instrument for the assessment of severity. *European Archives of Psychiatry and Clinical Neuroscience*, *257*, 153-163.
- Wetherell, J. L., Ayers, C. R., Nuevo, R., Stein, M. B., Ramsdell, J., & Patterson, T. L. (2010). Medical conditions and depressive, anxiety, and somatic symptoms in older adults with and without generalized anxiety disorder. *Aging & Mental Health*, *14*, 764-768.
- Weiss, F., Kleinstuber, M., & Rief, W. (2016). Health care utilization in outpatients with somatoform disorders: Descriptives, interdiagnostic differences, and potential mediating factors. *Journal of General Hospital Psychology*, *44*, 22-29.
- Wisniewski, J. J., Naglieri, J. A., & Mulick, J. A. (1988). Psychometric properties of a Children's Psychosomatic Symptom Checklist. *Journal of Behavioral Medicine*, *11*, 497-507.
- World Health Organization. (1993). *The ICD-10 classification of mental and behavioural disorders. Diagnostic criteria for research*. Geneva, Switzerland: Author.
- Yesavage, J. A., Brink, T. L., Rose, T. L., Lum, O., Huang, V., Adey, M., & Leirer, V. O. (1983). Development and validation of a Geriatric Depression Screening Scale: A preliminary report. *Journal of Psychiatric Research*, *17*, 37-39.
- Zijlema, W., Stolk, R., Lowe, B., Rief, W., BioSHaRE, White, P. D., & Rosmalen, J. G. M. (2013). How to assess common somatic symptoms in large-scale studies: A systematic review of questionnaires. *Journal of Psychosomatic Research*, *4*, 459-468.
- Zweig, R. A., & Türkel, E. (2007). The social adjustment scale-self-report: Psychometric properties for older adults. *Psychological Reports*, *101*, 920-926.