

and Diabetes Laboratory, University of British Columbia, Vancouver, British Columbia, Canada, 6. University of British Columbia, Vancouver, British Columbia, Canada, 7. Center for Hip Health and Mobility, Vancouver Coastal Health Research Institute, Vancouver, Canada

We assessed the efficacy of the home-based Otago Exercise Program (OEP) as a secondary falls prevention strategy in seniors referred to a falls prevention clinic after an index fall. We conducted a 12-month randomized controlled trial of 344 adults, aged 70 years and older, with = or > 1 fall resulting in medical attention in the prior 12 months. Participants were randomized to OEP or standard of care (CON). The OEP is a home-based strength and balance training program delivered by a physical therapist. All participants received AGS Guideline Care for falls prevention from a geriatrician. Differences in falls rate was tested with a negative binomial regression model. The rate of falls was lower in the OEP group vs the CON group (incident rate ratio [IRR] = 0.64, 95% CI 0.46 to 0.90). The estimated incidence rate of falls per person-year was 1.4 (95% CI 0.1 to 2.0) in the OEP group and 2.1 (95% CI 0.1 to 3.2) in the CON group, with an absolute incidence rate difference of 0.74 (95% CI 0.04 to 1.78) falls per person-year. DSST performance also increased in the OEP group by a mean change of 1.1 points (95% CI 0.02 to 2.1) vs the CON group. Improved DSST was associated with fewer falls (IRR = 0.80, 95% CI 0.68 to 0.95). These findings support the use of the OEP for secondary falls prevention.

MULTIPLE MEDICATION USE AND RISK OF TREATED FALL INJURY: THE HEALTH ABC STUDY

Lingshu Xue¹ Robert M. Boudreau,² Julie M. Donohue² Janice C. Zgibor³ Zachary A. Marcum⁴ Tina Costacou,² Anne B. Newman,² and Elsa Strotmeyer², 1. *University of Pittsburgh, Pittsburgh, United States*, 2. *University of Pittsburgh, Pittsburgh, Pennsylvania, United States*, 3. *University of South Florida, Tampa, Florida, United States*, 4. *University of Washington, Seattle, Washington, United States*

Multiple medication use within one year is associated with increased fall injury risk in older adults. However, chronically using multiple medications and treated fall injury have rarely been explored, particularly in cohort studies linked with claims data. We examined using >5 medications in 2 or more consecutive years (chronic medication use) as a risk factor for treated fall injury in 1,898 community-dwelling adults (age 73.6±2.9 years; 53% women; 37% black) with linked Medicare Fee-For-Service (FFS) claims from the Health, Aging and Body Composition Study since 1997/98 clinic visit. Incident fall injury (N=546) was the first claim from 1998/99 clinic visit to 12/31/08 with an ICD-9 fall code and non-fracture injury code, or fracture code with/without a fall code. Stepwise Cox models with a time-varying predictor of chronic medication use before fall injury or censoring (N=414) vs. not using >5 medications at the same time (N=1008) were adjusted for baseline demographics, lifestyle factors, fall history, quadriceps strength, cardiovascular disease (CVD), diabetes, sensory nerve impairment, and kidney function. Fall injury risk increased for chronic medication users (37%) vs. non-users (29%) (HR=1.25[1.00-1.57]), though was attenuated after adjustment for CVD and diabetes (HR=1.18[0.93-1.51]). Sensitivity analyses

excluding fall-risk-increasing drugs (FRIDs) from medication counts (HR=1.32[0.54-3.20]), or including those using >5 medications non-chronically (N=365) in referent groups (HR=1.22[0.96-1.55]) had consistent findings. Unmeasured comorbidity differences may confound associations of chronic medication use and treated fall injury risk in older adults with Medicare FFS. Considering both chronic diseases and medication use in fall risk assessments is needed.

PROGRAM CHARACTERISTICS ASSOCIATED WITH REDUCED FEAR OF FALLING: A SYSTEMATIC REVIEW AND META-ANALYSIS OF RCTS

Marlot Kruisbrink¹ Kim Delbaere² Gertrudis Kempen,¹ Rik Crutzen,¹ and G.A. Rixt Zijlstra¹, 1. *Maastricht University, Care and Public Health Research Institute (CAPHRI), Department of Health Services Research, Maastricht, Netherlands*, 2. *Neuroscience Research Australia, School of Public Health and Community Medicine, UNSW, Randwick, New South Wales, Australia*

Fear of falling (FOF) is common among older people and can result in activity avoidance and decreased physical functioning. Different types of interventions have demonstrated significant small reductions in FOF. To optimize effect sizes, we sought to identify characteristics of interventions that were associated with a change in FOF. Five scientific databases were searched for articles using randomized controlled trial designs in community-dwelling older people without medical conditions. Data extraction included intervention type, setting, group format, type of supervision, provider, delivery format, duration, number of sessions, contact time, and risk of bias (assessed with the Cochrane Collaboration's Risk of Bias Tool). After screening of titles, abstracts, and full texts, 55 unique studies – reporting on 68 interventions – were systematically reviewed. The majority of interventions focused on exercise (n=50). Interventions were performed at home (n=21) or in a community setting (n=23), were delivered in a group (n=26) or individual (n=30) format, and were often supervised (n=60) and delivered face-to-face (n=56). Duration ranged from 1 to 52 weeks and total contact time with the provider from 2 to 56 hours. Results of 42 interventions were suitable for meta-analysis. Univariate meta-regressions to evaluate associations between intervention characteristics and intervention effects directly after the intervention yielded no significant results. Due to self-reported outcomes and difficulties with blinding, risk of bias was high in all studies. To conclude, intervention characteristics were not associated with changes in FOF in this study. Possible reasons for an absence of associations and future research directions will be discussed.

SESSION 570 (SYMPOSIUM)

SEDENTARY BEHAVIOR AND PHYSICAL ACTIVITY IN THE ADULT CHANGES IN THOUGHT (ACT) STUDY

Chair: Dori E. Rosenberg, *Kaiser Permanente Washington Health Research Institute, Seattle, Washington, United States*

Co-Chair: Andrea Z. LaCroix, *University of California San Diego, La Jolla, California, United States*

Discussant: Jack Guralnik, *University of Maryland School of Medicine, Baltimore, Maryland, United States*