

Mucosal Invasion by Fusobacteria is a Common Feature of Acute Appendicitis in Germany, Russia, and China

Alexander Swidsinski, Yvonne Dörffel¹, Vera Loening-Baucke, Alexander Tertychnyy²,
Salbakay Biche-ool^{2,3}, Sergei Stonogin³, Yi Guo⁴, Ning-Dong Sun⁵

Departments of Gastroenterology, Laboratory for Molecular Genetics, Polymicrobial Infections and Bacterial Biofilms, Charité Universitätsmedizin Berlin, CCM, ¹Internal Medicine, Outpatient Clinic, Luisenstrasse 11-13, Charité Universitätsmedizin Berlin, 10117 Berlin, Germany, ²Russian State Medical University, Department of Pathology, Moscow Faculty, Ostrovityanova Steet. 1, 117997 Moscow, ³Department of Pediatrics, Tushinskaya Children's Hospital, Geroev Panfilovtsev Steet. 28, 125480 Moscow, Russian Federation, ⁴The State Key Laboratory of Virology 2004DA105204 and Department of Epidemiology, Wuhan University School of Public Health, Wuhan 430071, P.R. China, ⁵Department of Surgery, No. 457 Hospital of People's Liberation Army, Wuhan, Hubei Province, China

Address for correspondence:

Dr. Alexander Swidsinski, Laboratory for Molecular Genetics, Polymicrobial Infections and Bacterial Biofilms, Charité Universitätsmedizin Berlin, Campus Mitte, 10098 Berlin, Germany. E-mail: alexander.swidsinski@charite.de

ABSTRACT

Background/Aim: To investigate the geographic occurrence of mucosa-invading Fusobacteria in acute appendicitis. **Patients and Methods:** Carnoy- and formalin-fixed appendices from Germany, Russia, and China were comparatively investigated. Bacteria were detected using fluorescent *in situ* hybridization. Cecal biopsies from patients with inflammatory bowel disease and other conditions were used as disease controls. **Results:** Fusobacteria represented mainly by *Fusobacterium nucleatum* were the major invasive component in bacterial infiltrates in acute appendicitis but were completely absent in controls. The occurrence of invasive Fusobacteria in Germany, Russia, and China was the same. The detection rate in Carnoy-fixed material was 70–71% and in formalin-fixed material was 30–36%. **Conclusions:** Acute appendicitis is a polymicrobial infectious disease in which *F. nucleatum* and other Fusobacteria play a key role.

Key Words: Acute appendicitis, fluorescence *in situ* hybridization, Fusobacteria, *Fusobacterium necrophorum*, *Fusobacterium nucleatum*, inflammatory bowel disease, infection, intestinal microbiota, invasive bacteria, mucosal flora, polymicrobial infection

Received: 10.07.2011, Accepted: 31.10.2011

How to cite this article: Swidsinski A, Dörffel Y, Loening-Baucke V, Tertychnyy A, Biche-ool S, Stonogin S, *et al.* Mucosal invasion by fusobacteria is a common feature of acute appendicitis in Germany, Russia, and China. Saudi J Gastroenterol 2012;18:55-8.

Acute appendicitis is a common potentially life-threatening purulent disease in which multiple bacterial species are involved; the etiology, however, is unknown.

We recently published a study on mucosa-invading bacteria in acute appendicitis.^[1] The tissue samples from patients obtained during surgery for acute appendicitis were compared with biopsies from inflamed cecum of patients with inflammatory bowel disease (IBD), self-limiting colitis, diverticulosis, and biopsies from healthy controls. Bacteria were assessed using fluorescence *in situ* hybridization (FISH). Fusobacteria were found to be invasive in 62% of patients with proven appendicitis but in none of the controls.

Studying the literature for data indicating the infectious

Access this article online	
	Quick Response Code:
	Website: www.saudijgastro.com
	DOI: 10.4103/1319-3767.91734

origin of the disease, we found an article by Guo *et al.* published in 2004 in *The American Journal of Surgery* "Cluster of acute hemorrhagic appendicitis among high school students in Wuhan, China."^[2] Those authors concluded that the observed "cluster of acute appendicitis represent a special kind of appendicitis, with features of an infectious disease in epidemiology."

To compare our findings of fusobacterial invasion in acute appendicitis, we asked the authors to send us sections of the resected appendices. In addition, we asked a pathologist from Moskau, Russia, to provide us the samples from patients with acute appendicitis fixated both in Carnoy solution and formalin.

PATIENTS AND METHODS

Appendices

Materials from appendices removed during laparoscopic emergent appendectomy were investigated. None of the patients had received preoperative antibiotic treatment. All patients were operated within 24 h after onset of symptoms for suspected acute appendicitis. They had no history of previous episodes, ruling out chronic appendicitis.

The surgeries in China and Russia as our previous German study were performed without preoperative antibiotics.

Germany

The material of appendices and biopsies obtained in Germany were described in a previous publication.^[1] Altogether 52 appendices with proven acute appendicitis were investigated.

Cecal biopsies from 100 patients with ulcerative colitis, 100 patients with Crohn's disease, 50 patients with self-limiting colitis, 50 patients with diverticulosis, 50 patients with irritable bowel syndrome, and 50 healthy controls were investigated.^[1]

Russia

Twenty-three appendices with intraoperatively confirmed acute appendicitis were removed during laparoscopic surgery in Russia Tushinskaya Children's Hospital, Moscow. Patients were 6–14 years old (mean age, 11 years). The study was prospective and included a general surgeon and a pathologist. The resected appendices were divided into 2 parts. One part was fixed in Carnoy solution (6/6/1 vol. ethanol/glacial acetic acid/chloroform) and the other in formalin, and then both were embedded in paraffin using standard techniques. All patients were operated within 24 h after onset of symptoms for suspected acute appendicitis. Forty cecal biopsies were taken from children, 12–18 years of age (mean age, 14 years), with confirmed active IBD from macroscopically inflamed

regions. All biopsies were fixated in Carnoy and embedded in paraffin.

China

Sections of 11 appendices were received from the Department of Surgery, No. 457, Hospital of People's Liberation Army, where the patients were operated during an outbreak of acute appendicitis among high school students in China.^[2] The resected appendices were previously fixated in formalin and then embedded in paraffin using standard techniques.

The initial paraffin blocks from the resections were not available, since they were used up for the extended histopathologic evaluation of an appendicitis outbreak published previously.^[2] Of the initial sections of material 1–3 sections of each patient had remained intact and were used for this study.

Fluorescent *in situ* hybridization

Sections of 4 µm thickness were placed on SuperFrost plus slides (R. Langenbrinck, Emmendingen, Germany). Multicolor FISH was performed according to previously described protocols for evaluation of tissue specimens and identification of bacteria^[3,4] with a Nikon e600 fluorescence microscope (Nikon, Tokyo, Japan) and Nikon DXM1200 camera. Altogether 7 FISH probes were applied.

The Cy3-labeled probe (orange fluorescence) that characterized a bacterial group of interest: *F. nucleatum* (Fnuc),^[5] *F. necrophorum* (Fnec),^[5] or Fusobacteria (Fuso)^[5] was simultaneously hybridized with the universal for virtually all bacteria *Eubacteria*-specific (Eub388)^[6] Alexa488-labeled probe (green fluorescence) and a mix of Cy5-labeled probes (red fluorescence) that included the main faecal bacterial groups such as *Faecalibacterium prausnitzii* (Fprau),^[7] *Bacteroides* (Bac303),^[8] *Clostridium* group and XIVa/Roseburia (Erec482).^[9] The fluorescence signals of specific probes, universal probe, and probes for Fprau+Bac303+Erec482 groups were overlaid within the same microscopic field.

RESULTS

Histopathology

Germany

Twenty-seven of 52 patients with the clinical and intraoperative diagnosis of acute appendicitis had suppurative appendicitis and 25 had catarrhal appendicitis by histology.

Russia

Eleven of 23 patients with clinical and intraoperative diagnosis of acute appendicitis had suppurative appendicitis and 12 had catarrhal appendicitis by histology.

China

The initial paraffin blocks from the resections were not available, since they were used up for extended histopathologic examination of an appendicitis outbreak published previously. One to three sections of the initial resected appendices from each patient remained intact and were used for this study. All 11 patients had a suppurative appendicitis.

Infiltration of the mucosa by Fusobacteria in acute appendicitis

The occurrence of invasive Fusobacteria in Carnoy-fixed material from appendices in Germany and Russia were nearly the same [Table 1]. In the German material, the invasive Fusobacteria were mainly represented by *F. nucleatum* (79%), *F. necrophorum* (12%), and Fusobacteria (9%) that could not be determined at species level as they were positive with Fuso but negative with the Fnuc and Fnec FISH probes. *F. nucleatum* composed 85% and *F. necrophorum* 15% of the invasive Fusobacteria in the appendical material from Russia.

The invasion of Fusobacteria had an appearance of a needle-like infiltration of single epithelial cells within an intact appearing epithelial layer in patients with catarrhal appendicitis. We called them “pinned cells” because of the marked appearance of these infiltrates [Figure 1]. A deep infiltration of the submucosa with Fusobacteria was seen in suppurative appendicitis [Figure 2].

The detection of bacteria in formalin-fixed appendices was difficult. The hybridization signals even with the universal bacterial probe were often very weak. However, using the *F. nucleatum*-specific probe, we could definitively detect invasive Fusobacteria in 30% of appendices from Russia and in 36% appendices from China [Table 1 and Figure 3].

Invasive Fusobacteria were found in none of the Carnoy-fixed cecal biopsies from Germany (N=400) or Russia (N=40) [Table 1].

DISCUSSION

We had previously demonstrated that the occurrence and the mean percent of Fusobacteria in mucosal lesions correlated positively with the severity of acute appendicitis when investigating material of appendices resected for acute appendicitis in Germany and suggested an infectious origin of the disease.^[1]

The present study raises nearly identical data in a group of patients undergoing surgery in a hospital in Moskau. Similar to our study in Germany, none of the biopsies taken from the inflamed regions of patients with IBD contained invasive




Figure 1: Infiltration of single epithelial cells by *Fusobacterium nucleatum* (pinned cells) in the appendix of a patient with catarrhal appendicitis (hybridization with the Fnuc Cy3 probe, orange fluorescence, $\times 400$)




Figure 2: Diffuse infiltration of the submucosa by *Fusobacterium nucleatum* in suppurative appendicitis (Fnuc Cy3 probe, orange fluorescence, $\times 400$)




Figure 3: Diffuse infiltration of the submucosa by *Fusobacterium nucleatum* (Fnuc Cy3 probe, orange fluorescence). The section is counterstained with DAPI stain, the nuclei of eukaryotic cells are seen as large blue spots

Table 1: Occurrence of Fusobacteria in the mucosa

	Suppurative appendicitis		Catarrhal appendicitis		All appendicitis		Inflamm. controls
	Carnoy (%)	Formalin (%)	Carnoy (%)	Formalin (%)	Carnoy (%)	Formalin (%)	Carnoy (%)
Germany	14/2/3/27* (70) [†]		11/2/0/25 (52)		25/4/3/52 (62)		0/400 (0)
Russia	7/2/0/11 (82)	3/1/0/11 (36)	4/0/0/12 (33)	3/0/0/12 (25)	11/2/0/23 (56)	6/1/0/23 (30)	0/40 (0)
China		4/0/0/11 (36)				4/0/0/11 (36)	

*Absolute numbers of patients with infiltrative *Fusobacterium nucleatum*/*Fusobacterium necroforum*/other Fusobacteria/all patients. [†]Percentage of patients with infiltrative Fusobacteria in the group

Fusobacteria. We have now comparatively investigated the appendices fixated simultaneously immediately after surgery in Carnoy and formalin solution. The formalin fixation proved to be detrimental for FISH. The FISH signals in formalin-fixated tissues were very weak. Nevertheless, despite the overall low accessibility of bacteria to FISH probes in formalin-fixated material, we could definitively identify *F. nucleatum* as mucosa infiltrating bacterium in 30% of formalin-fixated samples from Russia and 36% of samples from China. Our data indicates, that the invasive microbial process underlying acute appendicitis is the same in such different geographic regions as Germany, Russia, and China and is in further support of the infectious origin of acute appendicitis both in sporadic cases and in clustered outbreaks.

The study was performed according to the ethical rules included in the Declaration of Helsinki. The investigations were approved by the Institutional Review Board of the Charité Universitätsmedizin Berlin. The investigations were performed concomitantly to acute surgery using material from appendices that would have been otherwise discarded.

REFERENCES

- Swidsinski A, Dörffel Y, Loening-Baucke V, Theissig F, Rückert JC, Ismail M, *et al.* Acute appendicitis is characterized by local invasion with *Fusobacterium nucleatum*/necrophorum. *Gut* 2011;60:34-40.
- Guo Y, Xiao SY, Yan H, Sun ND, Jiang MS, Liu DY. Cluster of acute hemorrhagic appendicitis among high school students in Wuhan, China. *Am J Surg* 2004;188:115-21.
- Swidsinski A, Weber J, Loening-Baucke V, Hale LP, Lochs H. Spatial organization and composition of the mucosal flora in patients with inflammatory bowel disease. *J Clin Microbiol* 2005;43:3380-9.
- Swidsinski A, Loening-Baucke V, Vanechoutte M, Doerffel Y. Active Crohn's disease and ulcerative colitis can be specifically diagnosed and monitored based on the biostructure of the fecal flora. *Inflamm Bowel Dis* 2008;14:147-61.
- Sunde PT, Olsen I, Göbel UB, Theegarten D, Winter S, Debelian GJ, *et al.* Fluorescence *in situ* hybridization (FISH) for direct visualization of bacteria in periapical lesions of asymptomatic root-filled teeth. *Microbiology* 2003;149:1095-102.
- Amann R, Krumholz L, Stahl DA. Fluorescent-oligonucleotide probing of whole cells for determinative, phylogenetic, and environmental studies in microbiology. *J Bacteriol* 1990;172:762-70.
- Suau A, Rochet V, Sghir A, Gramet G, Brewaeys S, Sutren M, *et al.* *Fusobacterium prausnitzii* and related species represent a dominant group within the human fecal flora. *Syst Appl Microbiol* 2001;24:139-45.
- Manz W, Amann R, Ludwig W, Vancanneyt M, Schleifer KH. Application of a suite of 16S rRNA-specific oligonucleotide probes designed to investigate bacteria of the phylum cytophaga-flavobacter-bacteroides in the natural environment. *Microbiology* 1996;142:1097-106.
- Franks AH, Harmsen HJ, Raangs GC, Jansen GJ, Schut F, Welling GW. Variations of bacterial populations in human feces measured by fluorescent *in situ* hybridization with group-specific 16S rRNA-targeted oligonucleotide probes. *Appl Environ Microbiol* 1998;64:3336-45.

Source of Support: Nil, **Conflict of Interest:** None declared.