



Contents lists available at ScienceDirect

## International Journal of Surgery Case Reports

journal homepage: [www.casereports.com](http://www.casereports.com)

# Effectiveness of a barbed suture in the repair of bile duct injury during laparoscopic cholecystectomy: Report of two cases



Yusuke Takahashi\*, Naoyuki Yokoyama, Natsumi Matsuzawa, Daisuke Sato, Tetsuya Otani

Department of Digestive Surgery, Niigata City Hospital, Niigata, 950-1197, Japan

## ARTICLE INFO

## Article history:

Received 18 May 2016

Received in revised form 18 July 2016

Accepted 23 July 2016

Available online 27 July 2016

## Keywords:

Cholecystectomy

Bile duct injury

Barbed suture

## ABSTRACT

**INTRODUCTION:** Bile duct injury during laparoscopic cholecystectomy occurs in rare cases. We report two cases using barbed suture for repair in bile duct injury during laparoscopic cholecystectomies.

**PRESENTATION OF CASES:** The first patient was a 73-year-old woman who underwent elective laparoscopic cholecystectomy for cholecystolithiasis. When the gallbladder was dissected from the bed, bile spillage was observed and an injured small bile duct was detected. The bile duct could not be managed using a titanium clip. The second patient was an 83-year-old woman who underwent emergent laparoscopic cholecystectomy for gallbladder torsion. After the gallbladder was dissected from the bed, bile spillage was observed. In both cases, a running suture, of absorbable monofilament 3-0 barbed suture, was used to laparoscopically repair the injuries.

**DISCUSSION:** Laparoscopic repair of a bile duct injury is technically challenging, especially in the gallbladder bed where suturing is very difficult because of the tangential approach and the risk of additional liver laceration. Barbed sutures have the benefit of being knotless, thus, performing a running suture is not difficult, even in laparoscopic procedures. Further, absorbable and monofilament threads are generally better suited for biliary surgery, compared with non-absorbable and braided sutures, because of the potential association of the other types of materials with bile duct stone and stricture formation.

**CONCLUSION:** We believe that the V-Loc™ device is an effective and appropriate option for bile duct injuries that occur during laparoscopic cholecystectomies, particularly around the gallbladder bed, and it is especially useful for surgeons unfamiliar with intracorporeal knot tying.

© 2016 The Author(s). Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

While bile duct injuries (BDIs) that occur during laparoscopic cholecystectomies (LCs) are rare, they are a potentially serious complication. They have an incidence of approximately 0.4–1.3% [1,2]. When BDIs become apparent intraoperatively, repair should be performed laparoscopically or during conversion to open surgery. Laparoscopic repair is often difficult. In general, absorbable monofilament threads are preferable in the repair of BDIs because non-absorbable and braided suture materials may serve as the nidus for the formation of bile duct stones and biliary strictures [3,4].

Recently, knotless, barbed sutures have been used in various laparoscopic and open surgeries; however, their use in BDIs has

not been reported. This report describes the use of an absorbable, barbed suture material in the repair of a BDI that was discovered on the hepatic surface of the gallbladder bed during LC.

## 2. Presentation of cases

### 2.1. Case 1

A 73-year-old woman undergoing postoperative breast cancer treatment was found to have asymptomatic gallbladder and common bile duct stones on computed tomography (CT). After endoscopic sphincterotomy for choledocholithiasis, without any evident anatomical variation of the biliary tract in the endoscopic retrograde cholangiography, the patient underwent elective LC for cholecystolithiasis (Fig. 1).


Intraoperative cholangiography of the cystic duct did not reveal any common bile duct stones, BDI, or aberrant bile ducts.

During dissection of the gallbladder from the gallbladder bed, and after transection of the cystic duct and artery, bile spillage, without bleeding, was observed on the hepatic surface on the right side of the gallbladder body and an injured small bile duct with a


Abbreviations: BDI, bile duct injury; LC, laparoscopic cholecystectomy; CT, computed tomography.

\* Corresponding author at: Niigata City Hospital, Department of Digestive Surgery, 7-463 Shumoku, Chuo-ku Niigata City, 950-1197, Japan.

E-mail address: [niigata.takahashi@hosp.niigata.niigata.jp](mailto:niigata.takahashi@hosp.niigata.niigata.jp) (Y. Takahashi).



**Fig. 1.** Endoscopic retrograde cholangiography showed gallbladder stone (arrow), and common bile duct stones (arrow head). Anatomical variations of the biliary tract were not evident.



**Fig. 2.** Intraoperative findings: a) Bile spillage (arrow) was present in the gallbladder bed during gallbladder dissection, b) The injured small bile duct was evident (arrow head). Repair using a titanium clip was unsuccessful.

diameter of about 1 mm was detected (Fig. 2a,b). We considered performing cholangiography from the injured bile duct; however, the bile duct was too thin and narrow to perform cholangiography. We initially tried to repair the injury using a titanium clip. However, the clip could not fully close the injured bile duct owing to limited maneuverability. Moreover, we were concerned that a non-absorbable foreign body such as a titanium clip could be the nidus

for future stone formation. Thereafter, we attempted to laparoscopically suture the small bile duct using a running suture involving an absorbable, 3-0 barbed suture (V-Loc™) (Fig. 3a). The needle was inserted about 5 mm from the bile spillage, and an anchor was made through the loop at the tail of the material. A running suture from the anchor over the bile spillage point was performed, and an additional running suture back to the anchor was performed. Consequently, nine total passes of the needles across the injured bile duct were performed, and the material was cut without tying a knot. Following the repair, cessation of bile spillage was visually confirmed (Fig. 3b). The needle handling was performed so that the parenchyma of the liver was not injured. The operation was completed following placement of a drain below the gallbladder bed. The total repair time (suturing time) was 18 min.

Postoperative bile leakage was not observed, and blood tests did not detect liver dysfunction or biliary stasis. The drain was removed three days after surgery, and the patient was discharged on the following day, without further complications. Routine clinical follow-ups for the patient's breast cancer surgery, including the CT performed ten months after surgery, have not revealed any signs of biliary leakage or stasis.

## 2.2. Case 2


An 83-year-old woman with the chief complaint of fever and lumbar pain was transferred to our hospital. Abdominal CT showed findings of acute cholecystitis and gallbladder torsion with hemorrhagic necrosis and without gallbladder stones. Therefore, we performed emergent LC.

Intraoperative cholangiography was not performed owing to emergent LC and no evidence of gallbladder stones in the preoperative abdominal CT scan. During the dissection of the gallbladder from the bed, following release of the gallbladder torsion and transection of the cystic duct and artery, bile spillage without bleeding from the upper side of the gallbladder bed was observed (Fig. 4). 3-0 V-Loc™ was used laparoscopically for the repair of BDI, as in case 1. Following a running suture involving four passes of the needle across the bile spillage point, bile spillage ceased (Fig. 5). The operation was completed after drain placement below the liver. The total repair time (suturing time) was 14 min.


The drain was removed three days after surgery. She was discharged 14 days after surgery without any complication. Magnetic resonance cholangiopancreatography performed two months after surgery revealed no evidence of bile leakage or biloma.

## 3. Discussion


In both cases, during LC, we repaired BDIs at the gallbladder bed using an absorbable 3-0 barbed suture (V-Loc™). BDIs during LCs are relatively rare, but occasionally lead to life-threatening complications. The injured bile duct was observed during dissection using laparoscopic coagulating shears in both cases; therefore, these injuries might be due to thermal injury. Moreover, each bile spillage was observed with the absence of bleeding, suggesting that the injured small bile duct was not part of the Glisson system. Rather, it might have been a Type 3 or 4 subvesical bile duct, according to the reported classification system or liver parenchyma itself, particularly in case 2 [5]. When a BDI is confirmed during LC, repair is mandatory. However, laparoscopic repair is technically challenging, especially in the gallbladder bed where suturing is very difficult because of the tangential approach and the risk of additional liver laceration. Hence, according to the 12th Nationwide Survey of Endoscopic Surgery in Japan, more than half of the BDI cases observed during LCs were repaired during conversion to open surgery [6].



**Fig. 3.** Repair of bile duct injury: a) A running suture, using an absorbable 3-0 barbed suture (V-Loc™), was performed for the bile duct injury repair in the gallbladder bed, b) Bile spillage ceased after creating the running suture.



**Fig. 4.** Bile spillage (arrow) was observed at the upper side of the gallbladder bed.



**Fig. 5.** Following running suture using 3-0 V-Loc™, bile spillage ceased.

The effectiveness of barbed sutures has been recently described in gynecologic, gastrointestinal, and other types of surgeries [7–9]. These sutures have the benefit of being knotless. Thus, performing a running suture is not difficult, even in laparoscopic procedures. Above all, the barbed suture seems to be useful for surgeons unfamiliar with intracorporeal knot tying. In both cases, we used the 3-0 V-Loc™ device to laparoscopically repair a BDI at the level of the gallbladder bed, without intra- or post-operative complications. Indeed, the operators of both cases were trainees, and they were able to successfully repair the injured bile duct using 3-0 V-Loc™. To our knowledge, BDI repair using a barbed suture has not previously been reported. However, the use of this material for laparoscopic choledochotomy repair has been reported [10–12]. In these reports, this material was used for primary closure (running suture) of the common bile duct, with the safety and effectiveness of this material confirmed in both the short and long term.

Absorbable and monofilament threads are generally better suited for biliary surgery, compared with non-absorbable and braided sutures, because of the potential association of the other

types of materials with bile duct stone and stricture formation [3,4]. Thus, we believe that the V-Loc™ device is appropriate for BDI repair. Furthermore, barbed sutures, such as the one used in the present case, provide the advantage of allowing adjustable suture tension and may be easily used during laparoscopic surgery. Creating an anchor at some distance from the injured bile duct is important because the distance between the loop and the first V-Loc™ barb is approximately 1 cm. In case 1, the anchor was made approximately 5 mm from the point of injury, and a running suture was performed, involving nine needle handlings across the duct. Although more than half of the BDI cases that occur during LCs in Japan undergo open conversion, the use of absorbable barbed sutures may facilitate a future increase in the number of laparoscopic repairs performed.

#### 4. Conclusion

An absorbable monofilament barbed suture is effective and can be an option for the laparoscopic repair of BDI during LC. This mate-

rial is especially useful for surgeons unfamiliar with laparoscopic suturing.

### Consent

Written informed consent was obtained from the patients for publication of this case report and accompanying figures. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

### Author contributions

Study concept, design, and writing of this case report were by author: Yusuke Takahashi. Natsumi Matsuzawa and Naoyuki Yokoyama participated in the treatment of the patients and drafted the manuscript. Naoyuki Yokoyama critically revised the manuscript. All authors read and approved the final manuscript.

### Conflict of interest statement

Conflicts of interest: none.

### Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### Ethical approval

Ethical approval was not obtained for this case report.

### Guarantor

Yusuke Takahashi accepts full responsibility for the work and the conduct of the case report, had access to the data, and controlled the decision to publish.

### References

- [1] A. Viste, A. Horn, K. Øvrebø, B. Christensen, J.H. Angelsen, D. Hoem, Bile duct injuries following laparoscopic cholecystectomy, *Scand. J. Surg.* 104 (2015) 233–237, <http://dx.doi.org/10.1177/1457496915570088>.
- [2] W.R. Al-Kubati, Bile duct injuries following laparoscopic cholecystectomy: a clinical study, *Saudi J. Gastroenterol.* 16 (2) (2010) 100–104, <http://dx.doi.org/10.4103/1319-3767.61236>.
- [3] C. Maeda, N. Yokoyama, T. Otani, T. Katada, N. Sudo, Y. Ikeno, et al., Bile duct stone formation around a nylon suture after gastrectomy: a case report, *BMC Res. Notes* 6 (2013) 108, <http://dx.doi.org/10.1186/1756-0500-6-108>.
- [4] K.W. Sharp, C.B. Ross, V.N. Tillman, J.F. Dunn, Common bile duct healing. Do different absorbable sutures affect stricture formation and tensile strength? *Arch. Surg.* 124 (4) (1989) 408–414, <http://dx.doi.org/10.1001/arch.surg.1989.01410040018003>.
- [5] T. Schnelldorfer, M.G. Sarr, D.B. Adams, What is the duct of Luschka?—A systematic review, *J. Gastrointest. Surg.* 16 (3) (2012) 656–662, <http://dx.doi.org/10.1007/s11605-011-1802-5>.
- [6] Japan Society for Endoscopic Surgery. 12th nationwide survey of endoscopic surgery in Japan. *Japan Society for Endoscopic Surgery Journal*, 19 (2014) 498–558.
- [7] F. Alessandri, V. Remorgida, P.L. Venturini, S. Ferrero, Unidirectional barbed suture versus continuous suture with intracorporeal knots in laparoscopic myomectomy: a randomized study, *J. Minim. Invasive Gynecol.* 17 (6) (2010) 725–729, <http://dx.doi.org/10.1016/j.jmig.2010.06.007>.
- [8] S.V. Demyttenaere, P. Nau, M. Henn, C. Beck, J. Zaruby, M. Primaera, et al., Barbed suture for gastrointestinal closure: a randomized control trial, *Surg. Innov.* 16 (2009) 237–242, <http://dx.doi.org/10.1177/1553350609342988>.
- [9] C.M. Bush, J.D. Prosser, M.P. Morrison, G. Sandhu, K.H. Wenger, D.H. Pashley, et al., New technology applications: knotless barbed suture for tracheal resection anastomosis, *Laryngoscope* 122 (5) (2012) 1062–1066, <http://dx.doi.org/10.1002/lary.23229>.
- [10] K. Yokoyama, N. Tanigawa, A. Ogata, T. Nagai, M. Higashino, Laparoscopic technique and initial experiences of choledocholithotomy closure with knotless unidirectional barbed sutures after surgery for biliary stone disease, *Surg. Laparosc. Endosc. Percutan. Tech.* 25 (4) (2015) 129–133, <http://dx.doi.org/10.1097/SLE.0000000000000122>.
- [11] L.C. Fernandez, A. Toriz, J. Hernandez, N. Sanchez, E. Linares, M. Zenteno, et al., Knotless choledochorraphy with barbed suture, safe and feasible, *Surg. Endosc.* 5 (2015), <http://dx.doi.org/10.1007/s00464-015-4620-0>.
- [12] J.S. Lee, Y.C. Yoon, Laparoscopic common bile duct exploration using V-Loc suture with insertion of endobiliary stent, *Surg. Endosc.* 30 (6) (2016) 2530–2534, <http://dx.doi.org/10.1007/s00464-015-4518-x>.

### Open Access

This article is published Open Access at [sciendo.com](http://sciendo.com). It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.