

Supplemental Information

Natural Killer Cells Degenerate Intact

Sensory Afferents following Nerve Injury

Alexander J. Davies, Hyoungh Woo Kim, Rafael Gonzalez-Cano, Jahyang Choi, Seung Keun Back, Seung Eon Roh, Errin Johnson, Melanie Gabriac, Mi-Sun Kim, Jaehye Lee, Jeong Eun Lee, Yun Sook Kim, Yong Chul Bae, Sang Jeong Kim, Kyung-Mi Lee, Heung Sik Na, Priscilla Riva, Alban Latremoliere, Simon Rinaldi, Sophie Ugolini, Michael Costigan, and Seog Bae Oh

Table S1. Related to Figure 1. Primer oligonucleotide sequences.

Primers	Forward sequence	Reverse sequence
PCR: <i>Raet1a-e</i>	GCTGTTGCCACAGTCACATC	CCTGGGTCACCTGAAGTCAT
PCR: <i>NK1.1</i>	GGAACAGAGCAGAGCATTCA	CCAATCAGGGTCAGGACAAG
PCR: <i>Advillin</i>	GCTACATCGTCCTCTCGACC	CATTTCACCTCCGTGGCTT
PCR: <i>Gapdh</i>	AACAGCAACTCCCACTCTTC	TGGGTGCAGCGAACTTTAT
Real time: <i>Raet1a-e</i>	AACGGGCTGGATGATGCAC	TGGGGTAGGATCCTTGATGGT
Real time: <i>Gapdh</i>	TCCATGACAACCTTTGGCATTG	CAGTCTTCTGGGTGGCAGTGA