



Article

Effect of Wastewater Treatment on Bacterial Community, Antibiotic-Resistant Bacteria and Endoparasites

Ingrid Papajová ^{1,*}, Júlia Šmigová ¹, Gabriela Gregová ² , Jindřich Šoltys ¹ , Ján Venglovský ², Ján Papaj ³ , Tatiana Szabóová ², Nikola Dančová ², Lukáš Ihnacik ^{1,2}, Ingrid Schusterová ⁴, Jana Sušinková ⁵, Jana Raková ⁵ and Ivana Regecová ⁶

- ¹ Institute of Parasitology, Slovak Academy of Sciences, 040 01 Košice, Slovakia; bystrianska@saske.sk (J.Š.); soltys@saske.sk (J.Š.); ihnacik@saske.sk (L.I.)
 - ² Department of Public Veterinary Medicine and Animal Welfare, University of Veterinary Medicine and Pharmacy in Košice, 041 81 Košice, Slovakia; gabriela.gregova@uvlf.sk (G.G.); jan.venglovsky@uvlf.sk (J.V.); tatiana.szaboova@uvlf.sk (T.S.); nikola.dancova@student.uvlf.sk (N.D.)
 - ³ Faculty of Electrical Engineering and Informatics, Technical University in Košice, 042 00 Košice, Slovakia; jan.papaj@tuke.sk
 - ⁴ 1st Department of Cardiology, East Slovak Institute of Cardiovascular Diseases, A Joint-Stock Company St., 040 11 Košice, Slovakia; schusterovai@gmail.com
 - ⁵ Faculty of Medicine, Pavol Jozef Šafárik University in Košice, 040 11 Košice, Slovakia; jana.susinkov@upjs.sk (J.S.); jana.rakova@upjs.sk (J.R.)
 - ⁶ Department of Food Hygiene, Technology and Safety, University of Veterinary Medicine and Pharmacy in Košice, 041 81 Košice, Slovakia; ivana.regecova@uvlf.sk
- * Correspondence: papaj@saske.sk; Tel.: +421-55-633-1411



Citation: Papajová, I.; Šmigová, J.; Gregová, G.; Šoltys, J.; Venglovský, J.; Papaj, J.; Szabóová, T.; Dančová, N.; Ihnacik, L.; Schusterová, I.; et al. Effect of Wastewater Treatment on Bacterial Community, Antibiotic-Resistant Bacteria and Endoparasites. *Int. J. Environ. Res. Public Health* **2022**, *19*, 2750. <https://doi.org/10.3390/ijerph19052750>

Academic Editors:
Annalaura Carducci and
Panagiotis Karanis

Received: 15 December 2021

Accepted: 24 February 2022

Published: 26 February 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Wastewater and wastewater treatment plants serve as urban reservoirs of pathogenic microorganisms. Wastewaters frequently contain bacteria, antibiotic-resistant bacteria, and developmental stages of parasites with significant zoonotic potential. Five wastewater treatment plants in the central part of Slovakia were investigated to determine the effect of treatment on bacterial community, antibiotic-resistant bacteria, and the occurrence of helminth eggs. Although all monitored chemical factors (chemical oxygen demand, biochemical oxygen demand, N-NH₄, total nitrogen, and total phosphorus) in the effluent were in line with the legislative standards for discharge into public waterways, the results of minimal inhibitory concentrations show that reclaimed water harbors *E. coli* resistant to several commonly used antibiotics (ampicillin, piperacillin, and tazobactam, combine ampicillin and sulbactam, cefotaxime, tetracycline). The presence of endoparasite developmental stages in wastewater and sludge (*Ascaris* spp., *Hymenolepis nana*, eggs from the Ancylostomatidae family, *Giardia duodenalis*) indicates potential health risks for humans and workers at these sites. Treatment such as composting before applying sludge to land is necessary to reduce human pathogens.

Keywords: wastewater; antimicrobial resistance; *E. coli*; protozoan cysts; helminth eggs; sanitation

1. Introduction

Wastewater and wastewater treatment plants serve as urban reservoirs for pathogenic microorganisms [1,2]. Wastewaters frequently contain developmental stages of parasites with likely zoonotic potential. Many viruses and bacteria are devitalized during the treatment process in a wastewater treatment plant, but sometimes not entirely [3,4].

Frequent use of antibiotics in human and veterinary medicine leads to the spread of antibiotic-resistant bacteria into the environment. Antibiotic-resistant strains enter the environment through human feces and the liquid manure of animals. Antibiotics consumption and the increase of microorganisms' resistance can be seen by higher concentrations in wastewater, agriculture, livestock farming, and the human population [5–7]. Antibiotic-resistant bacteria and their genes are significantly emerging environmental pollutants.

The wastewater and sludge can be transmitted to agricultural land via agricultural runoff into the aquatic ecosystem and poses the risk of transmission into the food chain [8]. The prevalence of antibiotics in municipal wastewater and surface waters can lead to the development of antibiotic-resistant bacteria due to long-term exposure to low concentrations of antibiotics in the ng/L to µg/L range [9].

E. coli is typically chosen as the representative indicator of antimicrobial resistance in Gram-negative bacteria and is responsible for infections in humans and animals [10,11]. As part of the endogenous microbiota, *E. coli* can easily acquire resistance against antimicrobials consumed by humans and animals [12,13].

Antibiotic resistance is primarily caused by antibiotic use, which has led to initiatives to restrict antibiotic prescriptions and curtail antibiotic use in agriculture. Despite this knowledge, antibiotics are still the most commonly used drugs in Slovakia. According to the National Action Plan on Antimicrobial Resistance to antibiotics for outpatients in the Slovak Republic, for the period 2019–2021, antibiotic consumption was slightly higher than in other EU countries [14]. Antibiotics use in healthcare facilities is at the EU level. The significant problem is the resistance of Gram-negative microorganisms such as enterobacteria, pseudomonads, and acinetobacter to the third and higher generation of cephalosporins, fluorinated quinolone, and aminoglycosides. In Slovakia, the immediate and specific problem is the increasing resistance of enterobacteria to carbapenems. Current data on resistance in the Slovak Republic (national surveillance) are available on the website of the Public Health Authority of the Slovak Republic [15]. Furthermore, it is known that helminths cause the most parasitic infections in humans and animals among all pathogenic intestinal parasites. Helminths pose severe health risks due to their lower levels of bacteria, thus enabling very high egg survival rates and inducing resistance to common disinfectants and ultraviolet irradiation [16,17].

Helminths are also often considered as the primary constraint for the reuse of wastewater in agriculture because of their low infective dose and prolonged survival rate in the environment [18–20]. Developmental stages of endoparasites have been found in raw wastewater [20]. Wastewaters frequently contain developmental stages of parasites with zoonotic potential (e.g., *Cryptosporidium* spp., *Giardia* spp., *Toxocara* spp., *Echinococcus* spp.) [19]. It is reported that 1% of the healthy human population eliminates pathogenic agents who pass with wastewaters to wastewater treatment plants. The most prevalent in the influent segment in different treatment plants are protozoan (oo)cysts *Giardia* spp., *Cryptosporidium* spp., and *Trichostrongylus* spp., *Ascaris lumbricoides*, *Enterobius vermicularis*, *Trichuris trichiura*, and *Hymenolepis* spp. [21–28]. Many factors influence the amount: the population density, the rates of transmission, the economic status of the society, geographical regions, and climatic conditions [29–31].

The most important are eggs soil-transmitted helminths (STH). According to the World Health Organization [19], the upper limits for the STH eggs in sludge are 0.25 and 1 helminth egg/g total solids (TS), respectively. Feachem et al. [32] reported a high prevalence of STH eggs in fecal sludge in developing countries (67–735 eggs/g TS) and a lower prevalence in developed countries (2–13 eggs/g TS). According to Buitrón and Galván [16], it is essential to know if the remaining eggs after the treatment are viable because they pose a severe health risk. According to some sources, the sewage sludge also contains as many as 10^6 microorganisms in 1 mL, and about 10% is pathogenic to animals and humans [33].

Contaminated environmental components thus can lead to the spread of microorganisms, which can further directly negatively impact human and animal health. The high tenacity of the endoparasitic stages in the outer environment compared to the other microorganisms increases the health risks. For example, pathogenic viruses and bacteria survive in the external environment for several hours to days, protozoan cysts for several months up to the year, while thick-shelled helminthic eggs may remain viable for several years due to the very high resistance of the eggs to the adverse environmental conditions [34–38]. The reason is that the cellular wall of eggs contains stabilizing proteins (keratin, elastin),

lipids, and in the nematode eggs also chitin [39]. All these pathogens are potentially dangerous for public health because the infection dose is low, and in some parasites, it can be accounted for by only one egg [40–42]. Wang et al. [20] reported that conventional onsite wastewater treatment systems could potentially contribute to the transmission of infectious diseases caused by waterborne pathogenic microorganisms and become an essential human health concern. On the other hand, sewage sludge can be a valuable source of organic matter for agriculture as it contains functional agrochemical-nutrient components applicable to the soil. Sludge has a high amount of organically bound nitrogen and phosphorus [43,44]. However, it also contains components that may pose a risk to the soil, water, and the human food chain, particularly at higher concentrations. Thus, this material requires qualified processing and utilization to prevent soil contamination and avoid the accumulation of risk components in the soil.

In the Slovak Republic, the annual production of sludge reaches close to 340 thousand tons. This material is included within the category of special wastes [45], which must be safely disposed of mainly due to the presence of pathogens. On the other hand, sludge is a valuable source of nitrogen, phosphorus, potassium, and some trace elements.

This work aimed to study the effect of wastewater treatment on bacterial communities, antibiotic-resistant bacteria, and the occurrence of developmental stages of endoparasites.

2. Materials and Methods

2.1. Description of Investigated Wastewater Treatment Plants

Five wastewater treatment plants (WWTPs) located in central Slovakia were investigated to determine the effect of treatment on bacterial communities, antibiotic-resistant bacteria, and helminth eggs. This study was performed in five wastewater treatment plants that remain anonymous and are labeled as “WWTP A–E”. All wastewater treatment plants receive municipal wastewater and discharge the effluents into streams or rivers. The characteristics of studied wastewater treatment plants are described in Table 1. The generated sludge was treated by aerobic stabilization (WWTP B) or anaerobic stabilization (WWTP C) and applied to the compost. Sludge from WWTP A, D, and E was not treated in the plants. Unstabilized raw sludge in WWTP B was treated by aerobic stabilization. WWTP B was shut down during the monitored period, so obtaining samples from the influent portion was not possible.

Table 1. Characteristics of the wastewater treatment plants studied.

	WWTP				
	A	B	C	D	E
Number of residents	630	7529	87,126	454	881
Discharge (m ³ /day)	550.8	3299.6	24,698.1	112.1	73.9
Discharge (l/s)	6.38	38.06	283.41	1.29	0.86
Recipient river	Turieč	Teplica	Váh	Vrúca	Blatnický potok
Cleaning method			Mechanical and biological		
Treatment of sludge	Unstabilized sludge (treatment at WWTP B)	Aerobic stabilization, sludge applied to the compost	Anaerobic stabilization, sludge applied to the compost	Unstabilized sludge (treatment together WWTP B)	Unstabilized sludge (treatment at WWTP B)
Sewage type	Domestic	Domestic	Domestic	Domestic	Domestic

WWTP—wastewater treatment plant.

2.2. Sample Collection

Samples were collected in the summer of 2020 (June–August). During the tested period, 9 samplings were performed and 3 samples were taken from each sampling place (influent, effluent, sludge).

Two liters of influent and effluent samples were collected to chemically clean bottles for the physicochemical evaluation, and 1 L was placed into a sterile bottle for microbiological and parasitological examination. A total of 2 L of raw sludge and 1 kg of treated sludge

were collected. Samples were stored without any conservation at 4 °C and transferred to the laboratory for microbiological, parasitological, and chemical examination, performed within 24–48 h.

2.3. Chemical Examination of Samples

Chemical examination of wastewater (input and output) and sewage sludge included determination of pH, the levels of total nitrogen (N_t), water-soluble ammonium (NH_4-N), total phosphorus (P_t), and chemical oxygen demand (COD_{Mn}). According to STN ISO 10523 [46], the pH was determined with a pH-meter (HACH Company, Loveland, CO, USA) and a WATERPROOF pH Tester 30. COD_{Mn} was determined by oxidation with $KMnO_4$ according to STN EN ISO 8467 [47] and NH_4-N by titration [48,49]. A portion of samples for N_t determinations was digested using a HACH-Digesdahl apparatus (HACH Company, Loveland, CO, USA). N_t was distilled with NaOH (40%) [50,51], and P_t was determined by the vanadomolybdate method [52,53]. Efficiency of chemical pollutants was calculated according to the formula:

$$\% \text{ of efficiency} = \left(\frac{\text{concentration of pollutant in influent} - \text{concentration of pollutant in effluent}}{\text{concentration of pollutant in influent}} \right) \times 100\% \quad (1)$$

According to National Regulation SR [54], the biological process of wastewater treatment is focused only on the reduction of chemical pollutants, which have to meet the maximal limit of chemical factors in the effluent grab samples of wastewater discharged as described in Table 2.

Table 2. Maximal limit of chemical factors in the effluent grab samples of wastewater discharged to the recipient rivers [54].

Number of Recipient	N_t (mg/L)	NH_4-N (mg/L)	P_t (mg/L)	COD_{Mn} (mg/L)	IS (mg/L)
up to 50	–	–	–	–	–
51–2000	–	–	–	170	60
2001–10,000	–	40	–	170	50
10,001–25,000	40	30	–	140	50
25,001–100,000	30	20	5	125	40
over 100,000	25	10	4	125	40

N_t —total nitrogen; NH_4-N —water-soluble ammonium nitrogen; P_t —total phosphorus; COD_{Mn} —chemical oxygen demand; IS—insoluble substances; – means no maximal limit.

2.4. Microbiological Examination of Samples

Isolation of Bacteria

The influent, effluent, and sludge wastewater samples were prepared by a serial dilution from 10^1 to 10^6 . All samples were examined in duplicate. Determination of relevant bacterial counts (total count of bacteria, coliform bacteria, fecal coliform bacteria, and fecal enterococci) was carried out in compliance with the Slovak Republic Government Regulation [55]. The total bacterial count (TCB) was determined according to STN EN ISO 6222 [56]. The pour-plate method on meat-peptone agar was followed by aerobic cultivation at 37 °C for 24 h for bacteria determination.

Coliform bacteria (CB) and fecal coliform bacteria (FCB) were cultivated according to STN EN ISO 9308-1 [57] using Endo agar (HiMedia, Mumbai, India) and incubated for 24 h at 37 and 43 °C, respectively. Then, the characteristic colonies were counted. A lactose fermentation test confirmed coliform bacteria presence.

Determination of fecal enterococci counts (FE) was carried out according to STN EN ISO 7899-2 [58]. The cultivation of respective bacteria was performed on Slanetz Bartley agar (Merck, Darmstadt, Germany) at 37 °C for 48 h.

2.5. *E. coli* Isolation

The suspect *E. coli* colonies from Endoagar (or Chromagar, McConkey agar) were identified by biochemical ENTEROtest 24 (Erba Lachema, Brno, Czech Republic), intended to identify the critical species of the family Enterobacteriaceae. Pure 24 h bacterial culture was inoculated in sterile saline solution. The homogenized suspension must have turbidity equal to No. 1 of the McFarland turbidity scale. The microtitration plates with 24 biochemical tests (urease, arginine, ornithine, lysine, hydrogen sulfide, simmons citrate, malonate, β -galactosidase, salicin, sorbitol, melibiose, cellobiose, lactose, trehalose, mannitol, β -glucuronidase, dulcitol, adonitol, arabitol, sucrose, inositol, raffinose, esculin, β -xylosidase) were inoculated with 0.1 mL of culture suspension. After 24 h of incubation at 37 °C, microtitration plates were analyzed for color reaction by reader ErbaScan and identification program ErbaExpert. The identification was supplemented by the paper strip tests: OXItest, COLItest, and PYRAtest (Erba Lachema, Brno, Czech Republic).

2.6. Determination of Minimal Antibiotics Inhibitory Concentrations in *E. coli* Isolates

A total of 142 isolates of *E. coli* (one sample, one strain) were analyzed for antibiotic susceptibility. Minimal inhibitory concentrations (MIC) were determined according to VET01-S2 [59] and EUCAST [60] by a Miditech system (Miditech, Bratislava, Slovakia) with the interpretative reading of MIC [61]. MIC GX expresses geometric mean MIC values for an antibiotic agent (mg/L) in *E. coli* isolates. The antibiotics used in the presented study were as follows: ampicillin (AMP), ampicillin and sulbactam (SAM), piperacillin + tazobactam (TZP), cefuroxime (CXM), cefotaxime (CTX), ceftazidime (CAZ), ceftazidime with clavulanic acid (CAC), cefoperazone and sulbactam (SPZ), cefepime (FEP), ertapenem (ETP), meropenem (MEM), gentamicin (GEN), tobramycin (TOB), amikacin (AMI), ciprofloxacin (CIP), tetracycline (TET), tigecycline (TGC), chloramphenicol (CHL), colistin (COL), trimethoprim and sulfonamides (COT) and nitrofurantoin (NIT). The antibiotic reading system automatically identifies common resistance mechanisms, such as “ESBL”, extended spectrum β -lactamases; “Multiresistance!”, current resistance in 3 or more unrelated ATB groups; “AAC(6’), AAC(3’), ANT(2’)”, enzyme-modifying aminoglycosides in G-bacteria; AGL AAC(6)—enzym AAC(6) modifying of aminoglycosides; “TEM-1, -2, SHV-1—low”, common beta-lactamases, with low enzyme expression; “TEM-1, -2, SHV-1—high”, common beta-lactamases with high enzyme expression; “Plasmidic AmpC!”, plasmid-transferred Amp C beta-lactamases; “Class A, Amp C, hyp.!”, derepressed chromosomal beta-lactamases!; “Carbapenem resistance”, resistance to carbapenems; Penicilinase high, penicilinase with high enzyme expression; Penicilinase low, penicilinase with low enzyme expression, etc.

2.7. Parasitological Examination of Samples

To determine protozoa (oo) cysts and helminth eggs count in influent and effluent samples from WWTP and sewage sludge, 50 mL from each 1 L sample was taken and examined by sedimentation and flotation techniques [62].

The dry sludge samples were surveyed according to Kazacos [63]. Briefly, 100 g of the pooled sludge sample, 100 mL of water, and 0.5 mL of Tween 40 were mixed and decanted for 10 min. Subsequently, the samples were sieved and replenished with 1000 mL of water. After 1 h of sedimentation, the soil samples were centrifuged (Eppendorf 5804, Hamburg, Germany) and then floated with sucrose flotation solution (specific density of 1.3). Samples were examined under the light microscope at 20 \times and 40 \times magnification (Leica Microsystems, DM 5000B light microscope, Wetzlar, Germany) to detect the presence of protozoa (oo) cysts and helminth eggs.

3. Results

WWTPs (A, D, E) are small, representing up to 10,000 residents. Here, the regularly monitored value is only one chemical indicator, i.e., CODMn (70 mg/L in grab samples or 135 mg/L in composite samples). In WWTP D and WWTP E, these values exceeded

the norm (180 mg/L) and in WWTP E, this value exceeded the norm more than twice. Furthermore, in the biggest WWTP C, the COD_{Mn} was up to 282.8 mg/L, where only 125 mg/L is officially allowed. WWTP B was in reconstruction, and wastewater flowed directly to the river, and the COD level was exceeded twice. Here the maximal limit was 140 mg/L (Table 3).

Table 3. Physico-chemical analysis of wastewater before and after treatment in WWTPs.

	N _t (mg/L)	NH ₄ -N (mg/L)	P _t (mg/L)	COD _{Mn} (mg/L)	pH
WWTP A					
Influent	22.41	14.01	17.02	214.84	7.13
Effluent	14.01	3.50	1.05	152.65	7.01
Sludge	249.33	11.21	32.76	290.30	6.70
Efficiency	37.5%	75.0%	93.8%	28.9%	
WWTP B					
Influent	44.83	15.41	26.37	478.40	7.30
WWTP C					
Influent	39.22	51.83	67.82	537.35	7.45
Effluent	12.61	3.50	2.71	282.80	7.34
Sludge	424.41	11.80	129.65	2467.475	6.78
Efficiency	67.9%	93.2%	96.0%	47.4%	
WWTP D					
Influent	39.22	26.61	108.14	348.95	7.26
Effluent	18.21	4.90	2.57	180.41	7.60
Sludge	260.53	21.01	86.20	863.47	6.92
Efficiency	53.6%	81.6%	14.4%	48.3%	
WWTP E					
Influent	103.65	81.24	133.96	936.90	7.93
Effluent	82.64	35.02	5.36	414.83	7.16
Sludge	140.07	43.42	71.79	623.22	6.95
Efficiency	20.3%	56.9%	96.0%	55.7%	

WWTP—wastewater treatment plant; N_t—total nitrogen; NH₄-N—water-soluble ammonium nitrogen; P_t—total phosphorus; COD_{Mn}—chemical oxygen demand; IS—insoluble substances.

After mechanical and biological treatment in WWTPs, the effluent's chemical factors decreased within the limit in every monitored wastewater treatment plant.

The results of the microbial evaluation and the percentage of efficiency in the individual examined wastewater treatment plants are described in Figure 1. The log₁₀ in the total number of microorganisms in individual wastewater treatment plants ranged from 7 to 5.4 in influent and from 6 to 3.4 in the effluent, representing a decrease from 99 to 89%. The total coliform bacteria concentration in the sludge varied from 7.6 to 5.7. The log₁₀ of coliform bacteria ranged between 6.9 to 4 in the influent and in the effluent from 6.4 to 3.4, representing a 68 to 99% purification efficiency. The fecal coliform bacteria were 4.5 to 6 at the influent and from 2.6 to 4.9 in the effluent to express 95 to 99% efficiency. The log₁₀ of fecal enterococci ranged from 4 to 4.4 in the influent and 2.2 to 4.2 in the effluent, thus showing the efficiency from 79 to 99%. The concentration of the monitored bacteria in the sludge was approximately the same as in the samples from the wastewater influent.

A modified microdilution method with the VetMIC panel detected antimicrobial resistance in 142 *E. coli* strains. The highest incidence of beta-lactamase resistance was observed for ampicillin (60% and MIC GX 18.9 mg/L in sludge samples), which was followed by ampicillin and sulbactam (35.21% and MIC GX 7.8 mg/L), tetracycline (20% in influent and 40% in effluent, MIC GX 3.8 mg/L), piperacillin and tazobactam (8.6%, MIC GX 2.2 mg/L in influent). MICs of cefuroxime and cefotaxime were 5.8 in influent and

5.6 mg/L (in effluent); tigecycline, trimethoprim and sulfonamides was about 5.6% (in effluent). The percentage of resistance to ceftazidime, tobramycin, amikacin, ciprofloxacin was 2.87% in influent and effluent components, and sludge showed low resistance to those antibiotics. Resistance to meropenem, sulbactam, cefepime, meropenem, gentamicin, and nitrofurantoin was 0% in all wastewater and sludge samples (Figure 2).




Figure 1. Comparison of bacterial contamination (log₁₀) and treatment efficiency in four WWTPs (CB—coliform bacteria, FCB—fecal coliform bacteria, FE—fecal enterococci, TCB—total count bacteria).




Figure 2. Comparison of MIC GX and percentages of resistance in influent, effluent, and sludge from the WWTPs.

Among all 142 investigated *E. coli* strains, the high penicillinase phenotype was confirmed in 9 strains (12.68%) and 11.27% of carbapenemases (detected in two strains from the influent portion). Three phenotypes were penicillinase low (4.23%). An AGL

AAC(6') phenotype was also detected in samples. Multi-drug resistance was confirmed in one sample (Figure 3). Based on the MIC phenotypic identification of *E. coli* strains, 12.5% were high-level resistance penicillinase occurrence, 4% low-level, 1.4% were multiresistant isolates, and 1.4% were aminoglycosides AAC(6')I. The ESBLs were not detected in isolated *E. coli* strains.




Figure 3. The phenotype of resistance in *E. coli* strains.

The results obtained from the parasitological examination are presented in Table 4. *Giardia duodenalis* was detected only in WWTP C influent section collected in June 2020. No other cysts were observed by microscopy in water or the sludge.

Table 4. Parasitological examination of wastewater before and after treatment in WWTPs.

WWTP	Sample	Eggs/Oocysts
WWTP A	Influent	<i>Ascaris</i> spp., <i>Hymenolepis nana</i> , family Ancylostomatidae
	Effluent	Neg.
	Sludge	<i>Ascaris</i> spp., family Ancylostomatidae
WWTP B	Influent	<i>Hymenolepis nana</i> , <i>Ascaris</i> spp., family Ancylostomatidae
WWTP C	Influent	<i>Hymenolepis nana</i> , <i>Giardia duodenalis</i>
	Effluent	Neg.
	Sludge	<i>Ascaris</i> spp., family Ancylostomatidae
WWTP D	Influent	Neg.
	Effluent	Neg.
	Sludge	<i>Ascaris</i> spp., family Ancylostomatidae
WWTP E	Influent	<i>Ascaris</i> spp.
	Effluent	Neg.
	Sludge	family Ancylostomatidae

There were no helminth eggs found in effluent portions. *Ascaris* spp. eggs were confirmed in influent from WWTP A, B, and E. No *Ascaris* spp. eggs were detected in influent from WWTP C and D. It was impossible to differentiate between the species since only the optical microscopy detection was carried out. *Hymenolepis nana* eggs and eggs from the family Ancylostomatidae were also found in the influent portion. The most prevalent helminth eggs in sludge were *Ascaris* spp. eggs and eggs from the family Ancylostomatidae (Table 4).

4. Discussion

The treatment process in the WWTPs is primarily focused on the removal of chemical pollutants (COD, BOD5, N-NH4, total nitrogen, and total phosphorus) [54], which are checked in effluent values as a preventive measure against river pollution (eutrophication of water) negatively impacting river ecosystems [42,64].

During the studied period, almost all chemical factors in effluent from monitored WWTPs were in line with the standards and regulations for discharge into public waterways.

COD, the measure of the oxygen equivalent of the organic matter and microorganisms in the wastewater, was higher in smaller and bigger WWTPs and exceeded the maximum level of 125 mg/L. In untreated domestic wastewater, COD usually ranges between 250 and 1000 mg/L [65].

By comparing the results of chemical parameters and microbiological analyses in the effluent, the higher efficiency of wastewater treatment in WWTP A, C, and D is visible. Almost all monitored microorganisms here decreased by 2 to 3 logarithmic orders. The concentration of microorganisms in the sludge in individual WWTPs was comparable or slightly higher than at the influent. A comparison of the number of microorganisms in the sludge of the monitored WWTPs found no significant differences (ranging from 6–7 log in TCB, CB, and 4.5–6 log in FCB and FE).

The number of coliform bacteria in effluent ranged from 3.4 to 6.3 log CFU·mL⁻¹ which correlates with results of other works investigating wastewater in Slovak and Czech Republic (3.02–4.94 log CFU·mL⁻¹) [66]. In effluent wastewater from hospitals, the concentrations of a number of coliform bacteria can reach 7.18 log CFU·mL⁻¹ [67], which is in two orders higher when compared to the domestic and municipal WWTP, highlighting the importance of separated wastewater and sludge collection and treatment.

Interestingly, the effectiveness of the treatment process in small WWTP (up to 2000 recipient) is less than in big WWTP (up to 100,000 recipient). For example, in the WWTP E, the wastewater from 900 recipients had the highest concentration in chemical parameters and microbial concentration in influent decreased only by 0.5 log order. This signals low effectiveness of cleanability and possible failure in the treatment process. In addition, larger WWTPs employ more professional staff, the cleaning process is regularly controlled and inspected (including self-inspections), and the monitored parameters in the effluent are set up to be more strict.

Although the success of treatment by comparing the concentration of chemical parameters, microorganisms and parasites in effluent has been relatively high, it still poses a risk to rivers' pollution and habitats. It is not proven that the treatment is sufficient to remove antibiotic-resistant bacteria. It is known that some of these bacteria can carry resistance genes either in their core genome or on mobile genetic elements. These genes can be transferred from one bacterium to another via horizontal gene transfer under selection pressure [68]. There are also possibilities for plasmid transfer in wastewater treatment plants and surface waters [69,70].

In a recent study, the following *E. coli* resistances were detected: more than 50% to ampicillin, 35% were resistant to ampicillin and sulbactam, 20% to tetracycline, and 8.6% to piperacillin and tazobactam. *E. coli* isolates from the effluent also showed resistance to cefotaxime, cefepime, cefuroxime, tigecycline, trimethoprim, and sulfonamides (around 5%). There were only minor differences between antibiotic resistance results from influent, effluent, and sludge.

We confirmed *E. coli* strains the Penicillinase: high, carbapenemase, Penicillinase: low, and AGL AAC(6') phenotypes. A multidrug-resistant *E. coli* phenotype was detected in one sample.

In a previous study, Gregová and Kmet' [5] detected that the production of extended-spectrum β -lactamases in coliform isolates was encoded mainly by blaTEM, blaCTX-M-2, and blaCTX-M-8/25 genes. About 62% of resistant strains contained a combination of two or more extended-spectrum beta-lactamases (ESBL) genes.

Similarly, Reinthaler et al. [71] determined a 200-fold reduction of *E. coli* in WWTP effluent samples (from 10⁴ CFU·mL⁻¹ to approx. 10² CFU·mL⁻¹). *E. coli* strains were resistant to the penicillin group (ampicillin 18%, piperacillin 12%), cephalosporin group (cephalothin 35%, cefuroxime 11%), quinolones (nalidixic acid 15%), trimethoprim/sulfamethoxazole (13%), and tetracycline (57%).

In untreated sludge, Redhead et al. [72] found higher resistance of *E. coli* isolates to amoxicillin, trimethoprim, ciprofloxacin, and gentamicin. After thermal hydrolysis at WWTPs, the absolute abundance was markedly reduced in *dfrA1*, *dfrA5*, *dfrA7*, *aac(3)-1*, CTX-M-1, CTX-M-9, *bla-Imp*, *qnrS*, *tetM*, *sul1*, and *intl1* genes, which is similar to the data observed by Martins et al. [73] and Raven et al. [74].

Lépesová et al. [67] have found that more than half of the coliform bacteria from hospital wastewater were multidrug resistant and possessed a strong biofilm-formation ability.

A high concentration of the monitored bacteria and antibiotic-resistant *E. coli* isolates was found in the sludge, which still poses a risk of spreading microorganisms through the direct incorporation of sludge into agricultural land. However, the above indicates that sludge produced during municipal wastewater treatment must be subjected to additional processing to improve its hygiene level. Therefore, composting constitutes an economically advantageous and hygienically effective method of sludge processing. In addition, processing and organic recycling wastes by composting must ensure a product that fulfills a range of water management and hygiene requirements and complies with the principles of waste legislation [45].

Still, wastewater treatment plants play a vital role in minimizing the discharge of many water pollutants, including antibiotics [75,76]. Degradation of hormones, pesticides, antibiotics, antihistamines, and drugs is limited, and as a result, they commonly end up in the aquatic environment. Runoff and wastewater discharges may also contribute to the spread of organic and inorganic nutrients that may boost the growth and proliferation of indigenous or introduced pathogens.

Ascaris spp. eggs, viable eggs from the Ancylostomatidae family, and viable *H. nana* eggs were found in influent. This indicates that human feces and domestic animal excreta might contaminate wastewater treatment systems. During mechanical and biological treatment processes in the studied WWTPs, mainly bacteria, but not the endoparasites' developmental stages, were devitalized. Therefore, all above-mentioned helminths egg sediment and concentrate were in the sludge with suspended particles. In the case of poor sludge treatment, they again re-enter the environment and pose potential hygiene, epidemiological or epizootiological risks [19,36,77]. Finally, we can state that the helminth eggs in the sludge indicate the overall health status and the incidence of diseases in the population monitored in the WWTP area.

An exciting finding was confirmation of *G. duodenalis* cysts influent. No cysts were found in the effluent stage and sludge. However, information about the presence of protozoan pathogens is limited due to the inconsistencies in sampling as well as concentration and recovery procedures [44]. Despite our findings, it is necessary to pay special attention to this parasite because it has great zoonotic potential. *G. duodenalis* (syn. *G. intestinalis*, *G. lamblia*) is a unicellular parasite causing gastrointestinal disorders in a wide range of hosts, including wild and companion animals as well as humans. It is one of the most spread parasitic diseases in man, who can become infected via the fecal–oral route through contaminated water (“water-borne”) and food (“food-borne”), or by direct transmission from host to host. This parasitic disease is classified as neglected and occurs more frequently in areas with poor hygiene standards, where children are the most affected population.

Similarly, Dudlová et al. [78] in east Slovakia studied the incidence of endoparasite germ's developmental stages (cysts, oocysts, protozoa, and helminth eggs) as an indirect monitoring factor for endoparasite circulation in raw municipal wastewater, sludge, and biologically cleaned wastewater. The raw wastewater contained cysts of *Giardia* spp., and *Entamoeba* spp., the helminth eggs *Ascaris* spp., and strongyle-type eggs. No protozoa cysts or helminth eggs were found in the treated wastewater. The highest occurrence of endoparasite developmental stages was detected in drained stabilized sludge. Protozoan (oo) cysts (*Giardia* spp., *Cryptosporidium* spp., *Entamoeba* spp.) and helminth eggs (*Ascaris* spp., *Trichuris* spp., *Taenia* spp., *Hymenolepis* spp., or strongyle-type eggs) were found. In drained and stabilized sludge, the eggs of *Capillaria* spp. and *Toxocara* spp. were also detected. In comparison with our results, differences in parasite composition can be

because wastewater treatment plants in eastern Slovakia are located close to territories with low hygiene standards and a significant presence of residents belonging to marginalized populations. On the other hand, Amoah et al. [79], Chaoua et al. [80], and others have reported that wastewater treatment plants are not fully capable of helminth eggs removal from the water.

We confirmed that sedimentation is an effective removal mechanism for the helminth eggs. Similar results are reported by Kansime et al. [81] and Dai et al. [82]. In the low-quality water, the eggs are incorporated into particle flocs with different settling velocities, and the settling rate of eggs and particles is closely associated [83]. Environmental conditions can also affect the survival of the pathogen in wastewater. For example, the rate of removing bacterial indicators (e.g., *Salmonella* and fecal coliform bacteria) is higher in summer than in winter [84,85]. UV radiation by sunlight is an effective mechanism for pathogen removal, mainly in open-water treatment wetlands. Temperature, pH, insolation, and solar radiation influence the survival of *A. suum* eggs [86–88].

Based on our results, we can conclude that the use of untreated or insufficiently treated wastewater poses many risks. The use of untreated wastewater is also associated with a higher infection intensity, especially for helminth infection. Ensink et al. [89] found a significantly increased risk of *A. lumbricoides*, hookworm, and *T. trichiura* infection in farming communities irrigating with wastewater. Aquatic recipients contaminated with wastewater containing helminth eggs also have a high concentration of suspended particles resulting in flocculation of the suspended material, including eggs [83].

5. Conclusions

Wastewater treatment plants play a vital role in minimizing the discharge of many water pollutants and the protection of the environment. The results show that the pattern of resistance in *E. coli* isolates found in monitored WWTPs is comparable with resistance seen at the clinical health care level. Ampicillin (beta-lactams), gentamicin (aminoglycoside), and ciprofloxacin (fluoroquinolones) are frequently prescribed drugs in Slovakia.

However, the main wastewater and sludge reuse limitation is contamination with helminth eggs due to its low infective dose and prolonged survival in the environment. In this study, several parasite species were identified in influent segments and sludge, where nematodes of human and animal origin were predominantly identified. In addition, the wastewater and sludge indicates potential health risks for humans and workers at these sites.

Author Contributions: Conceptualization, I.P., J.Š. (Júlia Šmigová) and G.G.; methodology, I.P., J.Š. (Júlia Šmigová) and J.V.; software, J.P.; investigation, J.Š. (Júlia Šmigová), L.I., T.S., N.D. and I.R.; resources, I.P.; data curation, J.Š. (Júlia Šmigová), G.G. and J.P.; writing—original draft preparation, I.P. and G.G.; writing—review and editing, J.Š. (Jindřich Šoltys), I.S., J.S. and J.R.; supervision, I.P.; project administration, I.P.; funding acquisition, I.P. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by grant number VEGA 2/0138/21.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data are available from the corresponding author upon reasonable request.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Onda, K.; LoBuglio, J.; Bartram, J. Global access to safe water: Accounting for water quality and the resulting impact on MDG progress. *Int. J. Environ. Res. Public Health* **2012**, *9*, 880–894. [[CrossRef](#)] [[PubMed](#)]
2. Bain, R.E.; Gundry, S.W.; Wright, J.A.; Yang, H.; Pedley, S.; Bartram, J.K. Accounting for water quality in monitoring access to safe drinking-water as part of the Millennium Development Goals: Lessons from five countries. *Bull. World Health Organ.* **2012**, *90*, 228–235. [[CrossRef](#)] [[PubMed](#)]

3. Okoh, A.I.; Sibanda, T.; Gusha, S.S. Inadequately treated wastewater as a source of human enteric viruses in the environment. *Int. J. Environ. Res. Public Health* **2010**, *7*, 2620–2637. [CrossRef] [PubMed]
4. Čornejová, T.; Venglovský, J.; Gregová, G.; Kmeťová, M.; Kmeť, V. Extended spectrum beta-lactamases in *Escherichia coli* from municipal wastewater. *Ann. Agric. Environ. Med.* **2015**, *22*, 447–450. [CrossRef] [PubMed]
5. Gregová, G.; Kmeť, V. Antibiotic resistance and virulence of *Escherichia coli* strains isolated from animal rendering plant. *Sci. Rep.* **2020**, *10*, 17108. [CrossRef] [PubMed]
6. Gregová, G.; Kmeť, V.; Szabóová, T. New insight on antibiotic resistance and virulence of *Escherichia coli* from municipal and animal wastewater. *Antibiotics* **2021**, *10*, 1111. [CrossRef] [PubMed]
7. Manyi-Loh, C.; Mamphweli, S.; Meyer, E.; Okoh, A. Antibiotic use in agriculture and its consequential resistance in environmental sources: Potential public health implications. *Molecules* **2018**, *23*, 795. [CrossRef]
8. Elahi, E.; Abid, M.; Zhang, L.; Alugongo, G.M. The use of wastewater in livestock production and its socioeconomic and welfare implications. *Environ. Sci. Pollut. Res.* **2017**, *24*, 17255–17266. [CrossRef]
9. Kraemer, S.A.; Ramachandran, A.; Perron, G.G. Antibiotic pollution in the environment: From microbial ecology to public policy. *Microorganisms* **2019**, *7*, 180. [CrossRef]
10. Payment, P.; Waite, M.; Dufour, A. Introducing parameters for the assessment of drinking water quality. *Assess. Microb. Saf. Drink. Water Improv. Approaches Methods* **2003**, *4*, 47–77.
11. Simpson, J.M.; Santo Domingo, J.W.; Reasoner, D.J. Microbial source tracking: State of the science. *Environ. Sci. Technol.* **2002**, *36*, 5279–5298. [CrossRef] [PubMed]
12. Handrova, L.; Kmet, V. Antibiotic resistance and virulence factors of *Escherichia coli* from eagles and goshawks. *J. Environ. Sci. Health Part B* **2019**, *54*, 605–614. [CrossRef] [PubMed]
13. Drugdova, Z.; Kmet, V. Prevalence of β -lactam and fluoroquinolone resistance, and virulence factors in *Escherichia coli* isolated from chickens in Slovakia. *Biologia* **2013**, *68*, 11–17. [CrossRef]
14. ECDC. *National Action Plan on Antimicrobial Resistance in the Slovak Republic for the Period 2019–2021*; ECDC: Bratislava, Slovakia, 2018. (In Slovak)
15. Public Health Authority of the Slovak Republic. Available online: <http://www.snars.sk> (accessed on 5 December 2021).
16. Buitrón, G.; Galván, M. Effect of compression–decompression on helminth eggs present in sludge of a settling tank. *Water. Res.* **1998**, *32*, 1708–1712. [CrossRef]
17. Mun, S.; Cho, S.H.; Kim, T.S.; Oh, B.T.; Yoon, J. Inactivation of *Ascaris* eggs in soil by microwave treatment compared to UV and ozone treatment. *Chemosphere* **2009**, *77*, 285–290. [CrossRef] [PubMed]
18. Paruch, A.M. Possible scenarios of environmental transport, occurrence and fate of helminth eggs in light weight aggregate wastewater treatment systems. *Rev. Environ. Sci. Biotechnol.* **2010**, *9*, 51–58. [CrossRef]
19. World Health Organization. *Guidelines for the Safe Use of Wastewater Excreta and Greywater*; World Health Organization: Geneva, Switzerland, 2006; Volume 4.
20. Wang, M.; Zhu, J.; Mao, X. Removal of pathogens in onsite wastewater treatment systems: A review of design considerations and influencing factors. *Water* **2021**, *13*, 1190. [CrossRef]
21. Heitman, T.L.; Frederick, L.M.; Viste, J.R.; Guselle, N.J.; Morgan, U.M.; Thompson, R.C.; Olson, M.E. Prevalence of *Giardia* and *Cryptosporidium* and characterization of *Cryptosporidium* spp. isolated from wildlife, human, and agricultural sources in the North Saskatchewan River Basin in Alberta, Canada. *Can. J. Microbiol.* **2002**, *48*, 530–541. [CrossRef]
22. Caccio, S.M.; De Giacomo, M.; Aulicino, F.A.; Pozio, E. *Giardia* cysts in wastewater treatment plants in Italy. *Appl. Environ. Microbiol.* **2003**, *69*, 3393–3398. [CrossRef]
23. Sulaiman, I.M.; Jiang, J.; Singh, A.; Xiao, L. Distribution of *Giardia duodenalis* genotypes and subgenotypes in raw urban wastewater in Milwaukee, Wisconsin. *Appl. Environ. Microbiol.* **2004**, *70*, 3776–3780. [CrossRef]
24. Leelayoova, S.; Siripattanapipong, S.; Thathaisong, U.; Naaglor, T.; Taamasri, P.; Piyaraj, P.; Mungthin, M. Drinking water: A possible source of *Blastocystis* spp. subtype 1 infection in schoolchildren of a rural community in central Thailand. *Am. J. Trop. Med. Hyg.* **2008**, *79*, 401–406. [CrossRef] [PubMed]
25. Ajonina, C.; Buzie, C.; Ajonina, I.U.; Basner, A.; Reinhardt, H.; Gulyas, H.; Liebau, E.; Otterpohl, R. Occurrence of *Cryptosporidium* in a wastewater treatment plant in North Germany. *J. Toxicol. Environ. Health A* **2012**, *75*, 1351–1358. [CrossRef] [PubMed]
26. Kitajima, M.; Haramoto, E.; Iker, B.C.; Gerba, C.P. Occurrence of *Cryptosporidium*, *Giardia*, and *Cyclospora* in influent and effluent water at wastewater treatment plants in Arizona. *Sci. Total Environ.* **2014**, *484*, 129–136. [CrossRef] [PubMed]
27. Sangbari, N.; Dadban Shahamat, Y.; Abbasinejat, Z.; Sharbatkhori, M.; Rostami, M. Survey of parasitic contamination of sewage sludges in northern Iran. *J. Appl. Sci. Environ. Manag.* **2018**, *22*, 1277–1280. [CrossRef]
28. Stott, R.; May, E.; Mara, D.D. Parasite removal by natural wastewater treatment systems: Performance of waste stabilisation ponds and constructed wetlands. *Water Sci. Technol.* **2003**, *48*, 97–104. [CrossRef]
29. Jimenez, B.; Chavez, A.; Leyva, A.; Tchobanoglous, G. Sand and synthetic medium filtration of advanced primary treatment effluent from Mexico City. *Water Res.* **2000**, *34*, 473–480. [CrossRef]
30. Ben Ayed, L.; Schijven, J.; Alouini, Z.; Jemli, M.; Sabbahi, S. Presence of parasitic protozoa and helminth in sewage and efficiency of sewage treatment in Tunisia. *Parasitol. Res.* **2009**, *105*, 393–406. [CrossRef] [PubMed]
31. Paruch, A.M.; Mæhlum, T.; Robertson, L. Changes in microbial quality of irrigation water under different weather conditions in Southeast Norway. *Environ. Process.* **2015**, *2*, 115–124. [CrossRef]

32. Feachem, R.G.; Bradley, D.J.; Garelick, H.; Mara, D.D. *Sanitation and Disease Health Aspects of Excreta and Wastewater Management*; John Wiley & Sons Edition: New York, NY, USA, 1983; p. 534.
33. Sasáková, N.; Juriš, P.; Papajová, I.; Vargová, M.; Ondrašovičová, O.; Ondrašovič, M.; Kasková, A.; Szabová, E. Parasitological and bacteriological risks to animal and human health arising from waste-water treatment plants. *Helminthologia* **2005**, *42*, 137–142.
34. Gaspard, P.G.; Wiart, J.; Schwartzbrod, J. Urban sludge reuse in agriculture: waste treatment and parasitological risk. *Bioresour. Technol.* **1995**, *52*, 37–40. [[CrossRef](#)]
35. Nelson, K.L.; Darby, J. Inactivation of viable *Ascaris* eggs by reagents during enumeration. *Appl. Environ. Microbiol.* **2001**, *67*, 5453–5459. [[CrossRef](#)] [[PubMed](#)]
36. Lalander, C.; Diener, S.; Magri, M.E.; Zurbrügg, C.; Lindström, A.; Vinnerås, B. Faecal sludge management with the larvae of the black soldier fly (*Hermetia illucens*)—From a hygiene aspect. *Sci. Total Environ.* **2013**, *458–460*, 312–318. [[CrossRef](#)] [[PubMed](#)]
37. Papajová, I.; Pipiková, J.; Papaj, J.; Čižmár, A. Parasitic contamination of urban and rural environments in the Slovak Republic: Dog's excrements as a source. *Helminthologia* **2014**, *51*, 273–280. [[CrossRef](#)]
38. Zdybel, J.; Cencek, T.; Karamon, J.; Klapéc, T. Effectiveness of selected stages of waste water treatment in elimination of eggs of intestinal parasites. *Bull. Vet. Inst. Pulawy* **2015**, *59*, 51–57. [[CrossRef](#)]
39. Bruňanská, M. Histochemical topography of succinate dehydrogenase in the reproductive system of *Ascaris suum* females. *Helminthologia* **1989**, *26*, 43–49.
40. Stephenson, L.S.; Latham, M.C.; Ottesen, E.A. Malnutrition and parasitic helminth infections. *Parasitology* **2000**, *121*, 23–38. [[CrossRef](#)]
41. Toze, S. Reuse of effluent water—benefits and risks. *Agric. Water Manag.* **2006**, *80*, 147–159. [[CrossRef](#)]
42. Amoah, I.D.; Kumari, S.; Reddy, P.; Stendström, T.A.; Bux, F. Impact of informal settlements and wastewater treatment plants on helminth egg contamination of urban rivers and risks associated with exposure. *Environ. Monit. Assess.* **2020**, *192*, 713. [[CrossRef](#)]
43. Singh, R.P.; Agrawal, M. Potential benefits and risks of land application of sewage sludge. *Waste Manag.* **2008**, *28*, 347–358. [[CrossRef](#)]
44. Sreesai, S.; Peapueng, P.; Tippayamongkonkun, T.; Sthiannopkao, S. Assessment of a potential agricultural application of Bangkok-digested sewage sludge and finished compost products. *Waste Manag. Res.* **2013**, *31*, 925–936. [[CrossRef](#)]
45. ECDC. *National Regulation SR No. 188/2003, Laying Down the Requirements of the Application of Sew-Age Sludge and Bottom Sediments to the Soil*; ECDC: Bratislava, Slovakia, 2003. (In Slovak)
46. *STN ISO 10523; Water Quality. Determination of ph.* ISO: Geneva, Switzerland, 2010. (In Slovak)
47. *STN EN ISO 8467; Water Quality. Determination of Permanganate Index.* ISO: Geneva, Switzerland, 2000. (In Slovak)
48. *STN ISO 7150-1; Water Quality. Determination of Ammonium. Part 1: Manual Spectrometric Method.* ISO: Geneva, Switzerland, 1995. (In Slovak)
49. Mulvaney, R.L. Nitrogen—Inorganic forms. In *Methods of Soil Analysis*; Sparks, D.L., Ed.; SSSA Inc.: Madison WI, USA, 1996; pp. 1123–1184.
50. *STN EN 25663; Water Quality. Determination of Kjeldahl Nitrogen. Method after Mineralization with Selenium.* ISO: Geneva, Switzerland, 2000. (In Slovak)
51. Bremner, J.M. Nitrogen—Total. In *Methods of Soil Analysis*; Sparks, D.L., Ed.; SSSA Inc.: Madison, WI, USA, 1996; pp. 1085–1121.
52. *STN EN ISO 6878; Water Quality. Determination of Phosphorus. Ammonium Molybdate Spectrometric Method.* ISO: Geneva, Switzerland, 2005. (In Slovak)
53. Lu, R. *Soil Agricultural Chemical Analysis Methods*; China Agricultural Science and Technology Press: Beijing, China, 2000.
54. ECDC. *National Regulation SR No. 269/2010, Laying Down Requirements for Achieving Good Water Status*; ECDC: Bratislava, Slovakia, 2010. (In Slovak)
55. ECDC. *Regulation of the Slovak Government 496/2010 Amending the Regulation 354/2006 Defining the Requirements for Drinking Water Intended for Human Consumption, and for Drinking Water Quality Monitoring*; ECDC: Bratislava, Slovakia, 2010. (In Slovak)
56. *STN EN ISO 6222; Water Quality. Enumeration of Culturable Microorganisms. Colony Count by Inoculation in a Nutrient Agar Culture Medium.* ISO: Geneva, Switzerland, 2001. (In Slovak)
57. *STN EN ISO 9308-1; Water Quality. Detection and Enumeration of Coliform Organisms, Thermotolerant Coliform Organisms and Presumptive Escherichia coli. Membrane Filtration Method.* ISO: Geneva, Switzerland, 1990. (In Slovak)
58. *STN EN ISO 7899-2; Water Quality. Detection and Enumeration of Intestinal Enterococci. Part 2: Membrane Filtration Method.* ISO: Geneva, Switzerland, 2003. (In Slovak)
59. CLSI Document VET01-S2. *Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals*; Publ. Clinical and Laboratory Standards Institute: Wayne, IL, USA, 2013; pp. 1–168.
60. EUCAST. *Guidelines for Detection of Resistance Mechanisms and Specific Resistances of Clinical and/or Epidemiological Importance; Version 2.0*; EUCAST: Växjö, Sweden, 2017; pp. 1–43.
61. Gattringer, R.; Nikš, M.; Ostertág, R.; Schwarz, K.; Medvedovic, H.; Graninger, W.; Georgopoulos, A. Evaluation of MIDITECH automated colorimetric MIC reading for antimicrobial susceptibility testing. *J. Antimicrob. Chemother.* **2002**, *49*, 651–659. [[CrossRef](#)] [[PubMed](#)]
62. Cherepanov, A.A. *Methods of Laboratory Centrals of Cleaning Plants on Farms*; Kolos: Moscow, Russia, 1982. (In Russian)
63. Kazacos, K.R. Improved method for recovering ascarid and other helminth eggs from soil associated with epizootics and during survey studies. *Am. J. Vet. Res.* **1983**, *44*, 896–900. [[PubMed](#)]

64. Grant, S.B.; Saphores, J.D.; Feldman, D.L.; Hamilton, A.J.; Fletcher, T.D.; Cook, P.L.M.; Stewardson, M.; Sanders, B.F.; Levin, L.A.; Ambrose, R.F.; et al. Taking the “waste” out of “wastewater” for human water security and ecosystem sustainability. *Science* **2012**, *337*, 681–686. [[CrossRef](#)] [[PubMed](#)]
65. Bitton, G. *Wastewater Microbiology*, 3rd ed.; John Wiley & Sons: Hoboken, NJ, USA, 2005; pp. 213–217. ISBN 0-471-65071-4.
66. Lépesová, K.; Mackuľak, T.; Birošová, L. *Chapter 3: Nutrients, Wastewater and Leachate: Testing, Risks and Hazards*; Nova Science Publishers: New York, NY, USA, 2018.
67. Lépesová, K.; Olejníková, P.; Mackuľak, T.; Cverenkárová, K.; Krahulcová, M.; Birošová, L. Hospital wastewater-important source of multidrug resistant coliform bacteria with ESBL-production. *Int. J. Environ. Res. Public Health* **2020**, *17*, 7827. [[CrossRef](#)] [[PubMed](#)]
68. Feuerpfeil, J.; Lopez-Pila, J.; Schmidt, R.; Schneider, E.; Szewzyk, R. Antibiotic resistant bacteria and antibiotics in the environment. *Bundesgesundheitsblatt* **1999**, *42*, 37–50. [[CrossRef](#)]
69. Mach, P.A.; Grimes, J.D. R-plasmid transfer in a wastewater treatment plant. *Appl. Environ. Microbiol.* **1982**, *44*, 1935–1943. [[CrossRef](#)]
70. Fernandez-Astorga, A.; Muela, A.; Cisterna, R.; Iriberry, J.; Barcina, I. Biotic and abiotic factors affecting plasmid transfer in *Escherichia coli* strains. *Appl. Environ. Microbiol.* **1992**, *58*, 392–398. [[CrossRef](#)] [[PubMed](#)]
71. Reinthaler, F.F.; Posch, J.; Feierl, G.; Wüst, G.; Haas, D.; Ruckebauer, G.; Mascher, F.; Marth, E. Antibiotic resistance of *E. coli* in sewage and sludge. Affiliations expand. *Water Res.* **2003**, *37*, 1685–1690. [[CrossRef](#)]
72. Redhead, S.; Nieuwland, J.; Esteves, S.; Lee, D.H.; Kim, D.W.; Mathias, J.; Cha, C.J.; Toleman, M.; Dinsdale, R.; Guwy, A.; et al. Fate of antibiotic resistant *E. coli* and antibiotic resistance genes during full scale conventional and advanced anaerobic digestion of sewage sludge. *PLoS ONE* **2020**, *15*, e0237283. [[CrossRef](#)]
73. Martins Da Costa, P.; Vaz-Pires, P.; Bernardo, F. Antimicrobial resistance in *Escherichia coli* isolated in inflow, effluent and sludge from municipal wastewater treatment plants. *Urban Water J.* **2007**, *4*, 275–281. [[CrossRef](#)]
74. Raven, K.E.; Ludden, C.; Gouliouris, T.; Blane, B.; Naydenova, P.; Brown, N.M.; Parkhill, J.; Peacock, S.J. Genomic surveillance of *Escherichia coli* in municipal wastewater treatment plants as an indicator of clinically relevant pathogens and their resistance genes. *Microbial Genomics* **2019**, *5*, e000267. [[CrossRef](#)] [[PubMed](#)]
75. Michael, I.; Rizzo, L.; McArdeall, C.S.; Manaia, C.M.; Merlin, C.; Schwartz, T.; Dagot, C.; Fatta-Kassinos, D. Urban wastewater treatment plants as hotspots for the release of antibiotics in the environment: A review. *Water Res.* **2013**, *47*, 957–995. [[CrossRef](#)] [[PubMed](#)]
76. Mao, D.; Yu, S.; Rysz, M.; Luo, Y.; Yang, F.; Li, F.; Hou, J.; Mu, Q.; Alvarez, P.J. Prevalence and proliferation of antibiotic resistance genes in two municipal wastewater treatment plants. *Water Res.* **2015**, *85*, 458–466. [[CrossRef](#)]
77. Sidhu, J.P.S.; Toze, S.G. Human pathogens and their indicators in biosolids: A literature review. *Environ. Internat.* **2009**, *35*, 187–201. [[CrossRef](#)]
78. Dudlová, A.; Juriš, P.; Jarčuška, P.; Čisláková, L.; Papajová, I.; Krčméry, V. Epidemiological risks of endoparasitoses spread by municipal waste water. *Helminthologia* **2015**, *52*, 188–194. [[CrossRef](#)]
79. Amoah, I.D.; Reddy, P.; Seidu, R.; Stenstrom, T.A. Removal of helminth eggs by centralized and decentralized wastewater treatment plants in South Africa and Lesotho: Health implications for direct and indirect exposure to the effluents. *Environ. Sci. Pollut. Res.* **2018**, *25*, 12883–12895. [[CrossRef](#)] [[PubMed](#)]
80. Chaoua, S.; Boussaa, S.; Khadra, A.; Boumezzough, A. Efficiency of two sewage treatment systems (activated sludge and natural lagoons) for helminth egg removal in Morocco. *J. Infect. Public Health* **2018**, *11*, 197–202. [[CrossRef](#)]
81. Kansiime, F.; Van Bruggen, J.J.A. Distribution and retention of faecal coliforms in the Nakivubo wetland in Kampala, Uganda. *Water Sci. Technol.* **2001**, *44*, 199–206. [[CrossRef](#)]
82. Dai, X.; Boll, J. Settling velocity of *Cryptosporidium parvum* and *Giardia lamblia*. *Water Res.* **2006**, *40*, 1321–1325. [[CrossRef](#)]
83. Sengupta, M.E.; Thamsborg, S.M.; Andersen, T.J.; Olsen, A.; Dalsgaard, A. Sedimentation of helminth eggs in water. *Water Res.* **2011**, *45*, 4651–4660. [[CrossRef](#)]
84. Ulrich, H.; Klaus, D.; Irmgard, F.; Annette, H.; Juan, L.P.; Regine, S. Microbiological investigations for sanitary assessment of wastewater treated in constructed wetlands. *Water Res.* **2005**, *39*, 4849–4858. [[CrossRef](#)] [[PubMed](#)]
85. Morató, J.; Codony, F.; Sánchez, O.; Pérez, L.M.; García, J.; Mas, J. Key design factors affecting microbial community composition and pathogenic organism removal in horizontal subsurface flow constructed wetlands. *Sci. Total Environ.* **2014**, *481*, 81–89. [[CrossRef](#)] [[PubMed](#)]
86. Davies-Colley, R.J.; Donnison, A.M.; Speed, D.J.; Ross, C.M.; Nagels, J.W. Inactivation of faecal indicator micro-organisms in waste stabilisation ponds: Interactions of environmental factors with sunlight. *Water Res.* **1999**, *33*, 1220–1230. [[CrossRef](#)]
87. Mandi, L.; Houhoum, B.; Asmama, S.; Schwartzbrod, J. Wastewater treatment by reed beds an experimental approach. *Water Res.* **1996**, *33*, 2009–2016. [[CrossRef](#)]
88. Reinoso, R.; Torres, L.A.; Bécares, E. Efficiency of natural systems for removal of bacteria and pathogenic parasites from wastewater. *Sci. Total Environ.* **2008**, *395*, 80–86. [[CrossRef](#)]
89. Ensink, J.H.; Blumenthal, U.J.; Brooker, S. Wastewater quality and the risk of intestinal nematode infection in sewage farming families in Hyderabad, India. *Am. J. Trop. Med. Hygiene* **2008**, *79*, 561–567. [[CrossRef](#)]