

# Estradiol Reduces Susceptibility of CD4<sup>+</sup> T Cells and Macrophages to HIV-Infection

Marta Rodriguez-Garcia<sup>1\*</sup>, Nabanita Biswas<sup>1</sup>, Mickey V. Patel<sup>1</sup>, Fiona D. Barr<sup>1</sup>, Sarah G. Crist<sup>1</sup>, Christina Ochsenbauer<sup>2</sup>, John V. Fahey<sup>1</sup>, Charles R. Wira<sup>1</sup>

<sup>1</sup> Department of Physiology and Neurobiology, Geisel School of Medicine at Dartmouth, Lebanon, New Hampshire, United States of America, <sup>2</sup> Department of Medicine, University of Alabama, Birmingham, Alabama, United States of America

## Abstract

The magnitude of the HIV epidemic in women requires urgent efforts to find effective preventive methods. Even though sex hormones have been described to influence HIV infection in epidemiological studies and regulate different immune responses that may affect HIV infection, the direct role that female sex hormones play in altering the susceptibility of target cells to HIV-infection is largely unknown. Here we evaluated the direct effect of 17- $\beta$ -estradiol (E<sub>2</sub>) and ethinyl estradiol (EE) in HIV-infection of CD4<sup>+</sup> T-cells and macrophages. Purified CD4<sup>+</sup> T-cells and monocyte-derived macrophages were generated *in vitro* from peripheral blood and infected with R5 and X4 viruses. Treatment of CD4<sup>+</sup> T-cells and macrophages with E<sub>2</sub> prior to viral challenge reduced their susceptibility to HIV infection in a dose-dependent manner. Addition of E<sub>2</sub> 2 h after viral challenge however did not result in reduced infection. In contrast, EE reduced infection in macrophages to a lesser extent than E<sub>2</sub> and had no effect on CD4<sup>+</sup> T-cell infection. Reduction of HIV-infection induced by E<sub>2</sub> in CD4<sup>+</sup> T-cells was not due to CCR5 down-regulation, but was an entry-mediated mechanism since infection with VSV-G pseudotyped HIV was not modified by E<sub>2</sub>. In macrophages, despite the lack of an effect of E<sub>2</sub> on CCR5 expression, E<sub>2</sub>-treatment reduced viral entry 2 h after challenge and increased MIP-1 $\beta$  secretion. These results demonstrate the direct effect of E<sub>2</sub> on susceptibility of HIV-target cells to infection and indicate that inhibition of target cell infection involves cell-entry related mechanisms.

**Citation:** Rodriguez-Garcia M, Biswas N, Patel MV, Barr FD, Crist SG, et al. (2013) Estradiol Reduces Susceptibility of CD4<sup>+</sup> T Cells and Macrophages to HIV-Infection. PLoS ONE 8(4): e62069. doi:10.1371/journal.pone.0062069

**Editor:** Gilda Tachedjian, Burnet Institute, Australia

**Received:** November 1, 2012; **Accepted:** March 18, 2013; **Published:** April 17, 2013

**Copyright:** © 2013 Rodriguez-Garcia et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** Supported by NIH contract HHSN27220100001C, which was indirectly funded by Advanced BioScience Laboratories, Inc., and by NIH grants AI102838, AI071761 and AI013541 (CRW). CO was supported by the UAB Center for AIDS Research (P30-AI-27767) Virology and Developmental Cores. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interest exist.

\* E-mail: Marta.Rodriguez.Garcia@dartmouth.edu

## Introduction

Heterosexual transmission of HIV-1 remains a worldwide health challenge that is responsible for most HIV-1 transmissions to women (70–90%) [1]. Globally, young women are most vulnerable to HIV-1 infection, with rates of infection twice as high as young men, and as much as eight times higher in Sub-Saharan Africa, where women account for 59% of people living with HIV-1 [2,3]. Furthermore, globally HIV-1 is the leading cause of death for women of reproductive age [2].

Gender discrepancies regarding HIV-1 infection and disease progression have been repeatedly reported and are due to hormonal differences among other factors [4,5]. While plasma viral loads are lower in HIV-infected women compared to men, the rate of disease progression is greater in women [6,7]. Additionally, sex hormone fluctuations in women have been associated with both protective and adverse effects. For example, relative to the follicular and luteal phases of the menstrual cycle, decreases in plasma viral load at ovulation, when estradiol levels are high, have been previously described [8], although others did not find any effect of the menstrual cycle on HIV-RNA levels in blood [9]. In contrast, analysis of genital secretions throughout the menstrual cycle demonstrated increased HIV-1 shedding during the luteal phase, when progesterone levels are higher, in some reports [9,10] while others did not find any pattern of genital tract

shedding during the menstrual cycle [11]. More recently, significant positive associations were found between the number of days from the luteinizing hormone surge and the number of endocervical HIV-infected cells [12]. Furthermore, serum estradiol levels in women are inversely correlated with AIDS-induced dementia [13,14]. Additionally, studies conducted in macaques showed that intravaginal treatment with estriol for weeks prior to SIV vaginal challenge was able to protect. Protection was attributed to a cornification and thickening of the vaginal epithelia [15]. Despite the fact that associations between sex hormones and HIV-infection have been established, the underlying cellular and molecular mechanisms remain poorly understood.

17- $\beta$ -estradiol (E<sub>2</sub>) is the main estrogen found in blood of women and exerts its actions through binding to the estrogen receptors (ER) present in the reproductive tract tissues and in immune cells in peripheral blood, including CD4<sup>+</sup> T-cells and macrophages, the two main HIV-target cells [16,17]. Binding of E<sub>2</sub> to its receptors results in modulation of the expression of multiple genes. Studies by others and us illustrate the broad spectrum of actions of E<sub>2</sub> on immune cells and the innate and adaptive immune response, including molecules and pathways involved in anti-viral innate immune responses [18,19,20]. With the exception of studies with isolated cells from the central nervous system or cell lines [21,22,23], very little is known about the direct

effects of sex hormones on HIV-infection of immune cells. Interestingly, Asin *et al* examined the effects of sex hormones on HIV-infection and reported that different doses and combinations of estradiol and progesterone were able to regulate HIV-1 replication in peripheral blood mononuclear cells [24]. Therefore, a gap remains in our understanding of the direct effects of E<sub>2</sub> in modulating susceptibility of CD4<sup>+</sup> T-cells and macrophages to HIV-infection.

In this study we evaluated the effects of E<sub>2</sub> on HIV infection of CD4<sup>+</sup> T-cells and macrophages. We found a dose-dependent reduction of HIV-infection by E<sub>2</sub> in both cell types, through a mechanism that affected the early steps of the viral cycle but was not due to CCR5 down-regulation.

## Materials and Methods

### Study Subjects

All investigations involving human subjects were conducted according to the principles expressed in the Declaration of Helsinki and carried out with the approval from the Committee for the Protection of Human Subjects (CPHS), Dartmouth Hitchcock Medical Center, and with written informed consent obtained from volunteer healthy donors recruited at Dartmouth Hitchcock Medical Center. Blood donors were anonymous, so no information regarding age or hormonal status was available and only gender information was disclosed.

### Generation of CD4<sup>+</sup> T-cells and macrophages

Peripheral blood mononuclear cells (PBMC) were isolated by standard Ficoll density gradient centrifugation. To generate monocyte-derived macrophages, CD14<sup>+</sup> cells were positively selected with magnetic beads (Miltenyi Biotech, Auburn, CA) and incubated in ultra-low attachment 6-well plates (Corning, Corning, NY) with Xvivo 15 media (Lonza, Walkersville, MD) supplemented with 10% human AB serum (Valley Biomedical, Winchester, VA) for 4 days [25]. CD4<sup>+</sup> T-cells were purified from PBMC using magnetic negative selection (Miltenyi Biotech) and activated *in vitro* using the same media described above supplemented with Phytohemagglutinin (PHA) (2.5 µg/ml; Sigma, St Louis, MO) and IL-2 (50 U/ml) (AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH: Human rIL-2 from Dr. Maurice Gately, Hoffmann- La Roche Inc) [26] for 1 to 3 days prior to HIV-infection. Purity higher than 98% was obtained for both CD14<sup>+</sup> cells and CD4<sup>+</sup> T-cell populations after magnetic isolation (not shown).

### Hormone treatment

CD4<sup>+</sup> T-cells and macrophages were treated either with 17β-estradiol (E<sub>2</sub>; Sigma), 17α-ethinyl estradiol (EE) or Raloxifene (R<sub>x</sub>; Tocris Biosciences, Bristol, UK) as indicated in Results. CD4<sup>+</sup> T-cells were treated during activation prior to infection (1–3 days) and/or following infection (6–7 days). Macrophages were treated during differentiation period (4 days) and/or following infection (6–7 days). For all hormone experiments E<sub>2</sub>, EE or Raloxifene were dissolved in 100% ethanol for an initial concentration of 1×10<sup>-3</sup> M, evaporated to dryness and suspended in Xvivo 15 media (Invitrogen) containing 10% of charcoal dextran-stripped human AB serum (Valley Biomedical) to a concentration of 1×10<sup>-5</sup> M. Dilutions were made to achieve final working concentrations ranging from 5×10<sup>-8</sup> M to 5×10<sup>-10</sup> M for E<sub>2</sub> and EE. As a control, an equivalent amount of 100% ethanol without dissolved hormone was initially evaporated [27]. All control conditions contained evaporated ethanol as a control. For

experiments using estrogen receptor antagonist, Raloxifene was added 30 min before E<sub>2</sub> at a 100-fold excess concentration.

### Viruses

Seed stock for HIV-1<sub>BaL</sub> (R5) was obtained through the AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH, from Dr. Suzanne Gartner, Dr. Mikulas Popovic and Dr. Robert Gallo [28]. Laboratory adapted viral strain HIV-1<sub>IIIB</sub> was obtained from Dr. P. Gupta (University of Pittsburgh, PA). Viral stocks were prepared by infection of PBMC activated with PHA (2.5 µg/ml and 50 U/ml IL-2) for 6–8 days as previously described [29]. Stocks were harvested when p24 concentrations reached 100 ng/ml. Titration of viral stocks was performed using PHA-stimulated PBMC as described [29]. The replication-competent GFP-encoding infectious molecular clone (pNL<sub>ENG1i</sub>-BaL.ecto) [30] was derived from pNL<sub>ENG1i</sub>-ires [31] to express heterologous BaL *env* gene sequences in an isogenic backbone following the strategy previously described [32,33]. This reporter virus, collectively referred to as Env-IMC-GFP, expresses GFP upon infection of HIV-1 susceptible target cells. The env-defective, GFP encoding proviral plasmid pNL<sub>ENG1i</sub>-ES-ires was previously described [31], was co-transfected with a VSV-G expressing plasmid to yield infectious, non-replicating pseudovirions.

### HIV-Infection

Macrophages were infected as described previously with minor modifications [25]. Briefly, macrophages were exposed to HIV-BaL for 2 h at an MOI of 0.1 and then washed to remove residual virus. CD4<sup>+</sup> T-cells were infected with HIV-1 BaL or HIV-1 IIIB also for 2 h at an MOI of 0.1 and residual virus was washed away. Uninfected controls were incubated with medium without the virus for the same amount of time. After incubation, washed macrophage or CD4<sup>+</sup> T-cell targets were plated in round bottom ultra low attachment 96-well plates (Corning) with or without sex hormones as indicated. Cell cultures were maintained for 6–7 days, with half of the well media collected and replaced with fresh media on day 3. Levels of p24 were measured in conditioned media by p24 ELISA (Advanced Bioscience Laboratories, Rockville, MD), and intracellularly by flow cytometry (KC57-FITC; Coulter, Danvers, MA). As a control to prove that detected p24 corresponds to de novo infection and not residual viral inocula, CD4<sup>+</sup> T-cells or macrophages were incubated with Zidovudine (AZT; 10 µM) (AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH) during viral challenge (2 h) and throughout the post-infection period. No cytotoxicity was observed with this concentration of Zidovudine as measured by Trypan blue exclusion (Trypan Blue Solution, HyClone Laboratories, Inc; Logan, UT) at the end of the infection period.

### Flow cytometry

Prior to HIV infection, cells were stained for surface markers with the following antibody panels: The CD4<sup>+</sup> T-cell panel included CD4, CD3 (eBioscience, San Diego, CA), CCR5 (BD Biosciences, San Jose, CA), CXCR4 (R&D, Minneapolis, MN), CD25 (Biolegend, San Diego, CA) and HLA-DR (Miltenyi Biotech). The panel for macrophages included CD14, CD4 (eBioscience), CD163 (Biolegend), CCR5 (BD Biosciences) and DC-SIGN (R&D). After infection (6–7 days), intracellular levels of p24 were analyzed. Briefly, cells were washed, fixed and permeabilized following instructions with Cytofix/Cytoperm Plus kit (BD Biosciences) for 20 min and stained for intracellular p24 with KC57-FITC antibody (Beckman Coulter) for 30 min. Analysis was performed on BD FACSCanto flow cytometer (BD

Biosciences) using FACSDiva software and data were analyzed with the FlowJo software (Tree Star, Inc. Ashland, OR). Expression of surface markers was measured by the percentage of positive cells and the mean fluorescence intensity (MFI).

### RNA isolation and quantitative RT-PCR analysis

Total RNA was extracted from CD4<sup>+</sup> T-cells and macrophages using the RNeasy kit (QIAGEN, Valencia, CA) with on-column DNase digestion using the RNase-Free DNase set (Qiagen). RNA was quantified and cDNA was generated with the iScript cDNA synthesis kit (Bio-Rad, Hercules, CA) as described. PCR was conducted in duplicates using the 5' fluorogenic nuclease assay in real-time quantitative PCR using TaqMan chemistry on the ABI 7300 Prism real-time PCR instrument (Applied Biosystems, Carlsbad, CA). Primers and probe sets were obtained from Applied Biosystems assays-on-demand (GAPDH, CCR5, ER1, ER2). Amplification conditions and analysis were performed as described before [27]. Briefly, PCR was conducted using the following cycle parameters: 95°C, 12 min for 1 cycle (95°C, 20 s; 60°C, 1 min), for 40 cycles. Analysis was conducted using the sequence detection software supplied with the ABI 7300. The software calculates the threshold cycle (C<sub>t</sub>) for each reaction and this was used to quantify the amount of starting template in the reaction. The C<sub>t</sub> values for each set of duplicate reactions were averaged for all subsequent calculations. A difference in C<sub>t</sub> values ( $\Delta C_t$ ) was calculated for each gene by taking the mean C<sub>t</sub> of each gene of interest and subtracting the mean C<sub>t</sub> for the housekeeping gene GAPDH for each cDNA sample. Relative expression levels were calculated using the formula  $2^{-\Delta C_t}$  [34].

### ELISA

Secretion of RANTES and MIP-1 $\beta$  was measured in culture media from CD4<sup>+</sup> T-cells and macrophages 3 days after HIV-infection by using RANTES and MIP-1 $\beta$  DuoSet ELISA development system (R&D, Minneapolis, MN) according to the manufacturer's instructions.

### Statistical analysis

Data were analyzed with the GraphPad Prism 5.0 software. Non-parametric test U-Mann Whitney or Wilcoxon paired test was applied for the comparison of two groups. For the comparison of three or more groups, the non-parametric Kruskal-Wallis test followed by Dunns post-test was applied. In all cases, a two sided P value <0.05 was considered statistically significant.

## Results

### 1. E<sub>2</sub> reduces susceptibility of CD4<sup>+</sup> T-cells to HIV-infection

To examine the direct effect of E<sub>2</sub> on HIV-infection in an *in vitro* infection assay, purified CD4<sup>+</sup> T-cells (>98% purity) were activated *in vitro* in the presence or absence of E<sub>2</sub> for 3 days and infected with R5 (HIV-1 BaL) or X4 (IIIB) viral strains in parallel. Secreted and intracellular p24 was measured 7 days after infection as an indication of viral replication.

As shown in Figure 1A, when CD4<sup>+</sup>T cells were treated with E<sub>2</sub> prior to infection (pre) for 3 days, released p24 was significantly reduced 7 days after infection with HIV-1<sub>BaL</sub> (56% reduction;  $p = 0.024$ ). However, when E<sub>2</sub> was added for the entire length of incubation (prepost) or 2 h after infection (post) no differences were found compared to the control condition. This observation was confirmed by a significant reduction in the expression of intracellular p24 in CD4<sup>+</sup> T-cells pre-treated with E<sub>2</sub> before infection (Figure 1B, top row). As seen in Figure 1C, addition of

AZT, prevents de novo infection of CD4<sup>+</sup> T-cells indicating that intracellular p24 detected is not residual inocula.

In contrast, pre-incubation of CD4<sup>+</sup> T-cells with E<sub>2</sub> did not reduce infection levels with HIV-1<sub>IIIB</sub>, measured as released p24 (Figure 1A, right panel) or intracellular p24 (Figure 1B, bottom row). As seen in Figure 1A (right side), addition of E<sub>2</sub> 2 h after infection (post) slightly reduced p24 levels, but this tendency did not reach statistical significance ( $p = 0.077$ ). Since no differences were found with HIV-1<sub>IIIB</sub> the following experiments focused on HIV-1<sub>BaL</sub>.

The experiments described above were performed at a high dose of E<sub>2</sub> ( $5 \times 10^{-8}$  M), which under physiological conditions, is reached during the periovulatory phase. Since this estradiol peak lasts between 1 to 3 days, we wanted to know if exposure of CD4<sup>+</sup> T-cells to a high dose of E<sub>2</sub> for 1 day would be sufficient to suppress HIV-infection. Figure 1D shows that 1 day pretreatment with E<sub>2</sub> significantly reduces both released and intracellular p24 levels relative to controls. Overall the same effect was observed with 3 days or 1 day of pre-treatment with E<sub>2</sub> (Figures 1A and 1D respectively). Equivalent to the results obtained after activation and E<sub>2</sub> treatment for 3 days (Figure 1A), only pre-incubation with E<sub>2</sub> for 1 day had a suppressive effect on HIV infection, and no inhibition was measured in released or intracellular p24 (Figure 1D, left and right panels) when E<sub>2</sub> was added after infection.


Since these experiments were conducted with cells from both female and male donors, we analyzed if any differences could be found between them. As shown in Figure 1E, p24 levels in the infected controls from female donors were lower than male donors, with median p24 values of 38,085 versus 71,541 respectively ( $p = 0.03$ ). To determine whether E<sub>2</sub> inhibited HIV infection of CD4<sup>+</sup> T-cells from men and women, cells were pretreated with E<sub>2</sub> for 1 day prior to infection. As seen in Figure 1E, treatment of CD4<sup>+</sup> T-cells from women significantly reduced HIV-infection (54.9% reduction;  $p = 0.03$ ), but had no effect on CD4<sup>+</sup> T-cells from men.

We investigated whether the reduction in HIV-susceptibility induced by E<sub>2</sub> could be due to differences in activation. CD4<sup>+</sup> T-cells treated or not with E<sub>2</sub> showed similar expression of CD25 and HLA-DR before HIV challenge (data not shown).

### 2. E<sub>2</sub> reduces susceptibility of macrophages to HIV-infection

Recognizing that CD4<sup>+</sup> T-cells and macrophages are the most likely targets for HIV infection, we next focused on the effect of E<sub>2</sub> on macrophage infection. Monocyte-derived macrophages were differentiated *in vitro* in the presence of E<sub>2</sub> for 4 days and infected with HIV-1<sub>BaL</sub>. As shown in Figure 2A, similar to our results with CD4<sup>+</sup> T-cells, viral replication was significantly reduced in macrophages differentiated in the presence of E<sub>2</sub> ( $p < 0.0003$ ). This reduction in susceptibility to HIV-infection was induced by differentiating the macrophages in the presence of E<sub>2</sub> before infection (pre; 69% reduction) and maintained when E<sub>2</sub> was added back to the culture 2 h after infection (prepost; 73% reduction). In contrast, E<sub>2</sub> had no effect when added immediately after infection (post) (Figure 2A). Intracellular p24 analysis confirmed a significant reduction in the percent of p24 positive cells when macrophages were differentiated in the presence of E<sub>2</sub> (Figure 2B and 2C;  $p < 0.05$ ). Figure 2D demonstrates that intracellular p24 values correspond to de novo infection, since macrophages infected in the presence of AZT had intracellular p24 values equal to uninfected controls.

To determine whether HIV infection of macrophages was influenced by gender, blood derived macrophages from women and



**Figure 1. Effect of E<sub>2</sub> on HIV-infection of CD4<sup>+</sup> T-cells.** A) Released p24 levels in the culture media after 7 days of infection when cells were pre-treated with E<sub>2</sub> for 3 days (pre E<sub>2</sub>), treated with E<sub>2</sub> before and after infection (prepost E<sub>2</sub>) or only after infection (post E<sub>2</sub>). Bars represent mean ± SEM from 7 independent experiments with different donors. \*P<0.05. B) Intracellular p24 levels after 7 days of infection, representative of n=7. C) Intracellular p24 levels after 7 days of infection in the presence of Zidovudine (AZT). D) Released p24 levels in the culture media (left panel) and intracellular p24 (right panel) after 7 days of infection when cells were pre-treated with E<sub>2</sub> for 1 day (pre E<sub>2</sub>), treated with E<sub>2</sub> before and after infection (prepost E<sub>2</sub>) or only after infection (post E<sub>2</sub>). Bars represent mean ± SEM from 6 independent experiments with different donors. \*P<0.05. E) Released p24 in the culture media after 7 days of infection in CD4<sup>+</sup> T-cells from women (black bars; N=6) or men (grey bars; N=5). \*P<0.05. doi:10.1371/journal.pone.0062069.g001

men were analyzed for HIV infection and responsiveness to E<sub>2</sub>. In contrast to CD4<sup>+</sup> T-cells (Figure 1E), irrespective of donor origin, no differences in HIV infection were observed. Interestingly, infection of macrophages derived from female and male donors was significantly reduced when cells were pretreated with E<sub>2</sub>.

When Figures 1 and 2 are compared, E<sub>2</sub> appears to be more effective in reducing susceptibility to HIV-infection in macrophages than CD4<sup>+</sup> T-cells. Further, the suppressive effect was maintained when E<sub>2</sub> was present after infection. This maintenance of suppression represents a difference with respect to the effect observed in CD4<sup>+</sup> T-cells in that the inhibitory effect induced by pre-treatment with E<sub>2</sub> is lost when E<sub>2</sub> is present after infection (Figure 1).


### 3. E<sub>2</sub> and ethinyl estradiol (EE) have different effects on CD4<sup>+</sup> T-cell and macrophage susceptibility to HIV-infection

Recognizing that ethinyl estradiol is the main estrogenic component in most oral contraceptives and that contraceptives

may be a risk factor for HIV infection [10,35], we investigated if EE would have the same effect as E<sub>2</sub> in preventing HIV-infection of target cells.

Following the same experimental design described above (Figure 1), CD4<sup>+</sup> T-cells were activated in the presence of EE and infected with R5 and X4 viral strains. As seen in Figure 3A, irrespective of the time (pre) and length (prepost/post) of hormone treatment, and unlike E<sub>2</sub> which inhibited HIV infection, EE had no effect on CD4<sup>+</sup> T-cell infection with HIV-1<sub>BaL</sub>. Similar to the lack of inhibition with E<sub>2</sub>, we observed a pattern of reduced HIV-1<sub>IIIb</sub> viral replication when EE was added 2 h after infection, but no statistical significance (Figure 3A, right panel).

In contrast, EE was able to decrease macrophage susceptibility to HIV-infection. As shown in Figure 3B, when macrophages were differentiated in the presence of EE, released p24 was significantly reduced 7 days after infection (p<0.005). When CD4<sup>+</sup> T-cells and macrophages from the same donors were analyzed in parallel to compare the effects of E<sub>2</sub> and EE, intracellular p24 staining



**Figure 2. Effect of E<sub>2</sub> on macrophage HIV-infection.** A) Released p24 levels in the culture media after 7 days of infection when cells were pre-treated with E<sub>2</sub> (pre E<sub>2</sub>), treated with E<sub>2</sub> before and after infection (prepost E<sub>2</sub>) or only after infection (post E<sub>2</sub>). Bars represent mean ± SEM from 5 independent experiments with different donors. \*\*P<0.01;\*\*\*P<0.001. B) Representative contour plot of intracellular p24 levels after 7 days of infection and C) Fold change respect infected control of intracellular p24 from 4 different donors. Mean ± SEM is represented. \*p<0.05. D) Intracellular p24 levels after 7 days of infection in the presence of Zidovudine (AZT). E) Released p24 in the culture media after 7 days of infection in CD4<sup>+</sup> T-cells from women (black bars; N=3) or men (grey bars; N=3). \*P<0.05. doi:10.1371/journal.pone.0062069.g002

confirmed the lack of effect on CD4<sup>+</sup> T-cells but a decrease in the % of infected macrophages for both E<sub>2</sub> and EE pre-treatments (Figure 3C). Pre-treatment with E<sub>2</sub> however, was consistently 10–20% more effective than EE in suppressing viral replication (Figure 3D).

#### 4. E<sub>2</sub> suppresses HIV-infection in a dose-dependent manner

To determine if the observed suppression of viral replication was dose-dependent, serial 10-fold concentrations of E<sub>2</sub> were tested. CD4<sup>+</sup> T-cells were incubated with the different doses of E<sub>2</sub> for 24 h and then infected with HIV-1<sub>BaL</sub>. As shown in Figure 4A, E<sub>2</sub> suppressed HIV-infection of CD4<sup>+</sup> T-cells in a dose-dependent fashion, with 10<sup>-8</sup> M being the tested concentration that most effectively suppressed viral replication.


To evaluate the effect of E<sub>2</sub> on macrophages, monocytes were differentiated into macrophages in the presence of serial 10-fold concentrations of E<sub>2</sub> for 4 days and their effect on viral replication evaluated 7 days post infection. Similar to the effects observed in CD4<sup>+</sup> T-cells, viral replication in macrophages was inhibited by E<sub>2</sub> in a dose-dependent manner, with 10<sup>-8</sup> M being the most effective concentration (Figure 4B, black bars). As a part of these studies, we also evaluated the effect of EE on macrophages and found that EE at 10<sup>-8</sup> M inhibited viral replication. Consistent

with our previous results, the levels of inhibition were slightly lower than those obtained with E<sub>2</sub>.

We then asked if the reduction in HIV-infection induced by E<sub>2</sub> was dependent on the dose of viral inoculum used. To investigate this, we used replication-competent HIV reporter viruses that encodes BaL *env* and expresses GFP upon infection. CD4<sup>+</sup> T-cells were treated with E<sub>2</sub> for 24 h and then infected with two different concentrations of virus. As seen in Figure 4C (upper panel), infection with a MOI of 0.1 corroborated our earlier results, showing approximately a 50% reduction (5.29 to 2.83%) in number of GFP+ cells following pre-treatment with E<sub>2</sub>, but not when E<sub>2</sub> was present both pre- and post-infection (Figure 4C, top row). In contrast, we found that at higher viral inocula (MOI 1) (Figure 4C, lower panel), the inhibitory effect of E<sub>2</sub> was abrogated. These findings suggest that E<sub>2</sub> induces a saturable mechanism of HIV-restriction.

#### 5. E<sub>2</sub> restriction of viral replication is not due to CCR5 down-regulation

To investigate the possible mechanisms involved in E<sub>2</sub> suppression of viral replication, we first examined the expression of CCR5, the main coreceptor for R5 viral strains. CD4<sup>+</sup> T-cells were activated in the presence of E<sub>2</sub> and CCR5 expression was assayed by RNA and flow cytometry immediately prior to HIV-



**Figure 3. Effect of EE on HIV-infection of CD4<sup>+</sup> T-cells and macrophages.** A) Released p24 levels in the culture media after 7 days of infection when CD4<sup>+</sup> T-cells were pre-treated with EE (pre EE), treated with EE before and after infection (prepost EE) or only after infection (post EE). Bars represent mean  $\pm$  SEM from 4 independent experiments with different donors. B) Released p24 levels in the culture media after 7 days of infection when macrophages were pre-treated with EE (pre EE), treated with EE before and after infection (prepost EE) or only after infection (post EE). Bars represent mean  $\pm$  SEM from 4 independent experiments with different donors. \*\* $P < 0.01$ . C) Comparison between CD4<sup>+</sup> T-cells and macrophages from the same donors pre-treated with E<sub>2</sub> and EE. Dot plots represent percent of p24<sup>+</sup> cells. D) Percent of suppression of HIV-infection respect to infected control in macrophages from the same donors pre-treated with E<sub>2</sub> (black dots) or EE (grey dots) in parallel. Each dot represents one individual.


doi:10.1371/journal.pone.0062069.g003

infection. No differences were observed in CD4<sup>+</sup> T-cell CCR5 gene expression after E<sub>2</sub> treatment, relative to controls (Figure 5A). When cells were analyzed for % positive cells and MFI by flow cytometry, surface expression of CCR5 in response to E<sub>2</sub> was either decreased or increased depending on the donor (Figure 5B) and showed no correlation with the levels of viral suppression. Surface expression of CD4 was also measured, but no differences were found between controls and E<sub>2</sub>-treated cells (not shown).

To measure viral entry at 2 h post infection [36], CD4<sup>+</sup> T-cells pre-treated with E<sub>2</sub> for 24 h were incubated with HIV-1<sub>BaL</sub> for 2 h at 37°C, after which free virus was removed by repeated washes. Cells were then trypsinized, washed and lysed prior to measuring p24 in the cell lysates by ELISA. As a control, cells were exposed to the virus for 2 h at 4°C to account for residual extracellular binding; this background was subtracted from the p24 values measured at 37°C. Half of the cells were kept in culture and HIV-infection was followed for 7 days to confirm the inhibitory activity of E<sub>2</sub>. As shown in Figure 5C, after 2 h of exposure to the virus, intracellular levels of p24 were not affected by E<sub>2</sub> pre-treatment, but determination of infection levels 7 days after viral exposure confirmed suppression of HIV-infection by E<sub>2</sub> (Figure 5C).

Macrophages differentiated in the presence of E<sub>2</sub> or EE were also analyzed for the expression of CCR5 at the time of infection. As seen in Figure 5A, neither E<sub>2</sub> nor EE had any effect on CCR5 gene expression relative to controls. Surface expression of CCR5 in macrophages also showed a donor variable scattered pattern after treatment with E<sub>2</sub> that did not correlate with viral suppression (Figure 5B). Likewise, no differences were found regarding CD4 or DC-SIGN surface expression (not shown). Viral entry was then assessed in the same manner described above for CD4<sup>+</sup> T-cells. In contrast to CD4<sup>+</sup> T-cells, differentiation of macrophages in the presence of E<sub>2</sub> resulted in a significant reduction of intracellular p24 after viral exposure for 2 h (Figure 5D). Interestingly, EE had no effect in reducing viral entry 2 h after viral challenge, but both E<sub>2</sub> and EE reduced infection levels 7 days after (Figure 5D).

To better define the role of E<sub>2</sub> in affecting viral entry, we pre-treated CD4<sup>+</sup> T-cells and macrophages with E<sub>2</sub> prior to infection with a single cycle, VSV-G pseudotyped virus, which enters the cells by endocytosis and bypasses receptor/coreceptor attachment and fusion [37]. As shown in Figure 5E, no differences in secreted p24 levels were found between the control cells and cells pre-treated with E<sub>2</sub>, strongly



**Figure 4. Dose response effect of E<sub>2</sub> on HIV-infection.** A) Released p24 levels in the culture media after 7 days of infection when CD4<sup>+</sup> T-cells where pre-treated with indicated doses of E<sub>2</sub>. Bars represent mean ± SEM from triplicates. Representative of 3 independent experiments with different donors. \*P<0.05 \*\*P<0.01. B) Released p24 levels in the culture media after 7 days of infection when macrophages where pre-treated with indicated doses of E<sub>2</sub> (black bars) or EE (grey bars). Bars represent mean ± SEM from triplicates. Representative of 3 independent experiments with different donors. \*P<0.05 \*\*P<0.01. C) Percent of GFP positive cells after infection with HIV-GFP at MOI 1 or MOI 0.1 in CD4<sup>+</sup> T-cells pre-treated with E<sub>2</sub> (pre E<sub>2</sub>) or treated before and after infection (prepost E<sub>2</sub>). doi:10.1371/journal.pone.0062069.g004

suggesting that E<sub>2</sub> inhibits infection at the steps of viral attachment or fusion and not during reverse transcription.

## 6. Effect of E<sub>2</sub> on CCR5 ligand secretion

Since our results suggest that E<sub>2</sub> is affecting viral entry, without down-regulating CD4 and CCR5 expression, we explored the possibility that E<sub>2</sub> treatment alters ligand secretion of RANTES and MIP-1β, known to block HIV-infection through CCR5 binding. CD4<sup>+</sup> T-cells and macrophages were treated with E<sub>2</sub> prior to infection with HIV<sub>BaL</sub> after which secretion of RANTES and MIP-1β was analyzed. As shown in Figure 6A, E<sub>2</sub> treatment had no effect on RANTES secretion by CD4<sup>+</sup> T-cells or macrophages. In contrast, secretion of MIP-1β was significantly increased in macrophages (Figure 6B), but unaffected by E<sub>2</sub> treatment of CD4<sup>+</sup> T-cells (Figure 6C, normalized values). These findings suggest that E<sub>2</sub> may reduce viral entry in macrophages by increasing MIP-1β secretion, which in turn interferes with HIV binding to CCR5 in macrophages. Whether other ligands are involved in CD4<sup>+</sup> T-cell suppression on HIV infection remains to be determined.

## 7. Inhibition of HIV-infection by E<sub>2</sub> is mediated through the estrogen receptor (ER)

To further examine the mechanisms involved in the reduction of HIV-infection by E<sub>2</sub>, we investigated the role of the estrogen

receptor. First we analyzed expression levels of ERα and ERβ in CD4<sup>+</sup> T-cells and macrophages (Figure 7A). Both CD4<sup>+</sup> T-cells and macrophages expressed ERα, with macrophages expression values approximately 10-fold higher than CD4<sup>+</sup> T-cells (median expression of 0.001 vs 0.00013 respectively; p = 0.005). In contrast, ERβ expression was low compared to ERα in CD4<sup>+</sup> T-cells, and undetectable in macrophages (Figure 7A). To establish that the effects of E<sub>2</sub> are mediated through ER, CD4<sup>+</sup> T-cells and macrophages were treated with E<sub>2</sub> in the presence or not of Raloxifene, a known preferential ERα antagonist [38]. As seen in Figure 7B, when cells were pretreated with Raloxifene for 30 min prior to the addition of E<sub>2</sub>, Raloxifene reversed the inhibitory effects of E<sub>2</sub> on p24 secretion by CD4<sup>+</sup> T-cells and macrophages. Importantly, Raloxifene alone had no effect on HIV-infection.

## Discussion


In the present study we demonstrate a reduction in HIV-susceptibility of CD4<sup>+</sup> T-cells and macrophages induced by E<sub>2</sub>. To the best of our knowledge the data presented here are the first to show that E<sub>2</sub> is involved by a direct mechanism in reducing the susceptibility to HIV-infection of CD4<sup>+</sup> T-cells and macrophages, the main target cells of HIV. We found that restriction of HIV-replication in target cells is not due to CCR5 down-regulation and appears to be mediated through a saturable mechanism of restriction.

Gender differences regarding HIV-infection are now well established. At the time of seroconversion women present lower viral titers than men and, compared to men at similar stages of the disease, women maintain lower viral loads [4,5]. Despite this evidence and the known interactions between the immune system and sex hormones, little is known about the direct effect of E<sub>2</sub> in modulating susceptibility of HIV-target cells to infection. A paucity of reports previously investigated the direct effects of E<sub>2</sub> on HIV infection, but most focused on cells that are not the main HIV-target cells [23,39,40] or used cell line models that were already infected [21]. Only one previous report approached the study of PBMC [24], leaving open the question if the contribution of E<sub>2</sub> was on CD4<sup>+</sup> T-cells, macrophages or other cell subsets such as CD8<sup>+</sup>T cells or NK cells.

In our study we observed suppression of HIV-infection in CD4<sup>+</sup> T-cells when they were pre-treated with E<sub>2</sub> prior but not after infection. This would be in agreement with a previous study in which, when E<sub>2</sub> was present throughout the experiment, no hormonal effect was observed with HIV-producing ACH-2 lymphocytes [21].

Our results also demonstrate that pre-treatment with E<sub>2</sub> was able to reduce susceptibility of macrophages to HIV infection with a more pronounced effect than the one observed for CD4<sup>+</sup> T-cells. Macrophages play a major role in HIV pathogenesis by contributing to HIV dissemination and establishment of viral reservoirs. As migratory cells, macrophages can reach enclosed environments such as the central nervous system that T cells cannot reach [41]. Our *in vitro* studies support the hypothesis that exposure of monocytes to high levels of E<sub>2</sub> reached in peripheral blood during ovulation would render them less susceptible to HIV infection when they migrate to the tissues and differentiate into macrophages.

An unexpected finding in our study was that whereas pre-treatment with E<sub>2</sub> protected CD4<sup>+</sup> T-cells from HIV infection, prolonged exposure post infection, had no effect. Others have shown that the long terminal repeat of HIV contains a steroid hormone-responsive element [42] through which E<sub>2</sub> enhances viral replication [22]. This could explain why addition of E<sub>2</sub> before




**Figure 5. Effect of E<sub>2</sub> on CCR5 expression and viral entry.** A) CCR5 expression by RT-PCR. Bars represent mean  $\pm$  SEM from 4 different donors. B) Surface expression of CCR5 after treatment with E<sub>2</sub> or EE compared to control. C) p24 detection in cell lysates from CD4<sup>+</sup> T-cells 2 h after infection (N=4) and intracellular p24 expression 7 days after infection. D) p24 detection in cell lysates from macrophages 2 h after infection (N=3) and intracellular p24 levels 7 days after infection. E) Released p24 in the culture media 3 days after infection with VSV-G pseudotyped HIV. Bars represent mean  $\pm$  SEM from 4 independent experiments with CD4<sup>+</sup> T-cells and 3 with macrophages from different donors. doi:10.1371/journal.pone.0062069.g005

and after infection did not have the same effect as pre-treatment alone in CD4<sup>+</sup> T-cells. Alternatively, others have shown cell surface expression of ER $\alpha$  in T cells, corresponding to the ER $\alpha$ 46 isoform, which in response to E<sub>2</sub> induces rapid phosphorylation of ERK and proliferation of T cells [43]. The plasma membrane-associated form of ER has the ability to rapidly signal in response to E<sub>2</sub> and can ultimately regulate transcriptional activation [44]. These findings suggest that E<sub>2</sub> added again after infection, signaling through membrane-associated ER, could mask the inhibitory effect of E<sub>2</sub> pre-treatment on HIV infection. In macrophages, however, pre-treatment as well as pre- and post-treatment inhibited HIV infection while post-infection treatment

with E<sub>2</sub> had no effect. Just why these cells differ in their responsiveness to E<sub>2</sub> remains uncertain. What is likely is that an explanation resides in the unique characteristics of macrophages given that both CD4<sup>+</sup> T-cells and macrophages received the same virus. Our results indicate that CD4<sup>+</sup> T-cells and macrophages display different ER expression profiles. Macrophages expressed significantly higher levels of ER $\alpha$  but lacked ER $\beta$  expression, while CD4<sup>+</sup> T-cells expressed both forms of ER. Taking into consideration that inhibition of HIV-infection in our system is most likely mediated through ER $\alpha$ , as also indicated by others [45], it is possible that the differences in ER expression between cell types may account for the observed differences. Additionally, previous






**Figure 6. Effect of E<sub>2</sub> on CCR5 ligand secretion.** A) Levels of RANTES detected in the culture media from CD4<sup>+</sup> T-cells (N=6) and macrophages (N=6) 3 days after infection with HIV-BaL. B) Levels of MIP-1β detected in culture media from CD4<sup>+</sup> T-cells (N=8) and macrophages (N=6) 3 days after infection with HIV-BaL. Each dot represents a different donor. C) Box plot of normalized levels of MIP-1β in E<sub>2</sub>-treated CD4<sup>+</sup> T-cells (white box) and macrophages (grey box) as compared to the control. Whiskers represent maximum and minimum values. \*P<0.05. doi:10.1371/journal.pone.0062069.g006

studies have shown that macrophages [46] and T cells [43] express different isoforms of ERα, ERα66 and ERα46, and that their relative expression is modified by E<sub>2</sub> [46,47]. Since the ERα46 isoform functions as a repressor of ERα66, differences in the expression of these isoforms will influence gene regulation [46]. The relative contribution of the different ERα isoforms to the reduction in HIV-infection induced by E<sub>2</sub> remains to be elucidated. While E<sub>2</sub> treatment had no effect on the internalization of virions two hours after viral challenge in CD4<sup>+</sup> T-cells, macrophages differentiated in the presence of E<sub>2</sub> had reduced viral entry at two hours. These findings could account for the E<sub>2</sub> sustained inhibition of HIV infection in macrophages compared to CD4<sup>+</sup> T-cells. As a part of these studies we investigated the early steps in HIV-infection by using a VSV-G pseudotyped virus, which does not require coreceptor-mediated attachment and fusion to enter the cells [37]. Since the viral cycle of VSV-G pseudotyped virus is the same as replicant competent HIV once the virus is inside the cell, the lack of differences in infection levels that we observed between control and E<sub>2</sub>-treated cells infected with VSV-G pseudotyped HIV strongly suggests that E<sub>2</sub> reduces HIV-infection by affecting some step before reverse transcription. Since E<sub>2</sub> had no effect on CD4 or CCR5 expression, these findings suggest that E<sub>2</sub> may inhibit HIV infection by stimulating CCR5-ligand secretion, which on binding to the receptor CCR5, interferes with HIV binding to this receptor. We found a significant increase in MIP-1β secretion by E<sub>2</sub>-treated macrophages, but not in CD4<sup>+</sup> T-cells, which could account for the

differences observed between cell types. Since MIP-1β, is one of many ligands that bind CCR5, further studies are needed to more fully understand the complex mechanisms through which E<sub>2</sub> functions to alter susceptibility of HIV-target cells to infection.

An interesting finding in our study was the differences in HIV-infection and responsiveness to E<sub>2</sub> found in immune cells from the blood of women and men. In the absence of E<sub>2</sub>, CD4<sup>+</sup> T-cells from women had lower levels of infection than that found in men. Additionally, CD4<sup>+</sup> T-cells from women were more responsive to E<sub>2</sub> than cells from men. Differential immune responsiveness to E<sub>2</sub> treatment by CD4<sup>+</sup> T-cells from women and men have been recently described by Moulton *et al* [48]. In their study, E<sub>2</sub> increased CREMα mRNA expression and down-regulated IL-2 secretion in T cells from women more frequently than from men. In contrast to CD4<sup>+</sup> T-cells, we found no significant differences between monocyte-derived macrophages from women and men. Further studies are needed to more fully define the differences in susceptibility of HIV-target cells from men and women.

Previous studies by us led to the hypothesis of a “window of vulnerability” in the menstrual cycle during which women are more susceptible to HIV infection [49]. This hypothesis was based on studies of the effects of E<sub>2</sub> and progesterone on immune responses in the female reproductive tract both *in vitro* and during the menstrual cycle. The present study extends this hypothesis by demonstrating that E<sub>2</sub> directly affects susceptibility of the target cells to HIV infection in a way that confers protection at a time when many aspects of the adaptive and innate immune systems are dampened, to optimize conditions for successful fertilization, implantation and pregnancy. Our dose response studies indicate that inhibition in CD4<sup>+</sup> T-cells and macrophages is maximal at  $5 \times 10^{-8}$  M, the concentration known to saturate estrogen receptors in target cells [50]. This concentration is comparable to E<sub>2</sub> levels at ovulation, and during part of the secretory phase of the menstrual cycle. Under these conditions, we hypothesize that while uterine CTL activity as well as innate protection in the cervix and vagina is suppressed [49,51,52], HIV-target cells would have reduced susceptibility to infection. Further studies are needed to determine whether HIV-target cell susceptibility to HIV infection varies with the menstrual cycle in the female reproductive tract (FRT). Our results are relevant to understand HIV pathogenesis, not only as a surrogate *in vitro* model for the cells present in the mucosa at the time of exposure, but also given the fact that CD4<sup>+</sup> T-cell and monocyte migration from peripheral blood into the FRT is hormone and inflammation responsive. Importantly, our results suggest a protective role for E<sub>2</sub> throughout the FRT at the time when fecundation is likely to occur and HIV, present in semen, is most likely to be present.



**Figure 7. Role of estrogen receptor in E<sub>2</sub>-mediated suppression of HIV-infection.** A) Expression of ERα (black bars) and ERβ (grey bars) detected by real time RT-PCR in CD4<sup>+</sup> T-cells (N=3) and macrophages (N=4). Bars represent mean ± SEM from 3 and 4 different donors for CD4<sup>+</sup> T-cells and macrophages respectively. ND: non-detected. B) Released p24 detected by ELISA in the culture media of CD4<sup>+</sup> T-cells and macrophages after 7 days of infection. Bars represent mean ± SEM from 4 replicates. Representative of 3 independent experiments. Rx: Raloxifene. \*P<0.05. doi:10.1371/journal.pone.0062069.g007

Our study indicates that ethinyl estradiol (EE), a known synthetic estrogen used in many chemical contraceptive formulations (oral, ring, etc) exerts effects on HIV-target cells that are different from those seen with E<sub>2</sub>, the naturally occurring estrogen. While many studies have investigated the interactions between EE and antiretroviral drugs in HIV-infected women, to the best of our knowledge, ours is the first study to evaluate the direct effect of EE in influencing susceptibility of target cells to HIV-infection. In the present study, while E<sub>2</sub> had a clear effect in reducing susceptibility of CD4<sup>+</sup> T-cells to HIV-infection, no significant effect was observed with EE at the same concentration. Additionally, even though EE reduced susceptibility of macrophages to HIV-infection, it was consistently less effective than E<sub>2</sub>. Lastly, our studies indicate that EE, unlike E<sub>2</sub>, played no role in reducing entry of the virus in macrophages within 2 h. These results demonstrate that EE is less effective than E<sub>2</sub> in reducing susceptibility of HIV-target cells to infection. EE has a greater affinity for ER than E<sub>2</sub> [50], but the mechanism(s) responsible for the differential effects between E<sub>2</sub> and EE remains to be determined. As oral contraceptives are a part of multifaceted

preventive strategies, these findings highlight the importance of a careful evaluation of the impact of hormonal contraception in HIV-acquisition.

In conclusion, our data indicate that E<sub>2</sub> acts directly on CD4<sup>+</sup> T-cells and macrophages to inhibit HIV-infection. These results should contribute to our understanding of HIV-acquisition and highlight the importance of considering the influence of menstrual cycle and hormonal contraceptives when planning new preventive strategies in women.

## Acknowledgments

We thank Richard Rossoll and Zheng Shen for technical assistance.

## Author Contributions

Conceived and designed the experiments: MRG CRW. Performed the experiments: MRG NB MVP FDB SGC. Analyzed the data: MRG NB MVP FDB SGC CO JVF CRW. Contributed reagents/materials/analysis tools: CO. Wrote the paper: MRG CRW.

## References

- NIAID (2010) HIV infection in Women.
- UNAIDS (2010) Women, girls, gender equality and HIV.
- WHO (2011) Progress report 2011: Global HIV/AIDS response.
- Gandhi M, Bacchetti P, Miotti P, Quinn TC, Veronese F, et al. (2002) Does patient sex affect human immunodeficiency virus levels? *Clin Infect Dis* 35: 313–322.
- Gilad J, Walfisch A, Borer A, Schlaeffer F (2003) Gender differences and sex-specific manifestations associated with human immunodeficiency virus infection in women. *Eur J Obstet Gynecol Reprod Biol* 109: 199–205.
- Anastos K, Gange SJ, Lau B, Weiser B, Detels R, et al. (2000) Association of race and gender with HIV-1 RNA levels and immunologic progression. *J Acquir Immune Defic Syndr* 24: 218–226.
- Sterling TR, Lyles CM, Vlahov D, Astemborski J, Margolick JB, et al. (1999) Sex differences in longitudinal human immunodeficiency virus type 1 RNA levels among seroconverters. *J Infect Dis* 180: 666–672.
- Cu-Uvin S (2005) Effect of the menstrual cycle on virological parameters. *J Acquir Immune Defic Syndr* 38 Suppl 1: S33–34.
- Reichelderfer PS, Coombs RW, Wright DJ, Cohn J, Burns DN, et al. (2000) Effect of menstrual cycle on HIV-1 levels in the peripheral blood and genital tract. WHS 001 Study Team. *AIDS* 14: 2101–2107.
- Baeten JM, Lavreys L, Overbaugh J (2007) The influence of hormonal contraceptive use on HIV-1 transmission and disease progression. *Clin Infect Dis* 45: 360–369.
- Mostad SB, Jackson S, Overbaugh J, Reilly M, Chohan B, et al. (1998) Cervical and vaginal shedding of human immunodeficiency virus type 1-infected cells throughout the menstrual cycle. *J Infect Dis* 178: 983–991.
- Benki S, Mostad SB, Richardson BA, Mandaliya K, Kreiss JK, et al. (2008) Increased levels of HIV-1-infected cells in endocervical secretions after the luteinizing hormone surge. *J Acquir Immune Defic Syndr* 47: 529–534.
- Clark RA, Bessinger R (1997) Clinical manifestations and predictors of survival in older women infected with HIV. *J Acquir Immune Defic Syndr Hum Retrovirol* 15: 341–345.
- Grinspoon S, Corcoran C, Miller K, Biller BM, Askari H, et al. (1997) Body composition and endocrine function in women with acquired immunodeficiency syndrome wasting. *J Clin Endocrinol Metab* 82: 1332–1337.
- Smith SM, Mefford M, Sadora D, Klase Z, Singh M, et al. (2004) Topical estrogen protects against SIV vaginal transmission without evidence of systemic effect. *AIDS* 18: 1637–1643.
- McKinnon LR, Kaul R (2012) Quality and quantity: mucosal CD4<sup>+</sup> T cells and HIV susceptibility. *Curr Opin HIV AIDS* 7: 195–202.
- Cobos-Jimenez V, Boiman T, Hamann J, Kootstra NA (2011) Macrophages and HIV-1. *Curr Opin HIV AIDS* 6: 385–390.
- Escribese MM, Rodriguez-Garcia M, Sperling R, Engel SM, Gallart T, et al. (2011) Alpha-defensins 1–3 release by dendritic cells is reduced by estrogen. *Reprod Biol Endocrinol* 9: 118.
- Hickey DK, Patel MV, Fahey JV, Wira CR (2011) Innate and adaptive immunity at mucosal surfaces of the female reproductive tract: stratification and integration of immune protection against the transmission of sexually transmitted infections. *J Reprod Immunol*.
- Escribese MM, Kraus T, Rhee E, Fernandez-Sesma A, Lopez CB, et al. (2008) Estrogen inhibits dendritic cell maturation to RNA viruses. *Blood* 112: 4574–4584.
- Bourinbaier AS, Nagorny R, Tan X (1992) Pregnancy hormones, estrogen and progesterone, prevent HIV-1 synthesis in monocytes but not in lymphocytes. *FEBS Lett* 302: 206–208.
- Katagiri D, Hayashi H, Victoriano AF, Okamoto T, Onozaki K (2006) Estrogen stimulates transcription of human immunodeficiency virus type 1 (HIV-1). *Int Immunopharmacol* 6: 170–181.
- Wilson ME, Allred KF, Bisotti AJ, Bruce-Keller A, Chuahan A, et al. (2006) Estradiol negatively regulates HIV-LTR promoter activity in glial cells. *AIDS Res Hum Retroviruses* 22: 350–356.
- Asin SN, Heimberg AM, Eszterhas SK, Rollenhagen C, Howell AL (2008) Estradiol and progesterone regulate HIV type 1 replication in peripheral blood cells. *AIDS Res Hum Retroviruses* 24: 701–716.
- Rodriguez-Garcia M, Porchis F, de Jong OG, Levi K, Diefenbach TJ, et al. (2011) Expression of PD-L1 and PD-L2 on human macrophages is up-regulated by HIV-1 and differentially modulated by IL-10. *J Leukoc Biol* 89: 507–515.
- Lahm HW, Stein S (1985) Characterization of recombinant human interleukin-2 with micromethods. *J Chromatogr* 326: 357–361.
- Patel MV, Ghosh M, Fahey JV, Wira CR (2012) Uterine epithelial cells specifically induce interferon-stimulated genes in response to polyinosinic-polycytidylic acid independently of estradiol. *PLoS One* 7: e35654.
- Gartner S, Markovits P, Markovits DM, Kaplan MH, Gallo RC, et al. (1986) The role of mononuclear phagocytes in HTLV-III/LAV infection. *Science* 233: 215–219.
- van 't Wout AB, Schuitemaker H, Kootstra NA (2008) Isolation and propagation of HIV-1 on peripheral blood mononuclear cells. *Nat Protoc* 3: 363–370.
- Ochiel DO, Ochsenbauer C, Kappes JC, Ghosh M, Fahey JV, et al. (2010) Uterine epithelial cell regulation of DC-SIGN expression inhibits transmitted/founder HIV-1 trans infection by immature dendritic cells. *PLoS One* 5: e14306.
- Gelderblom HC, Vatakis DN, Burke SA, Lawrie SD, Bristol GC, et al. (2008) Viral complementation allows HIV-1 replication without integration. *Retrovirology* 5: 60.
- Ochsenbauer C, Edmonds TG, Ding H, Keele BF, Decker J, et al. (2012) Generation of transmitted/founder HIV-1 infectious molecular clones and characterization of their replication capacity in CD4 T lymphocytes and monocyte-derived macrophages. *J Virol* 86: 2715–2728.
- Edmonds TG, Ding H, Yuan X, Wei Q, Smith KS, et al. (2010) Replication competent molecular clones of HIV-1 expressing Renilla luciferase facilitate the analysis of antibody inhibition in PBMC. *Virology* 408: 1–13.
- Rodriguez-Garcia M, Climent N, Oliva H, Casanova V, Franco R, et al. (2010) Increased alpha-defensins 1–3 production by dendritic cells in HIV-infected individuals is associated with slower disease progression. *PLoS One* 5: e9436.
- Heffron R, Donnell D, Rees H, Celum C, Were E, et al. (2011) Hormonal contraceptive use and risk of HIV-1 transmission: A prospective cohort analysis 6th IAS Conference on HIV Pathogenesis, Treatment and Prevention Rome, ItalyWEAX0206.
- Yu D, Wang W, Yoder A, Spear M, Wu Y (2009) The HIV envelope but not VSV glycoprotein is capable of mediating HIV latent infection of resting CD4 T cells. *PLoS Pathog* 5: e1000633.
- Anderson JL, Hope TJ (2005) Intracellular trafficking of retroviral vectors: obstacles and advances. *Gene Ther* 12: 1667–1678.
- Hickey DK, Fahey JV, Wira CR (2013) Estrogen receptor alpha antagonists mediate changes in CCL20 and CXCL1 secretions in the murine female reproductive tract. *Am J Reprod Immunol* 69: 159–167.

39. Bruce-Keller AJ, Barger SW, Moss NI, Pham JT, Keller JN, et al. (2001) Pro-inflammatory and pro-oxidant properties of the HIV protein Tat in a microglial cell line: attenuation by 17 beta-estradiol. *J Neurochem* 78: 1315–1324.
40. Lee YW, Eum SY, Nath A, Toborek M (2004) Estrogen-mediated protection against HIV Tat protein-induced inflammatory pathways in human vascular endothelial cells. *Cardiovasc Res* 63: 139–148.
41. Orenstein JM, Fox C, Wahl SM (1997) Macrophages as a source of HIV during opportunistic infections. *Science* 276: 1857–1861.
42. Kolesnitchenko V, Snart RS (1992) Regulatory elements in the human immunodeficiency virus type 1 long terminal repeat LTR (HIV-1) responsive to steroid hormone stimulation. *AIDS Res Hum Retroviruses* 8: 1977–1980.
43. Pierdominici M, Maselli A, Colasanti T, Giammarioli AM, Delunardo F, et al. (2010) Estrogen receptor profiles in human peripheral blood lymphocytes. *Immunol Lett* 132: 79–85.
44. Moriarty K, Kim KH, Bender JR (2006) Minireview: estrogen receptor-mediated rapid signaling. *Endocrinology* 147: 5557–5563.
45. Heron PM, Turchan-Cholewo J, Bruce-Keller AJ, Wilson ME (2009) Estrogen receptor alpha inhibits the estrogen-mediated suppression of HIV transcription in astrocytes: implications for estrogen neuroprotection in HIV dementia. *AIDS Res Hum Retroviruses* 25: 1071–1081.
46. Murphy AJ, Guyre PM, Wira CR, Pioli PA (2009) Estradiol regulates expression of estrogen receptor ERalpha46 in human macrophages. *PLoS One* 4: e5539.
47. Cunningham M, Gilkeson G (2011) Estrogen receptors in immunity and autoimmunity. *Clin Rev Allergy Immunol* 40: 66–73.
48. Moulton VR, Holcomb DR, Zajdel MC, Tsokos GC (2012) Estrogen upregulates cyclic AMP response element modulator alpha expression and downregulates interleukin-2 production by human T lymphocytes. *Mol Med* 18: 370–378.
49. Wira CR, Fahey JV (2008) A new strategy to understand how HIV infects women: identification of a window of vulnerability during the menstrual cycle. *AIDS* 22: 1909–1917.
50. Blair RM, Fang H, Branham WS, Hass BS, Dial SL, et al. (2000) The estrogen receptor relative binding affinities of 188 natural and xenochemicals: structural diversity of ligands. *Toxicol Sci* 54: 138–153.
51. White HD, Musey LK, Andrews MM, Yeaman GR, DeMars LR, et al. (2001) Human immunodeficiency virus-specific and CD3-redredirected cytotoxic T lymphocyte activity in the human female reproductive tract: lack of correlation between mucosa and peripheral blood. *J Infect Dis* 183: 977–983.
52. White HD, Yeaman GR, Givan AL, Wira CR (1997) Mucosal immunity in the human female reproductive tract: cytotoxic T lymphocyte function in the cervix and vagina of premenopausal and postmenopausal women. *Am J Reprod Immunol* 37: 30–38.