

Association of Biomarkers with Individual and Multiple Body Sites of Pain: The Johnston County Osteoarthritis Project

Katherine S Norman^{1,2}, Adam P Goode¹⁻³, Carolina Alvarez^{4,5}, David Hu^{4,5}, Steven Z George^{2,3}, Todd A Schwartz^{4,6}, Stephanie T Danyluk², Rebecca Filippo⁷, Virginia B Kraus^{2,8,9}, Janet L Huebner⁸, Rebecca J Cleveland^{4,5}, Joanne M Jordan^{4,5,10,11}, Amanda E Nelson^{4,5}, Yvonne M Golightly^{4,11,12}

¹Department of Population Health Sciences, Duke University, Durham, NC, USA; ²Department of Orthopedic Surgery, Duke University School of Medicine, Durham, NC, USA; ³Duke Clinical Research Institute, Duke University, Durham, NC, USA; ⁴Thurston Arthritis Research Center, University of North Carolina, Chapel Hill, NC, USA; ⁵Department of Medicine, University of North Carolina, Chapel Hill, NC, USA; ⁶Department of Biostatistics, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, USA; ⁷Duke Department of Physical Therapy and Occupational Therapy, Duke University Health System, Durham, NC, USA; ⁸Duke Molecular Physiology Institute, Duke University School of Medicine, Durham, NC, USA; ⁹Duke Department of Medicine, Duke University School of Medicine, Durham, NC, USA; ¹⁰Department of Orthopedics, University of North Carolina, Chapel Hill, NC, USA; ¹¹Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, USA; ¹²College of Allied Health Professions, University of Nebraska Medical Center, Omaha, NE, USA

Correspondence: Adam P Goode, Department of Orthopedic Surgery, Duke Clinical Research Institute, Duke University, Durham, NC, USA, Tel +1 919-681-6157, Fax +1 919-684-1846, Email Adam.Goode@duke.edu

Introduction: Biochemical biomarkers may provide insight into musculoskeletal pain reported at individual or multiple body sites. The purpose of this study was to determine if biomarkers or pressure-pain threshold (PPT) were associated with individual or multiple sites of pain.

Methods: This cross-sectional analysis included 689 community-based participants. Self-reported symptoms (ie, pain, aching, or stiffness) were ascertained about the neck, upper back/thoracic, low back, shoulders, elbows, wrist, hands, hips, knees, ankles, and feet. Measured analytes included CXCL-6, RANTES, HA, IL-6, BDNF, OPG and NPY. A standard dolorimeter measured PPT. Logistic regression was used to determine the association between biomarkers and PPT with individual and summed sites of pain.

Results: Increased IL-6 and HA were associated with knee pain (OR=1.30, 95% CI 1.03, 1.64) and (OR=1.32, 95% CI 1.01, 1.73) respectively; HA was also associated with elbow/wrist/hand pain (OR=1.60, 95% CI 1.22, 2.09). Those with increased NPY levels were less likely to have shoulder pain (OR=0.56, 95% CI 0.33, 0.93). Biomarkers HA (OR=1.50, 95% CI 1.07, 2.10), OPG (OR=1.74, 95% CI 1.00, 3.03), CXCL-6 (OR=1.75, 95% CI 1.02, 3.01) and decreased PPT (OR=3.97, 95% CI 2.22, 7.12) were associated with multiple compared to no sites of pain. Biomarker HA (OR=1.57, 95% CI 1.06, 2.32) and decreased PPT (OR=3.53, 95% CI 1.81, 6.88) were associated with multiple compared to a single site of pain.

Conclusion: Biomarkers of inflammation (HA, OPG, IL-6 and CXCL-6), pain (NPY) and PPT may help to understand the etiology of single and multiple pain sites.

Keywords: epidemiology, pain, musculoskeletal, population health

Introduction

Musculoskeletal pain is a highly prevalent condition and considered the largest source of disability worldwide.¹⁻³ Research has shown individuals with specific sites of musculoskeletal pain are at increased risk of multiple sites of pain or exhibiting higher pain sensitivity, although there remains a lack of understanding on the relationship and development of pain at other sites.^{4,5} Increased number of pain sites is associated with worse physical, mental, and emotional functioning, as well as increased economic burden for both patients and health care systems.^{2,6-9} Moreover, musculoskeletal pain is complicated by chronic conditions, such as cardiovascular disease (CVD), diabetes (DM), and obesity, though evidence is lacking on how the association occurs, particularly in multiple sites of pain.¹⁰ Other risk factors and conditions for multiple sites of pain include older age, female sex, higher body mass index (BMI), lower physical activity measures, lower education, depression and anxiety.^{9,11,12}

As a growing number of individuals report musculoskeletal pain at more than one site, a better understanding of biologic contributors to pain may be necessary to inform pain management strategies.^{13–15} Recently, biochemical markers have been increasingly investigated for their role in chronic pain and heightened pain perception.¹⁶ Inflammatory biomarkers, including interleukin-6 (IL-6), hyaluronan acid (HA), C-C Motif Chemokine Ligand 5 (RANTES), osteoprotegerin (OPG), and C-X-C Motif Chemokine Ligand 6 (CXCL-6), have all been associated with pain intensity and more physical limitations found in chronic pain states.^{17–21} Biochemical markers may provide additional information useful for understanding the experience of individual or multiple sites of pain, particularly when adjusting for common chronic comorbidities. There is a lack of research that examines the associations of demographics, individual level characteristics (including common comorbidities like obesity), and inflammatory biomarkers with characteristics of individual site or multiple sites of pain. More research is needed to better understand the associations between individual sites of pain and pain at multiple sites, especially examining these relationships within the community since, historically, the majority of pain-related studies sample primarily from care-seeking populations.

To our knowledge, no other community-based study has examined the relationship between biochemical biomarkers with known associations with inflammation and pain/stress, and multiple sites of pain. As such, the purposes of this study were to: 1) describe the frequency of specific (individual) and multiple pain sites within a community-based cohort study; and 2) determine the association between demographic, individual level characteristics, and biochemical markers with specific as well as multiple sites of pain. We chose inflammatory biomarkers IL-6, HA, RANTES, OPG, and CXCL-6 as they are commonly studied in pain and chronic pain states, including chronic neuropathic pain, low back pain, fibromyalgia and osteoarthritis.^{17,18,21,22} We also chose biomarkers NPY and BDNF, which are common biomarkers for pain/stress, secondary to known associations with pain regulation and modulation, especially with the structural degradation associated with low back pain and chronic pain.^{22–24} We hypothesize that these certain biochemical biomarkers will be associated with multiple pain sites, regardless of the anatomical location (ie, common factor), and there will be differences among the demographic and individual level characteristics for different individual sites of pain. In addition, we explored factors that were specific to a given anatomical location in order to provide context for the multiple pain site findings and to generate hypotheses for future studies. Although our primary intention was identification of biomarkers associated with pain at single and multiple sites, we also adjusted for other health-related factors, such as medical comorbidities and social determinants of health. This provided an analysis plan to more fully inform our understanding of risk factors associated with pain distribution in a sample of community-based adults.

Methods

The data for this cross-sectional study were collected as part of the Johnston County Osteoarthritis Project (JoCoOA). The JoCoOA project aimed to determine the incidence, prevalence and progression of osteoarthritis among Black and White men and women 45+ years old in a community-based sample ascertained from six townships of Johnston County, North Carolina. Information regarding the sampling strategy and recruitment methods have been described in previous JoCoOA publications.²⁵ Briefly, JoCoOA participants were selected from the community independently of reports of pain site(s), diagnosis of OA, and biomarker status. Participants were enrolled and data were collected via interviews and clinical examinations during 1991–1997 and 2003–2004; individuals who identified as Black were over-sampled. Approximately every 5 years following initial enrollment, follow-up assessments were performed. Data for the present study were collected from 1697 participants during the second follow-up study visit (2006–2010). All participants in JoCoOA have provided informed consent for participation, and JoCoOA has been continuously approved by the institutional review boards of the University of North Carolina and the Centers for Disease Control and Prevention, which conforms to the Declaration of Helsinki.

Pain Outcomes

Outcomes for these analyses were generated from self-reported pain at multiple locations throughout the body during the clinical examination. Presence of pain at each body location was collected at clinical interview by asking participants to answer “yes” or “no” to the question, “On most days, do you have pain, aching or stiffness in your [neck, upper back/thoracic, low back, shoulder, elbow, wrist, hand, hip, knee, ankle, foot]?” For sites that have bilateral options, such as right and left knee, we counted reports of pain at either or both joints as one site of pain. Sites combined for analysis due to sparse counts of pain included: elbow/wrist/

hand, neck/upper back/thoracic, and ankle/foot. We examined two different outcome categories; individual pain (ie, reporting “yes” in one particular body location) and multiple pain sites (ie, reporting “yes” in more than one body location without regard to a specific body site location). Multiple pain sites were further quantified by summing from 0 to 7. We then categorized this into a single variable of no pain sites (0), single pain site (1) and multiple pain sites (2–7). Other studies have generally categorized multiple sites of pain as 3 or more sites to mirror the definition of chronic widespread pain; however, in this analysis we used more than one site of pain as the criterion for having multiple sites of pain to reflect increasing number of pain sites without regard for anatomic relationship.^{26,27} Including individuals without pain at one or more sites allows for a comparison that is not often an option, as most studies include only care-seeking cohorts.

Exposures

Demographic and Individual Level Characteristics

Demographic information including age, gender (male/female), race (Black/White) and educational status (<high school, high school, >high school) were collected by self-report. Age, gender, race, and educational status are traditional demographic features included in studies examining pain; these variables have shown associations with multiple sites of pain in previous research.^{7,9,11,12}

Comorbidities

Body mass index (BMI in kg/m²) was calculated at the time of clinical examination using height without shoes and weight as measured on a balance beam scale. Those with a BMI \geq 30.0 kg/m² were categorized as obese. The covariate DM was self-reported by asking participants a “yes” or “no” question if they had ever been diagnosed with “diabetes or high blood sugar.” CVD included self-reports of heart attack, angina or angina pectoris, congestive heart failure, other heart problems, cerebrovascular accident, and peripheral vascular disease. Diabetes, obesity, and CVD were included in the analysis due to the known associations with musculoskeletal pain, particularly increased number of pain sites.^{10,28–30}

Serum Biochemical Biomarkers

Details regarding collection of biospecimens, as well as the participants with biochemical samples have been described elsewhere.³¹ All samples were collected after completion of morning activity at a time (>1 hour after arising) when these serum markers have attained equilibrium. The biomarkers associated with inflammation selected for this study included interleukin-6 (IL-6), hyaluronan (HA), C-C Motif Chemokine Ligand 5 (RANTES), osteoprotegerin (OPG), and C-X-C Motif Chemokine Ligand 6 (CXCL-6). Serum biomarkers were measured in singlicate as our prior duplicate pilot work has demonstrated excellent reliability and validity (coefficient of variation (CV) below 15% representing good reliability).²² A human control serum sample was quantified in duplicate on all plates to determine intra- (within) and inter- (between) assay coefficients of variation. [Supplementary Table 1](#) includes additional information pertaining to the biomarkers including: biomarker with details of the manufacturer, dilution, means and distributions, intra-and inter-assay coefficients of variation. All samples had concentrations in the detectable range for all biomarkers.

Pressure Pain Threshold

Pressure pain threshold (PPT) is a well-validated measure for pain sensitivity.³² PPT was conducted using a standard mechanical pressure-based dolorimeter (measured in kilograms of pressure) at bilateral upper trapezius sites. All PPT clinical measurements were performed by a single research assistant utilizing an a-priori measurement protocol that has shown good reliability.³² The protocol started with a “practice trial” to show the device to the participant. Following the practice trial, pressure was applied to the upper trapezius at a rate of 1 kg per second until the patient reported pain, or until a maximum pressure of 4 kg was achieved without a report of pain (reported as “>4.0 kg”). Trials were repeated until two consecutive readings were within 0.4 kg, with a maximum of four trials per side. This procedure was performed first on the left trapezius, then on the right for a total of two measurements that were averaged into a single PPT score.³³ This protocol results in a dichotomous measure of PPT with a score of < 4.0 kg indicating a higher pain sensitivity.

Statistical Analysis

Means with standard deviations, medians and ranges were used to describe the distributions of biomarkers. Intra- and inter-assay CVs were used to describe the reliability of each assay; CVs less than 15% represented assays with good reliability. The lower limit of detection for each biomarker, quantification range, and the dilution used for each assay are also reported. Differences with respect to the absence and presence of each outcome were conducted using *t*-tests and chi-square tests as appropriate. Due to skewness, each biomarker was natural log-transformed prior to conducting regression analysis.

Descriptive statistics were used to summarize study subjects and all relevant variables. Counts and percentages were produced for categorical variables, while mean and standard deviation were computed for continuous variables. To assess associations of covariates with presence of pain by site, multivariable logistic regression was used to separately model the log odds of presence of pain at each of seven sites: 1) Neck/Upper Back/Thoracic; 2) Lower Back; 3) Hip(s); 4) Knee(s); 5) Foot/Ankle(s); 6) Shoulder(s); and 7) Elbow/Wrist/Hand(s). To assess associations of covariables with presence of pain at no sites, a single site, or multiple sites, two models were used: 1) Treating the three-level outcome of pain at no sites, a single site, or multiple sites as nominal, for which a multivariable generalized logits regression model was used; 2) Treating the three-level outcome as ordinal, for which a multivariable proportional odds model using cumulative logits was used.

Models included all previously defined demographic, comorbidity, and biomarker variables to produce adjusted odds ratios (aOR) and corresponding 95% confidence intervals (CIs). Complete cases were included for multivariable analyses, excluding subjects with missing covariates given that missing data (7.5%) was well below <10% (Figure 1) among participants with available serum data.

All analyses were performed with SAS version 9.4 (SAS institute Inc., Cary, NC). The significance level for these analyses was determined at 0.05.

Results

An illustration of the selection of the sample used for these analyses can be found in Figure 1. Of the 1697 participants that entered the second follow-up time point, 1686 (99.4%) had complete pain data. Based on the availability and volume of serum biospecimens remaining, biochemical analyses were conducted on a subsample of 689 (40.6%) individuals.

The demographics of the sample and outcome characteristics are presented in Table 1. The average age was 66.0. The majority of participants were women (69.5%) and 33.4% of participants identified as Black. The average BMI was 31.5 ± 6.1 kg/




Figure 1 Flow Diagram of Study Participants.

Table 1 Demographics and Clinical Characteristics of Biomarker Subsample with Complete Data, N=689

	Overall (N=689)
Covariables	
Age (range=50–88), years (mean ± SD)	66.0 ± 7.4
Gender, n (%)	
Male	210 (30.5%)
Female	479 (69.5%)
Race, n (%)	
Black	230 (33.4%)
White	459 (66.6%)
Education, n (%)	
Less than High School	91 (13.2%)
High School	341 (49.5%)
More than High school	257 (37.3%)
Comorbidities	
BMI (range=18–55), kg/m² (mean ± SD)	31.5 ± 6.1
Obese (≥ 30 kg/m², n (%))	373 (54.1%)
CVD, n (%)	158 (22.9%)
DM, n (%)	135 (19.6%)
Biomarkers (log transformed)	
OPG (range=2.2–5.7), ln(pg/mL) (mean ± SD)	4.5 ± 0.4
CXCL-6 (range=4.2–9.8), ln(pg/mL) (mean ± SD)	5.4 ± 0.5
RANTES (range=10–13), ln(pg/mL) (mean ± SD)	12.0 ± 0.5
IL-6 (range=-1.3–4.4), ln(pg/mL) (mean ± SD)	0.3 ± 0.7
HA (range=2.1–7.5), ln(ng/mL) (mean ± SD)	4.0 ± 0.7
BDNF (range=9–11), ln(pg/mL) (mean ± SD)	10.3 ± 0.3
NPY (range=1.1–4.5), ln(pg/mL) (mean ± SD)	3.1 ± 0.4
Dolorimeter (range=0.8–4.0), kg mean ± SD)	3.7 ± 0.7
Pressure pain sensitivity (<4.0 kgs, n (%))	193 (28.0%)
Outcomes	
Anatomic Pain Site, n (%)	
Neck/Upper Back/Thoracic	231 (33.6%)
Lower Back	316 (45.9%)
Hip(s)	259 (37.6%)
Knee (s)	291 (42.3%)
Foot/Ankle (s)	230 (33.4%)
Shoulder (s)	214 (31.1%)
Elbow/Wrist/Hand (s)	326 (47.4%)
Sum of Pain Sites, n (%)	
0	158 (23.0%)
1	95 (13.8%)
2	92 (13.4%)
3	98 (14.2%)
4	79 (11.5%)
5	63 (9.2%)
6	58 (8.4%)
7	45 (6.5%)
2+ Pain Sites	435 (63.2%)

Abbreviations: BMI, Body Mass Index; DM, diabetes mellitus; CVD, cardiovascular disease; OPG, osteoprotegerin; CXCL6, C-X-C Motif Chemokine Ligand 6; RANTES, C-C Motif Chemokine 5; IL-6, interleukin-6; HA, hyaluronan; BDNF, brain derived neurotrophic factor; NPY, neuropeptide-Y.

m², with over half (54.1%) of participants categorized as obese. Twenty-three percent of participants reported having no pain at any of the body sites, approximately 14% had pain at one body site, and 63% had pain at two or more body sites. The most common site of pain was the elbow/wrist/hand (47.4%), closely followed by the lower back (45.9%).

Results for the different demographic and individual level characteristics, and biomarker data at each individual pain site are presented in Table 2, adjusted for age, gender, race, education, the comorbidities of CVD, DM, and obesity. Generally, Black individuals reported less sites of pain, while women and those with lower education reported more pain in the included sites. Obesity and CVD were both associated with higher odds of pain at the low back, hip(s), knee(s), foot/ankle(s) compared to individuals without obesity or CVD. There were higher odds of reporting shoulder pain (OR=1.77, 95% CI 1.21 to 2.59) for obese individuals compared to non-obese individuals. Odds of elbow/wrist/hand pain (OR=1.77, 95% CI 1.15 to 2.74) were higher in those reporting DM compared to those not reporting DM. Increased serum concentration of IL-6 and HA were associated with higher odds of knee pain (OR=1.30, 95% CI 1.03 to 1.64 and OR=1.32, 95% CI 1.01 to 1.73, respectively). Increased serum concentration of HA was also associated with higher odds of elbow/wrist/hand pain (OR=1.60, 95% CI 1.22 to 2.09), while increased serum concentration of NPY was associated with lower odds of shoulder pain (OR=0.56, 95% CI 0.33, 0.93). Those with a PPT <4.0 kg indicating higher pain sensitivity, had higher odds of reporting pain for every site included in this study.

Table 3 describes the relationships between predictors and no, single, and/or multiple sites of pain. Generally, participants who identified as Black had lower odds of reporting pain compared to those who identified as White; while individuals with obesity had higher odds of reporting pain compared to non-obese individuals. Participants with increased serum levels of OPG (OR=1.74, 95% CI 1.00 to 3.03), CXCL-6 (OR=1.75, 95% CI 1.02 to 3.01), and HA (OR=1.50, 95% CI 1.07 to 2.10) had higher odds of reporting multiple sites of pain compared to no pain sites. Participants with increased serum levels of HA also had higher odds (OR=1.57, 95% CI 1.06 to 3.32) of having multiple sites of pain compared to a single site of pain. Participants with higher pain sensitivity on the PPT (<4 kg) had 3 to 4 times higher odds of reporting multiple sites of pain compared to no or single sites of pain. Table 3 also describes the proportional odds cumulated over no, individual, and multiple sites of pain. Largely, participants who identified as Black had lower odds of greater number of pain sites versus no sites of pain, and obese participants had higher odds of greater number of pain sites versus no sites of pain. Participants with increased serum levels of OPG (OR=1.58, 95% CI 1.01 to 2.47), CXCL-6 (OR=1.64, 95% CI 1.06 to 2.55), HA (OR=1.45, 95% CI 1.10 to 1.90), and lower PPT (higher pain sensitivity, OR=3.78, 95% CI 2.39 to 5.99) had higher odds of reporting pain as the number of pain sites increased.

Discussion

In this analysis, we investigated associations between demographics, individual level characteristics, and biochemical biomarkers for specific and multiple pain sites. To the best of our knowledge, this is one of the first community-based studies within the United States to examine these relationships. This unique, community-based sample was diverse and offered a variety of plausible biological markers for analysis of biomarker associations with pain sites. These findings offer valuable information regarding pain distributions to complement information provided by cohorts of patients seeking care for musculoskeletal pain conditions. In this community-based cohort, the majority of participants reported multiple sites of pain, while only a quarter reported no pain, and even fewer reported a single site of pain.

One objective during this analysis was to evaluate the frequency of specific and multiple pain sites within a community-based cohort. Our results related to the sum of pain sites are consistent with other studies that found a majority of people reported multiple sites of pain. Roughly two-thirds of the sample reported 2 or more sites of pain, which is slightly higher than the prevalence of multi-site pain in the general public.¹⁵ This increased prevalence may have been influenced by the older average age of our population (66 years old) compared to the general population. Consistent with our results, a study examining adults over the age of 65, observed that approximately 75% of adults reported multiple sites of pain.³⁴ These findings indicate increased musculoskeletal pain seen in aging population, which could contribute to functional limitations or disability.^{35,36} Participants who identified as Black demonstrated reduced odds of pain at most of the individual sites, as well as reduced odds of reporting both single and multiple sites of pain compared to White participants. Literature suggests that pain is underreported in individuals who identify as Black although pain severity tends to be higher in this population.³⁷ Due to the nature of the data collection, we are unable to

Table 2 Cross-Sectional Associations* Between Presence of Individual Pain Sites with Demographics, Comorbidities, and Biomarkers, N=689

Covariables	Outcomes						
	Neck/Upper Back/Thoracic	Lower Back	Hip(s)	Knee(s)	Foot/ Ankle (s)	Shoulder(s)	Elbow/Wrist/ Hand (s)
	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)
DEMOGRAPHICS							
Age (1 year increase)	0.99 (0.96, 1.01)	0.98 (0.96, 1.01)	0.97 (0.95, 0.99)	0.98 (0.96, 1.01)	0.98 (0.95, 1.01)	0.97 (0.94, 1.00)	0.97 (0.94, 0.99)
Gender (F vs M)	1.58 (1.04, 2.40)	1.16 (0.79, 1.69)	1.21 (0.82, 1.80)	1.35 (0.91, 2.00)	2.06 (1.34, 3.18)	1.17 (0.76, 1.79)	1.71 (1.16, 2.51)
Race (B vs W)	0.45 (0.27, 0.76)	0.43 (0.26, 0.70)	0.52 (0.31, 0.85)	0.76 (0.47, 1.24)	0.42 (0.25, 0.72)	0.66 (0.39, 1.12)	0.48 (0.29, 0.78)
Education: HS vs >HS	1.37 (0.93, 2.01)	1.19 (0.83, 1.71)	1.10 (0.76, 1.59)	1.08 (0.75, 1.57)	1.00 (0.68, 1.48)	1.15 (0.77, 1.71)	0.99 (0.69, 1.42)
Education: <HS vs >HS	2.28 (1.32, 3.93)	2.20 (1.28, 3.77)	1.83 (1.08, 3.12)	1.31 (0.76, 2.24)	1.95 (1.13, 3.36)	1.78 (1.02, 3.11)	1.57 (0.91, 2.70)
COMORBIDITIES							
Obesity	1.09 (0.75, 1.57)	1.67 (1.18, 2.35)	1.44 (1.01, 2.05)	2.11 (1.48, 2.99)	1.51 (1.04, 2.19)	1.77 (1.21, 2.59)	1.20 (0.85, 1.71)
CVD	1.23 (0.83, 1.84)	1.67 (1.13, 2.46)	1.48 (1.01, 2.18)	1.51 (1.02, 2.23)	1.53 (1.03, 2.28)	1.16 (0.77, 1.74)	1.23 (0.84, 1.82)
DM	1.23 (0.79, 1.91)	1.15 (0.75, 1.76)	1.26 (0.82, 1.93)	1.18 (0.77, 1.81)	1.30 (0.84, 2.03)	1.19 (0.76, 1.85)	1.77 (1.15, 2.74)
BIOMARKERS (for a 1 unit increase in ln(biomarker))							
OPG, ln(pg/mL)	1.20 (0.75, 1.90)	1.29 (0.84, 2.00)	1.53 (0.98, 2.41)	1.10 (0.70, 1.71)	0.69 (0.43, 1.09)	1.07 (0.67, 1.71)	1.35 (0.87, 2.10)
CXCL6, ln(pg/mL)	1.18 (0.76, 1.82)	1.30 (0.86, 1.97)	1.03 (0.67, 1.57)	1.16 (0.76, 1.77)	1.34 (0.86, 2.07)	1.14 (0.73, 1.80)	1.28 (0.84, 1.96)
RANTES, ln(pg/mL)	0.98 (0.64, 1.50)	1.01 (0.68, 1.51)	1.35 (0.89, 2.05)	1.22 (0.81, 1.84)	1.45 (0.93, 2.25)	1.24 (0.80, 1.93)	0.92 (0.61, 1.38)
IL-6, ln(pg/mL)	1.13 (0.89, 1.44)	1.08 (0.86, 1.36)	0.97 (0.76, 1.22)	1.30 (1.03, 1.64)	1.14 (0.89, 1.46)	1.15 (0.90, 1.48)	1.04 (0.82, 1.31)
HA, ln(ng/mL)	1.02 (0.78, 1.34)	1.10 (0.85, 1.43)	1.10 (0.85, 1.43)	1.32 (1.01, 1.73)	1.14 (0.87, 1.51)	1.14 (0.86, 1.51)	1.60 (1.22, 2.09)
BDNF, ln(pg/mL)	0.95 (0.50, 1.82)	0.71 (0.39, 1.31)	0.70 (0.37, 1.31)	0.60 (0.32, 1.13)	0.77 (0.40, 1.48)	0.72 (0.37, 1.39)	1.02 (0.55, 1.89)
NPY, ln(pg/mL)	0.78 (0.47, 1.28)	0.80 (0.50, 1.28)	0.81 (0.50, 1.32)	0.66 (0.41, 1.08)	0.80 (0.49, 1.32)	0.56 (0.33, 0.93)	0.66 (0.41, 1.07)
Pressure pain sensitivity: Dolorimeter <4 vs 4 kg/s	3.10 (2.12, 4.54)	2.35 (1.61, 3.43)	2.44 (1.68, 3.56)	2.46 (1.68, 3.61)	2.53 (1.72, 3.70)	3.13 (2.12, 4.62)	2.76 (1.87, 4.06)

Notes: *Results from seven, separate by site, multivariable logistic regression models for dichotomous presence of pain; aOR=Odds Ratio, represents the association between each covariate and the outcome while simultaneously adjusted for all other explanatory variables in the table; 95% CI= 95% Confidence Interval; **Bold** values indicate a 95% CI where the null effect of OR=1 is not included, at a 0.05 level of significance.

Abbreviations: F, Female; M, Male; B, Black; W, White; HS, High School Education; >HS, more than high school education; <HS, less than high school education; Obesity, Body Mass Index \geq 30 kg/m²; DM, diabetes mellitus; CVD, cardiovascular disease; OPG, osteoprotegerin; CXCL6, C-X-C Motif Chemokine Ligand 6; RANTES, C-C Motif Chemokine 5; IL-6, interleukin-6; HA, hyaluronan; BDNF, brain derived neurotrophic factor; NPY, neuropeptide-Y.

Table 3 Cross-Sectional Associations* Between Multiple, Single and No Pain Sites with Demographics, Comorbidities, and Biomarkers, N=689

Covariables	Generalized Logits Model Treating Three-Level Outcome as Nominal			Proportional Odds Model Using Cumulative Logits Treating Three-Level Outcome as Ordinal
	Single Site of Pain vs No Pain	Multiple Sites of Pain vs No Pain	Multiple Sites of Pain vs Single Site of Pain	Cumulated Over Multiple to No Sites of Pain
	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)
DEMOGRAPHICS				
Age (1 year increase)	0.97 (0.93, 1.01)	0.97 (0.94, 1.00)	1.00 (0.96, 1.03)	0.98 (0.95, 1.00)
Gender (F vs M)	1.17 (0.66, 2.06)	1.51 (0.97, 2.36)	1.30 (0.77, 2.17)	1.40 (0.97, 2.01)
Race (B vs W)	0.33 (0.15, 0.73)	0.25 (0.13, 0.45)	0.76 (0.37, 1.57)	0.33 (0.20, 0.54)
Education: HS vs >HS	0.89 (0.50, 1.57)	1.11 (0.71, 1.72)	1.25 (0.75, 2.08)	1.11 (0.78, 1.59)
Education: <HS vs >HS	1.39 (0.53, 3.68)	1.86 (0.88, 3.96)	1.34 (0.60, 2.99)	1.67 (0.93, 3.02)
COMORBIDITIES				
Obesity	1.77 (1.00, 3.12)	2.34 (1.52, 3.62)	1.32 (0.81, 2.17)	2.02 (1.43, 2.87)
CVD	1.39 (0.69, 2.77)	1.58 (0.93, 2.71)	1.14 (0.64, 2.03)	1.40 (0.92, 2.13)
DM	1.03 (0.45, 2.31)	1.60 (0.88, 2.91)	1.56 (0.79, 3.06)	1.51 (0.94, 2.43)
BIOMARKERS (for a 1 unit increase in ln(biomarker))				
OPG, ln(pg/mL)	1.36 (0.67, 2.73)	1.74 (1.00, 3.03)	1.28 (0.69, 2.38)	1.58 (1.01, 2.47)
CXCL6, ln(pg/mL)	1.22 (0.74, 2.00)	1.75 (1.02, 3.01)	1.21 (0.65, 2.25)	1.64 (1.06, 2.55)
RANTES, ln(pg/mL)	1.26 (0.66, 2.39)	1.22 (0.74, 2.00)	0.97 (0.54, 1.72)	1.18 (0.79, 1.76)
IL-6, ln(pg/mL)	1.10 (0.75, 1.60)	1.08 (0.81, 1.44)	0.99 (0.71, 1.38)	1.02 (0.81, 1.29)
HA, ln(ng/mL)	0.95 (0.61, 1.48)	1.50 (1.07, 2.10)	1.57 (1.06, 2.32)	1.45 (1.10, 1.90)
BDNF, ln(pg/mL)	0.61 (0.23, 1.63)	0.64 (0.30, 1.37)	1.06 (0.44, 2.55)	0.69 (0.37, 1.28)
NPY, ln(pg/mL)	0.89 (0.42, 1.88)	0.63 (0.35, 1.15)	0.72 (0.36, 1.41)	0.69 (0.43, 1.11)
Pressure pain sensitivity: Dolorimeter <4 vs 4 kg/s	1.13 (0.49, 2.57)	3.97 (2.22, 7.12)	3.53 (1.81, 6.88)	3.78 (2.39, 5.99)

Notes: *Results from a multivariable generalized logits model and a multivariable cumulative logits model for a three-level, polytomous outcome of no pain, single site pain, and multiple site pain; aOR=Odds Ratio, represents the association between each covariate and the outcome while simultaneously adjusted for all other explanatory variables in the table; 95% CI= 95% Confidence Interval; Bold values indicate a 95% CI where the null effect of OR=1 is not included, at a 0.05 level of significance.

Abbreviations: F, Female; M, Male; B, Black; W, White; HS, High School Education, >HS, more than high school education; <HS, less than high school education; Obesity, Body Mass Index>30 kg/m²; DM, diabetes mellitus; CVD, cardiovascular disease; OPG, osteoprotegerin; CXCL6, C-X-C Motif Chemokine Ligand 6; RANTES, C-C Motif Chemokine 5; IL-6, interleukin-6; HA, hyaluronan; BDNF, brain derived neurotrophic factor; NPY, neuropeptide-Y.

determine if participants who identify as Black in the JoCoOA sample underreport their pain or truly have less sites of pain. Female gender and lower educational status were also significantly associated with individual sites of pain, but these demographic features were not determined to be statistically significant for single and multiple site pain analyses.

Overwhelmingly, the most consistent factor in individual and multiple pain sites was the PPT, as it was statistically significant for every individual site analysis. Furthermore, those with multiple pain sites had significantly lower PPT thresholds (higher pain sensitivity) than those without pain or only a single site pain; however, there was no significant difference in PPT for single site compared to no pain. This finding indicates that those experiencing multiple pain sites may have increased pain sensitivity and altered pain processing.³⁸ Increased pain sensitivity often leads to worse outcomes and additional healthcare needs, so it is vital to identify objective measures to facilitate identification of individuals experiencing altered pain processing early in order to target appropriate treatment interventions.³⁹ This nonclinical, population-based sample combined with PPT as a form of quantitative sensory testing allows for improved understanding of musculoskeletal pain and pain sensitivity.⁴⁰ Use of PPT in the clinical setting is feasible as it is a low-cost application with high user and patient acceptability.^{32,41,42} Thus, use in the clinical setting with further development and understanding of PPT thresholds may help to identify individuals at risk for multiple sites of musculoskeletal pain.⁴² A deeper understanding of how pain sensitivity

contributes to perceived functional limitations will also improve clinical management and outcomes of patients with multiple sites of pain. Given the unique design of examining association with presence of pain and quantitative sensory testing in the form of PPT, we are contributing to an understanding of identification of potential risk factors for pain in a population-based cohort. Our results corroborate that lower PPT thresholds may identify patients experiencing altered pain processing to develop means of addressing their altered pain perceptions early, and even in community-based samples this higher pain sensitivity may be linked with having more sites of pain.

Even after accounting for PPT, we identified two biochemical biomarkers that were associated with multiple pain sites (OPG and CXCL-6), indicating potential roles of these biomarkers in more complex, multiple sites of pain. Higher levels of CXCL-6 have been associated with chronic neuropathic pain, and moderate to severe low back pain, implicating this biochemical marker in possible pain pathways.^{21,43} The biomarker OPG is a neuropeptide that has been shown to be related to neurogenic inflammation, especially in conditions such as complex regional pain syndrome, which may contribute to mechanisms for multiple sites of pain.⁴⁴ Our findings showed elevated HA levels were seen in both multiple sites of pain and individual pain sites. Increased levels of biomarker HA have been seen in patients with OA, particularly in the spine, hip, and knee joints; previous studies indicate that elevated levels of HA increase with higher levels of OA burden due to the role of HA in inflammatory pathways associated with symptoms of increased stiffness and swelling.^{45,46} Additionally, our results demonstrated an association of HA with elbow/wrist/hand pain which is consistent with previous research demonstrating elevated HA levels are associated with increased number of arthritic joints, especially in the hand, which can contribute to increased pain and stiffness.⁴⁷⁻⁴⁹ We found IL-6 and NPY to be associated with specific sites of pain, knee and shoulder pain respectively. Biomarker IL-6 is a cytokine, and has been found to play a role as an inflammatory mediator in several pain pathways which can exacerbate pain.⁵⁰ NPY has also been shown to play a role in pain perception, specifically in low back pain and chronic pain, but there is little research on the role of NPY, shoulder pain and OA symptoms.^{22,51} This cross-sectional analysis corroborates the hypothesis that biomarkers may play a role in both individual and multiple sites of pain. As such, our analyses support the further examination of the utility of biochemical biomarkers as objective indicators of individual pain sites or multiple sites of pain.

Several of the biomarkers associated with pain are also known to be associated with other comorbid, chronic conditions. Although we adjusted for the common comorbidities of DM, CVD, and obesity, since this was a cross-sectional analysis, we cannot determine a temporal or causal component of these conditions on the development of pain at individual or multiple sites. These conditions are hypothesized to be associated with multiple sites of pain due to compounding issues which contribute to comorbidities and pain such as poor diet, systemic inflammation, and altered ability to participate in physical activity.²⁸⁻³⁰ Our results show those with obesity have almost double the odds of having multiple sites pain compared to those with single site and no pain. Obesity and multiple sites of pain have been shown to have a strong cross-sectional association, particularly in OA-related pain due to increased mechanical loads on joints secondary to increased body mass.^{6,7,12,52} Altered biomarker levels, including IL-6, can signify dysregulation of inflammatory systems, which has been shown to be associated with increased pain and obesity in certain populations.^{53,54} Elevated BMI is associated with chronic inflammation and pain, which could be explained by biomarkers included in this study.⁵⁵ Again, our data are cross-sectional so we cannot determine a temporal relationship, though our results indicate further analysis of biomarkers in contribution to systemic inflammation which may exacerbate conditions of pain and/or obesity. Evidence also supports that musculoskeletal pain, especially OA pain, is associated with CVD, and individuals with increased number of pain sites are known to have an increased risk of cardiovascular death.^{28,30} Finally, a longitudinal study showed that patients with diabetes have a higher cumulative-incidence of musculoskeletal pain compared to non-diabetic groups over a 10-year period.²⁹ These conditions, especially obesity, are hypothesized to be linked with musculoskeletal pain through inflammatory pathways (ie, metabolic syndrome) which can also exacerbate conditions such as DM and CVD.^{54,56}

Our study has several strengths including the fact that it is a community-based study, examining the associations among demographic, individual level characteristics, and biochemical markers, and individual as well as multiple sites of pain. Our large sample size of participants of Black or White race of both genders allowed for several comparisons within the analyses. As such, these analyses may be more generalizable to the population outside of the more commonly studied clinically based cohorts by providing information from a community-based sample.⁵⁸ This furthers our understanding of pain sensitivity and pain patterns in a larger population which has been identified as a critical gap in the pain research literature.^{40,57} However, this study does have several limitations. Despite the fact that our biomarker findings were consistent with other studies that found

correlations with these biomarkers and musculoskeletal pain, several biomarkers included in this study did not demonstrate an association with individual pain sites or multiple sites of pain. The lack of significant findings for several biomarkers could be attributed to the fact that pain was defined as dichotomous (yes/no), whereas other biomarker studies use a continuum of pain intensity ratings, duration, or other measures as the definition of pain or pain interference.^{17,18,24,47} Another limitation relates to the cross-sectional design of this study resulting in our only being able to study the association between biomarkers and individual sites of pain or multiple sites of pain, not the causal relationships between these biomarkers and the development of multiple sites of pain. Additionally, biochemical markers were only measured on a subset of the population, and therefore may not be representative of the entire cohort. Our measurement of PPT was dichotomized per the protocol of this epidemiologic study differs from PPT values which can also be treated as continuous. Defining a cutoff of PPT for pain sensitivity may be more pragmatic and useful to identify patients at risk for increased pain sensitivity. Finally, we collected comorbidities from participant self-report on DM and CVD which may be prone to measurement error. In addition, we did not have information on participants' history of trauma, surgery, or other conditions which might contribute to pain. Future studies are warranted to examine whether these biochemical biomarkers may predict the progression from single to multiple sites of body pain.

In summary, community members having multiple pain sites are more likely to have higher pain sensitivity and have associations with certain biochemical markers, especially OPG, CXCL-6, and HA. In contrast, there were two biochemical markers (IL-6 and NPY), that were only associated with specific pain sites. These findings support our goal of identifying biochemical markers, particularly those related to inflammation, that may aid in objectively distinguishing risk factors for pain that progresses from one to multiple sites. Combining these findings with other clinical outcomes, and potentially diagnostic imaging, may provide further insight into the development of chronic, multi-site pain, and associated disabilities.

Acknowledgment

This work was supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS) R01AR071440 (Goode, George, Kraus, Jordan, Golightly, Cleveland, Schwartz), R01AR075399 (Goode, DeFrate, George, Kraus) and (Goode, K24 AR079594). The Johnston County Osteoarthritis Project is supported in part by cooperative agreements S043, S1734, and S3486 from the Centers for Disease Control and Prevention (CDC)/ Association of Schools of Public Health; the NIAMS Multipurpose Arthritis and Musculoskeletal Disease Center grant 5-P60-AR30701; the NIAMS Multidisciplinary Clinical Research Center grant 5 P60 AR49465-03 and NIAMS Core Center for Clinical Research P30-AR-072580; and the National Institute on Aging (NIA) P30-AG-028716. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the CDC, NIA, or NIAMS. The data for this study are held securely at the University of North Carolina, Chapel Hill and are restricted due to data sharing agreements so they are not publicly available.

Disclosure

Dr Adam P Goode reports grants from NIH, during the conduct of the study. Dr Steven Z George reports grants from NIH paid to his institution, during the conduct of the study; received honorarium and travel for Deputy Editor work from American Physical Therapy Association, royalties from Rehab Essentials for online education module, honorarium and travel for Scientific Advisory Board from Med Risk, outside the submitted work. All authors disclose they have no other financial or personal relationships with other people or organizations that could potentially and inappropriately bias their work and conclusions.

References

1. Blyth FM, Briggs AM, Schneider CH, Hoy DG, March LM. The global burden of musculoskeletal pain-where to from here? *Am J Public Health*. 2019;109(1):35–40. doi:10.2105/AJPH.2018.304747
2. Brooks PM. The burden of musculoskeletal disease—a global perspective. *Clin Rheumatol*. 2006;25(6):778–781. doi:10.1007/s10067-006-0240-3
3. Briggs AM, Woolf AD, Dreinhofer K, et al. Reducing the global burden of musculoskeletal conditions. *Bull World Health Organ*. 2018;96(5):366–368. doi:10.2471/BLT.17.204891
4. Felson DT, Niu J, Quinn EK, et al. Multiple nonspecific sites of joint pain outside the knees develop in persons with knee pain. *Arthritis Rheumatol*. 2017;69(2):335–342. doi:10.1002/art.39848

5. Neogi T, Frey-Law L, Scholz J, et al. Sensitivity and sensitisation in relation to pain severity in knee osteoarthritis: trait or state? *Ann Rheum Dis*. 2015;74(4):682–688. doi:10.1136/annrheumdis-2013-204191
6. Mundal I, Grawe RW, Bjorngaard JH, Linaker OM, Fors EA. Prevalence and long-term predictors of persistent chronic widespread pain in the general population in an 11-year prospective study: the HUNT study. *BMC Musculoskelet Disord*. 2014;15:213. doi:10.1186/1471-2474-15-213
7. Kamaleri Y, Natvig B, Ihlebaek CM, Benth JS, Bruusgaard D. Number of pain sites is associated with demographic, lifestyle, and health-related factors in the general population. *Eur J Pain*. 2008;12(6):742–748. doi:10.1016/j.ejpain.2007.11.005
8. Lacey RJ, Belcher J, Rathod T, Wilkie R, Thomas E, McBeth J. Pain at multiple body sites and health-related quality of life in older adults: results from the North Staffordshire Osteoarthritis project. *Rheumatology*. 2014;53(11):2071–2079. doi:10.1093/rheumatology/keu240
9. Mun CJ, Ruhlman L, Karoly P. Examining the adjustment patterns of adults with multiple chronic pain conditions and multiple pain sites: more pain, no gain. *J Pain Jan*. 2020;21(1–2):108–120. doi:10.1016/j.jpain.2019.06.002
10. van der Zee-Neuen A, Putrik P, Ramiro S, et al. Impact of chronic diseases and multimorbidity on health and health care costs: the additional role of musculoskeletal disorders. *Arthritis Care Res*. 2016;68(12):1823–1831. doi:10.1002/acr.22913
11. Larsson B, Bjork J, Borsbo B, Gerdle B. A systematic review of risk factors associated with transitioning from regional musculoskeletal pain to chronic widespread pain. *Eur J Pain*. 2012;16(8):1084–1093. doi:10.1002/j.1532-2149.2012.00117.x
12. Pan F, Laslett L, Blizzard L, et al. Associations between fat mass and multisite pain: a five-year longitudinal study. *Arthritis Care Res*. 2017;69(4):509–516. doi:10.1002/acr.22963
13. Kamaleri Y, Natvig B, Ihlebaek CM, Bruusgaard D. Localized or widespread musculoskeletal pain: does it matter? *Pain*. 2008;138(1):41–46. doi:10.1016/j.pain.2007.11.002
14. Croft P, Dunn KM, Von Korf M. Chronic pain syndromes: you can't have one without another. *Pain*. 2007;131(3):237–238. doi:10.1016/j.pain.2007.07.013
15. Hartvigsen J, Davidsen M, Hestbaek L, Sogaard K, Roos EM. Patterns of musculoskeletal pain in the population: a latent class analysis using a nationally representative interviewer-based survey of 4817 Danes. *Eur J Pain*. 2013;17(3):452–460. doi:10.1002/j.1532-2149.2012.00225.x
16. Klyne DM, Barbe MF, van den Hoorn W, Hodges PW. ISSLS PRIZE IN CLINICAL SCIENCE 2018: longitudinal analysis of inflammatory, psychological, and sleep-related factors following an acute low back pain episode-The good, the bad, and the ugly. *Eur Spine J*. 2018;27(4):763–777. doi:10.1007/s00586-018-5490-7
17. Koch A, Zacharowski K, Boehm O, et al. Nitric oxide and pro-inflammatory cytokines correlate with pain intensity in chronic pain patients. *Inflamm Res*. 2007;56(1):32–37. doi:10.1007/s00011-007-6088-4
18. Sowa GA, Perera S, Bechara B, et al. Associations between serum biomarkers and pain and pain-related function in older adults with low back pain: a pilot study. *J Am Geriatr Soc*. 2014;62(11):2047–2055. doi:10.1111/jgs.13102
19. Pernambuco AP, Schetino LP, Alvim CC, et al. Increased levels of IL-17A in patients with fibromyalgia. *Clin Exp Rheumatol*. 2013;31(6 Suppl 79):S60–3.
20. Xue JB, Zhan XL, Wang WJ, Yan YG, Liu C. OPG rs2073617 polymorphism is associated with upregulated OPG protein expression and an increased risk of intervertebral disc degeneration. *Exp Ther Med*. 2016;12(2):702–710. doi:10.3892/etm.2016.3342
21. Backryd E, Lind AL, Thulin M, Larsson A, Gerdle B, Gordh T. High levels of cerebrospinal fluid chemokines point to the presence of neuroinflammation in peripheral neuropathic pain: a cross-sectional study of 2 cohorts of patients compared with healthy controls. *Pain*. 2017;158(12):2487–2495. doi:10.1097/j.pain.0000000000001061
22. Goode AP, Schwartz TA, Kraus VB, et al. Inflammatory, structural, and pain biochemical biomarkers may reflect radiographic disc space narrowing: the Johnston County Osteoarthritis project. *J Orthop Res*. 2020;38(5):1027–1037. doi:10.1002/jor.24534
23. Malfait AM, Miller RE, Block JA. Targeting neurotrophic factors: novel approaches to musculoskeletal pain. *Pharmacol Ther*. 2020;211:107553. doi:10.1016/j.pharmthera.2020.107553
24. Diz JBM, de Souza Moreira B, Felicio DC, et al. Brain-derived neurotrophic factor plasma levels are increased in older women after an acute episode of low back pain. *Arch Gerontol Geriatr*. 2017;71:75–82. doi:10.1016/j.archger.2017.03.005
25. Goode AP, Marshall SW, Renner JB, et al. Lumbar spine radiographic features and demographic, clinical, and radiographic knee, Hip, and hand osteoarthritis. *Arthritis Care Res*. 2012;64(10):1536–1544. doi:10.1002/acr.21720
26. Mundal I, Grawe RW, Bjorngaard JH, Linaker OM, Fors EA. Psychosocial factors and risk of chronic widespread pain: an 11-year follow-up study—the HUNT study. *Pain*. 2014;155(8):1555–1561. doi:10.1016/j.pain.2014.04.033
27. Hunt IM, Silman AJ, Benjamin S, McBeth J, Macfarlane GJ. The prevalence and associated features of chronic widespread pain in the community using the 'Manchester' definition of chronic widespread pain. *Rheumatology*. 1999;38(3):275–279. doi:10.1093/rheumatology/38.3.275
28. McBeth J, Symmons DP, Silman AJ, et al. Musculoskeletal pain is associated with a long-term increased risk of cancer and cardiovascular-related mortality. *Rheumatology*. 2009;48(1):74–77. doi:10.1093/rheumatology/ken424
29. Pai LW, Hung CT, Li SF, Chen LL, Chung Y, Liu HL. Musculoskeletal pain in people with and without type 2 diabetes in Taiwan: a population-based, retrospective cohort study. *BMC Musculoskelet Disord*. 2015;16:364. doi:10.1186/s12891-015-0819-4
30. Williams A, Kamper SJ, Wiggers JH, et al. Musculoskeletal conditions may increase the risk of chronic disease: a systematic review and meta-analysis of cohort studies. *BMC Med*. 2018;16(1):167. doi:10.1186/s12916-018-1151-2
31. Goode AP, Marshall SW, Kraus VB, et al. Association between serum and urine biomarkers and lumbar spine individual radiographic features: the Johnston County Osteoarthritis Project. *Osteoarthritis Cartilage*. 2012;20(11):1286–1293. doi:10.1016/j.joca.2012.08.003
32. Wylde V, Palmer S, Learmonth ID, Dieppe P. Test-retest reliability of quantitative sensory testing in knee osteoarthritis and healthy participants. *Osteoarthritis Cartilage*. 2011;19(6):655–658. doi:10.1016/j.joca.2011.02.009
33. Goode AP, Shi XA, Gracely RH, Renner JB, Jordan JM. Associations between pressure-pain threshold, symptoms, and radiographic knee and Hip osteoarthritis. *Arthritis Care Res*. 2014;66(10):1513–1519. doi:10.1002/acr.22321
34. Patel KV, Guralnik JM, Dansie EJ, Turk DC. Prevalence and impact of pain among older adults in the United States: findings from the 2011 national health and aging trends study. *Pain*. 2013;154(12):2649–2657. doi:10.1016/j.pain.2013.07.029
35. Butera KA, Roff SR, Buford TW, Cruz-Almeida Y. The impact of multisite pain on functional outcomes in older adults: biopsychosocial considerations. *J Pain Res*. 2019;12:1115–1125. doi:10.2147/JPR.S192755
36. Chang AY, Skirbekk VF, Tyrovolas S, Kassebaum NJ, Dieleman JL. Measuring population ageing: an analysis of the global burden of disease study 2017. *Lancet Public Health*. 2019;4(3):e159–e167. doi:10.1016/S2468-2667(19)30019-2
37. Mossey JM. Defining racial and ethnic disparities in pain management. *Clin Orthop Relat Res*. 2011;469(7):1859–1870. doi:10.1007/s11999-011-1770-9

38. Sofat N, Ejindu V, Kiely P. What makes osteoarthritis painful? The evidence for local and central pain processing. *Rheumatology*. 2011;50(12):2157–2165. doi:10.1093/rheumatology/ker283
39. Coronado RA, Bialosky JE, Robinson ME, et al. Pain sensitivity subgroups in individuals with spine pain: potential relevance to short-term clinical outcome. *Phys Ther*. 2014;94(8):1111–1122. doi:10.2522/ptj.20130372
40. Waller R, Smith AJ, Graven-Nielsen T, et al. Role of population-based cohorts in understanding the emergence and progression of musculoskeletal pain. *Pain*. 2022;163(1):58–63. doi:10.1097/j.pain.0000000000002316
41. van Wilgen P, van der Noord R, Zwerver J. Feasibility and reliability of pain pressure threshold measurements in patellar tendinopathy. *J Sci Med Sport*. 2011;14(6):477–481. doi:10.1016/j.jsams.2011.05.004
42. Georgopoulos V, Akin-Akinyosoye K, Zhang W, McWilliams DF, Hendrick P, Walsh DA. Quantitative sensory testing and predicting outcomes for musculoskeletal pain, disability, and negative affect: a systematic review and meta-analysis. *Pain*. 2019;160(9):1920–1932. doi:10.1097/j.pain.0000000000001590
43. Grad S, Bow C, Karpinen J, et al. Systemic blood plasma CCL5 and CXCL6: potential biomarkers for human lumbar disc degeneration. *Eur Cell Mater*. 2016;31:1–10. doi:10.22203/ecm.v031a01
44. Littlejohn G. Neurogenic neuroinflammation in fibromyalgia and complex regional pain syndrome. *Nat Rev Rheumatol*. 2015;11(11):639–648. doi:10.1038/nrrheum.2015.100
45. Elliott AL, Kraus VB, Luta G, et al. Serum hyaluronan levels and radiographic knee and Hip osteoarthritis in African Americans and Caucasians in the Johnston County osteoarthritis project. *Arthritis Rheum*. 2005;52(1):105–111. doi:10.1002/art.20724
46. Wells AF, Klareskog L, Lindblad S, Laurent TC. Correlation between increased hyaluronan localized in arthritic synovium and the presence of proliferating cells. A role for macrophage-derived factors. *Arthritis Rheum*. 1992;35(4):391–396. doi:10.1002/art.1780350405
47. Sasaki E, Tsuda E, Yamamoto Y, et al. Serum hyaluronan levels increase with the total number of osteoarthritic joints and are strongly associated with the presence of knee and finger osteoarthritis. *Int Orthop*. 2013;37(5):925–930. doi:10.1007/s00264-013-1849-x
48. Aslam I, Perjar I, Shi XA, et al. Associations between biomarkers of joint metabolism, hand osteoarthritis, and hand pain and function: the Johnston County Osteoarthritis project. *J Rheumatol*. 2014;41(5):938–944. doi:10.3899/jrheum.130904
49. Filkova M, Senolt L, Braun M, et al. Serum hyaluronic acid as a potential marker with a predictive value for further radiographic progression of hand osteoarthritis. *Osteoarthritis Cartilage*. 2009;17(12):1615–1619. doi:10.1016/j.joca.2009.06.002
50. Greene MA, Loeser RF. Aging-related inflammation in osteoarthritis. *Osteoarthritis Cartilage*. 2015;23(11):1966–1971. doi:10.1016/j.joca.2015.01.008
51. Diaz-delCastillo M, Woldbye DPD, Heegaard AM. Neuropeptide Y and its involvement in chronic pain. *Neuroscience*. 2018;387:162–169. doi:10.1016/j.neuroscience.2017.08.050
52. Magnusson K, Osteras N, Mowinkel P, Natvig B, Hagen KB. No strong temporal relationship between obesity and multisite pain—results from a population-based 20-year follow-up study. *Eur J Pain*. 2014;18(1):120–127. doi:10.1002/j.1532-2149.2013.00338.x
53. Hainsworth KR, Simpson PM, Raff H, Grayson MH, Zhang L, Weisman SJ. Circulating inflammatory biomarkers in adolescents: evidence of interactions between chronic pain and obesity. *Pain Rep*. 2021;6(1):e916. doi:10.1097/PR9.0000000000000916
54. Collins KH, Herzog W, MacDonald GZ, et al. Obesity, metabolic syndrome, and musculoskeletal disease: common inflammatory pathways suggest a central role for loss of muscle integrity. *Front Physiol*. 2018;9:112. doi:10.3389/fphys.2018.00112
55. Seaman DR. Body mass index and musculoskeletal pain: is there a connection? *Chiropr Man Therap*. 2013;21(1):15. doi:10.1186/2045-709X-21-15
56. Esser N, Legrand-Poels S, Piette J, Scheen AJ, Paquot N. Inflammation as a link between obesity, metabolic syndrome and type 2 diabetes. *Diabetes Res Clin Pract*. 2014;105(2):141–150. doi:10.1016/j.diabres.2014.04.006
57. Davis KD, Aghaepour N, Ahn AH, et al. Discovery and validation of biomarkers to aid the development of safe and effective pain therapeutics: challenges and opportunities. *Nat Rev Neurol*. 2020;16(7):381–400. doi:10.1038/s41582-020-0362-2
58. Hernan MA, Robins JM. Using big data to emulate a target trial when a randomized trial is not available. *Am J Epidemiol*. 2016;183(8):758–764. doi:10.1093/aje/kwv254