

ORAL PRESENTATION

Open Access

Relationship between bone density and bone metabolism in adolescent idiopathic scoliosis (AIS)

Yoichi Aota*, Ko Ishida, Nobuyuki Tanabe, Tomoyuki Saito

From 10th International Conference on Conservative Management of Spinal Deformities - SOSORT 2013 Annual Meeting
Chicago, IL, USA. 8-11 May 2013

Purpose

Although osteopenia is often associated with AIS, studies on bone metabolism in relation to AIS have not yielded clear results. To characterize bone metabolism in AIS patients, a cross sectional study assessing bone metabolism and bone density was performed.

Methods

Using dual-energy X ray absorptiometry and bone metabolism markers (bone formation marker; BAP, bone resorption marker; TRAP5b), the bone mineral density of lumbar and bilateral proximal femurs were studied in 41 AIS patients aged 10 to 20 years old, with a mean of 15.2 ± 5.9 years old. Divided into two groups by levels of bone resorption marker (TRAP5b), BMD, BMI and age of menarche were compared statistically in each group.

Results

Among the AIS patients studied, osteopenia (-1 standard deviations to -2 standard deviations) was found in 34.1% of the patients and osteoporosis (below -2 standard deviations) was found in 24.4% of the patients. In 39 AIS patients (95.2%), BAP values were within normal range. On the other hand, TRAP5b values were significantly high in 65.9% of the patients. In high levels of the TRAP5b group, BMD values of the lumbar spine and right femoral neck were significantly lower than those of the TRAP5b group.

Conclusions and discussion

The bone resorption marker was high in 65.9% of the AIS patients, and high bone resorption in bone metabolism

was found to be a possible cause of low bone mineral density in patients with AIS.

Published: 18 September 2013

References

1. Cook SD, Harding AF, Morgan EI, et al: Trabecular bone mineral density in idiopathic scoliosis. *J Pediatr Orthop* 1987, **7**:168-174.
2. Hannon RA, Clowes JA, Eagleton AC, et al: Clinical performance of immunoreactive tartrate-resistant acid phosphatase isoform 5b as a marker of bone resorption. *Bone* 2004, **34**:187-194.
3. Cheung CS, Lee WT, Tse YK, et al: Generalized Osteopenia in Adolescent Idiopathic Scoliosis - Association With Abnormal Pubertal Growth, Bone Turnover, and Calcium Intake? *Spine* 2006, **31**:330-338.

doi:10.1186/1748-7161-8-S2-O4

Cite this article as: Aota et al.: Relationship between bone density and bone metabolism in adolescent idiopathic scoliosis (AIS). *Scoliosis* 2013 **8**(Suppl 2):O4.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: yaota@yokohama-cu.ac.jp
Department of Orthopaedic Surgery, Yokohama City University, Yokohama, Japan