

Scientific Article

# The long-lasting relationship of distress on radiation oncology-specific clinical outcomes



Justin Anderson MD, Alexander N. Slade MD, PhD\*,  
Philip Reed McDonagh MD, PhD, Whitney Burton MSW,  
Emma C. Fields MD

Virginia Commonwealth University Health System, Massey Cancer Center, Department of Radiation Oncology,  
Richmond, Virginia

Received 6 August 2018; revised 10 October 2018; accepted 2 November 2018

## Abstract

**Purpose:** The diagnosis and treatment of cancer can have significant mental health ramifications. The National Comprehensive Cancer Network currently recommends using a distress screening tool to screen patients for distress and facilitate referrals to social service resources. Its association with radiation oncology-specific clinical outcomes has remained relatively unexplored.

**Methods and materials:** With institutional review board approval, National Comprehensive Cancer Network distress scores were collected for patients presenting to our institution for external beam radiation therapy during a 1-year period from 2015 to 2016. The association between distress scores (and associated problem list items and process-related outcomes) and radiation oncology-related outcomes, including inpatient admissions during treatment, missed treatment appointments, duration of time between consultation and treatment, and weight loss during treatment, was considered.

**Results:** A total of 61 patients who received either definitive (49 patients) or palliative (12 patients) treatment at our institution and completed a screening questionnaire were included in this analysis. There was a significant association between an elevated distress score (7+) and having an admission during treatment (36% vs 11%;  $P = .04$ ). Among the patients treated with definitive intent, missing at least 1 appointment (71% vs 26%;  $P = .03$ ) and having an admission during treatment (57% vs 10%;  $P = .009$ ) were significantly associated with our institutional definition of elevated distress. We found no correlation between distress score and weight loss during treatment or a prolonged time between initial consult and treatment start.

**Conclusions:** High rates of distress are common for patients preparing to receive radiation therapy. These levels may affect treatment compliance and increase rates of hospital admissions. There remains equipoise in the best method to address distress in the oncology patient population. These results may raise awareness of the consequences of distress among radiation

Meeting information: A previous version of this paper was presented at the American Society for Radiation Oncology Meeting on October 1, 2017.

Sources of support: This work had no specific funding.

Conflicts of interest: The authors have no conflicts of interest to disclose.

\* Corresponding author. Virginia Commonwealth University, Massey Cancer Center, Department of Radiation Oncology, Box 980058, Richmond, VA 23298.

E-mail address: [alexander.n.slade@vcuhealth.org](mailto:alexander.n.slade@vcuhealth.org) (A.N. Slade).

<https://doi.org/10.1016/j.adro.2018.11.001>

2452-1094/© 2018 The Authors. Published by Elsevier Inc. on behalf of American Society for Radiation Oncology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

oncology patients. Specific interventions to improve distress need further study, but we suggest a more proactive approach by radiation oncologists in addressing distress.

© 2018 The Authors. Published by Elsevier Inc. on behalf of American Society for Radiation Oncology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

In 1997, the National Comprehensive Cancer Network (NCCN) established a multidisciplinary panel to examine how to integrate psychosocial care into routine cancer care.<sup>1</sup> Malignancies have long been known to disrupt a patient's family, friendships, finances, and work life, and these disruptions are often difficult to recognize and address by providers.<sup>2–4</sup> The NCCN Distress Thermometer is a tool that allows patients to score their level of distress on a scale of 0 to 10 (similar to the pain scale), with 0 being no distress and 10 being extreme distress. Physicians and others can use the tool to quickly identify patients with cancer who are experiencing distress and may benefit from intervention.

The tool includes a problem list on which patients can answer additional questions to help pinpoint the source of distress and allow for more targeted interventions. When the Distress Thermometer was originally released, a score of  $\geq 4$  was sufficient to trigger additional questioning and possible referral to psychosocial services.<sup>1</sup> This tool has provided a way for providers to quickly screen patients for distress and additional psychosocial concerns.

Approximately a third of patients with cancer are estimated to experience a significant degree of distress, which varies by cancer site.<sup>5</sup> High levels of distress on their own have been shown to be a poor prognostic factor,<sup>6,7</sup> but elevated distress levels may precipitate or exacerbate symptoms, such as loss of appetite, difficulty concentrating, and sleeplessness. These and other symptoms may undermine patients' ability to fight their own diseases.<sup>8</sup>

Some efforts have been made to correlate distress with specific cancer outcomes. One study found that high distress levels measured before and during radiation therapy (RT) prognostic with higher distress levels associated with decreased survival.<sup>9</sup>

This retrospective study explores the association between distress scores and radiation oncology-specific outcomes. A secondary goal was to determine what factors could be contributing to worse outcomes in patients with high distress.

## Methods and Materials

With institutional review board approval, we performed a retrospective review of our distress screening questionnaire and problem list (variation of the NCCN

version; Fig 1) collected from 129 patients receiving treatment at the Department of Radiation Oncology when they presented for their initial consultation between 2015 and 2016. Of these patients, 38 questionnaires had insufficient information to be included in the analysis (ie, did not report a distress score). An additional 30 patients were excluded because they did not ultimately receive RT at our institution (ie, it was received elsewhere or refused), were treated with brachytherapy as monotherapy (eg, for prostate cancer), or were treated for a benign condition (eg, Dupuytren's contracture). Of the 61 remaining patients, 49 were treated with definitive intent and 12 were treated with palliative (ie, noncurative) intent.

We used Fisher's exact test and *t* tests to evaluate the association between distress score and radiation oncology-specific outcomes, including inpatient admissions during treatment, missed or canceled treatment appointments, duration of time between consultation and treatment, and weight loss during treatment. The policy established at our institution specifies that patients who report distress scores of  $\geq 7$  are considered high distress and trigger a social work consult. Therefore, we divided patients into 2 categories: low distress with scores of  $\leq 6$  and high distress with scores of 7 to 10.

In addition to the ordinal 0-to-10 score, the NCCN scale also includes a problem list, which offers patients a yes/no response on 39 potential items that may have been problematic for the patient during the previous week. These items include practical problems, including child care and housing; family problems; emotional problems, such as depression; spiritual problems; and physical problems. Our institution makes minor modifications to this problem list (Fig 2). The overall number of reported problems was totaled according to their subsection of the problem list (ie, spiritual, family, practical, and physical). The breakdown of specific questions included in the categories is presented in Figure 2. Patients were given the option to check yes or no for each problem, but if a patient completed any of the problem list questions, missing answers were assumed to be no.

## Results

Of the 61 patients included, 44% were female and 72% were white. The most common malignancies were head

Distress is a term used to describe unhappiness and unpleasant feelings that make it difficult to deal with what is going on. Filling out this form helps your health care team know how you are doing and how we can help.


Keep in mind that levels of distress may change over time and patients are welcome to reach out to anyone on their health care team to request assistance from the Clinical Social Worker.

#### Step One

On a scale from 1 – 10, how much distress have you felt over the last week including today? Circle the number on the distress thermometer to show how you are feeling.

This tool is used only to help you find needed assistance and resources.

**Please note:** Depending on your level of distress, you may be contacted by a social worker to make sure you are aware of the services available. **This call will be made whether you have a cancer diagnosis or not.** We help all of the patients that come through our clinics regardless of the diagnosis. **Please continue to Step Two on the back of this page.**



**Figure 1** Distress thermometer for patients. (adapted from the National Comprehensive Cancer Network)

and neck (31%), breast (20%), and genitourinary (15%), with all stages of disease represented (Table 1).

In our sample, patients who reported higher distress scores were more likely to present with higher stages of disease ( $P = .018$ ). There was a significant association between a high distress score and admission during treatment ( $P = .041$ ). Other demographic and clinical factors, including age, race, and cancer stage or site, were not correlated with distress levels in our sample. Likewise, no correlation was detected between distress score and amount of weight loss/gain, or the time between initial consult and start of treatment.

Table 2 explores the categories of problems that may underlie higher levels of distress. Overall, the average distress score of our sample was 3.8. Patients with higher levels of distress reported significantly more practical, emotional, family-related, and physical problems. Practical problems showed the largest disparity based on distress level, with the reported number of problems averaging 2.4 among those reporting distress scores of 7 to 10 versus 0.8 for those with distress scores of 0 to 6 ( $P = .003$ ). The positive association between distress levels and these outcomes was robust to other cutoffs of distress scores (eg,  $>5$ ). A multivariate analysis to adjust for the variables noted in Table 1 was considered, but the small sample size (especially for the clinical variables such as cancer site) would make these results less reliable.

Table 3 explores the breakdown for patients who missed or canceled any of their treatment appointments and those who experienced hospital admissions during their RT course. Overall, patients who canceled or missed RT appointments had, on average, nearly double the reported distress score compared with those who did not have any such missed or canceled appointments (4.9

vs 3.2;  $P = .047$ ). The number of emotional problems and the number of physical problems were both associated with canceling or missing a treatment appointment. However, the remaining items of the problem list were not associated with the probability of a patient canceling or missing a treatment appointment. Patients who had a hospital admission during RT also reported elevated distress scores (3.4 vs 5.8;  $P = .024$ ), but they were not more likely to report an increased number of practical, emotional, family, or physical problems.

When evaluating patients treated with definitive intent, a higher proportion of patients with higher distress scores received a social work consult (71% vs 26%;  $P = .030$ ). When palliative and definitively treated patients were combined, there was no longer a significant difference in level of distress and likelihood of receiving a social work consult (50% vs 28%;  $P = .19$ ). Overall, however, patients treated with palliative intent had significantly higher distress scores relative to those treated with definitive intent (5.9 vs 3.2;  $P = .007$ ), and there was a weak association between palliative treatment intent and number of practical and physical problems ( $P < .10$ ).

## Discussion

This study identifies several practical issues surrounding the care of radiation oncology patients that may be more prevalent among patients with higher levels of distress at the time of initial consultation. Specifically, we find that higher levels of distress correlated with more missed appointments and more inpatient admissions during RT, especially among patients who received RT

Please check YES or NO for on the list below.

<u>My Personal Concerns:</u>		<u>My Physical Concerns:</u>	
YES	NO	YES	NO
<input type="checkbox"/>	<input type="checkbox"/> (Pr) I have housing concerns	<input type="checkbox"/>	<input type="checkbox"/> Appearance
<input type="checkbox"/>	<input type="checkbox"/> (Pr) I have insurance/financial issues	<input type="checkbox"/>	<input type="checkbox"/> Bathing/Dressing
<input type="checkbox"/>	<input type="checkbox"/> (Pr) I have legal issues*	<input type="checkbox"/>	<input type="checkbox"/> Breathing
<input type="checkbox"/>	<input type="checkbox"/> (Pr) I have transportation issues	<input type="checkbox"/>	<input type="checkbox"/> Changes in urination
<input type="checkbox"/>	<input type="checkbox"/> (Pr) I have work/school issues	<input type="checkbox"/>	<input type="checkbox"/> Constipation
<input type="checkbox"/>	<input type="checkbox"/> (Em) I feel depressed	<input type="checkbox"/>	<input type="checkbox"/> Diarrhea
<input type="checkbox"/>	<input type="checkbox"/> (Em) I feel fear or worry	<input type="checkbox"/>	<input type="checkbox"/> Drowsiness
<input type="checkbox"/>	<input type="checkbox"/> (Em) I feel sadness	<input type="checkbox"/>	<input type="checkbox"/> Eating/appetite
<input type="checkbox"/>	<input type="checkbox"/> (Em) I have a loss of interest in usual activities	<input type="checkbox"/>	<input type="checkbox"/> Fatigue/tiredness
<input type="checkbox"/>	<input type="checkbox"/> (Fa) I have changes in intimacy	<input type="checkbox"/>	<input type="checkbox"/> Feeling Swollen
<input type="checkbox"/>	<input type="checkbox"/> (Fa) I have concerns about fertility	<input type="checkbox"/>	<input type="checkbox"/> Fevers
<input type="checkbox"/>	<input type="checkbox"/> (Sp) I have spiritual/religious concerns	<input type="checkbox"/>	<input type="checkbox"/> Getting around
<input type="checkbox"/>	<input type="checkbox"/> (Fa) I have family problems	<input type="checkbox"/>	<input type="checkbox"/> Indigestion
<input type="checkbox"/>	<input type="checkbox"/> (Pr) I am struggling with a treatment decision	<input type="checkbox"/>	<input type="checkbox"/> Memory/concentration
		<input type="checkbox"/>	<input type="checkbox"/> Mouth Sores
		<input type="checkbox"/>	<input type="checkbox"/> Nausea
		<input type="checkbox"/>	<input type="checkbox"/> Nose dry/congestion
		<input type="checkbox"/>	<input type="checkbox"/> Pain
		<input type="checkbox"/>	<input type="checkbox"/> Sexual
		<input type="checkbox"/>	<input type="checkbox"/> Skin dry/itchy
		<input type="checkbox"/>	<input type="checkbox"/> Sleeping Issues
		<input type="checkbox"/>	<input type="checkbox"/> Substance Abuse
		<input type="checkbox"/>	<input type="checkbox"/> Tingling in hands/feet

**Figure 2** Distress thermometer for patients (adapted from the National Comprehensive Cancer Network). Abbreviations: Em = emotional problem; Fa = family problem; Pr = practical problem; Sp = spiritual problem. All problems in the right column are classified as physical problems. \*Problem list item used at our institution, but not in the National Comprehensive Cancer Network tool.

with definitive intent. Missed treatment appointments and inpatient admissions during treatment can prolong the duration of RT and possibly decrease the efficacy of treatment. For example, among head and neck cancer patients, each additional week of treatment prolongation was associated with a 14% reduction in local control.<sup>10</sup> Similar findings of worse outcomes with treatment prolongation have been reported in prostate, cervical, lung, anal, bladder, and breast cancers.<sup>11–16</sup> Increased rates of hospitalization for patients with higher levels of distress may also signify additional health problems that need to be addressed more thoroughly in these patients.

To prevent hospitalizations and missed appointments among patients with higher levels of distress, distress must be detected early and addressed by a multidisciplinary care team. We found that social work was frequently involved for patients who reported high distress, though social work involvement on its own was not associated with our targeted outcomes. Addressing distress within the radiation oncology setting is complicated, and barriers to proper treatment include lack of training for clinicians and staff in the management of psychosocial issues, time constraints, and limited patient reporting.

The results of several studies have been inconclusive on effective methods to improve patient-reported distress. One previously proposed method involved caregivers having patients rank the top 4 issues that contribute to their distress and giving the patients options for how they would like to address them, including outside referral.<sup>17</sup> Unfortunately, this method of addressing distress was shown to be cost ineffective and was not associated with improvement in patient mood states. There has been some success in nursing staff helping to address patient distress. In a study in which nurses received a training course before providing basic psychosocial care, minor interventions, and exploring referral possibilities for their patients, patients did report satisfaction with the nurses' care, but no effect on depressive symptoms or quality of life was seen.<sup>18</sup> A recent randomized controlled trial that evaluated brief psychosocial intervention for depressed patients with cancer concluded that it was insufficient as a stand-alone treatment.<sup>19</sup>

These studies show that more intensive and focused treatment is needed to help address patient distress. Sometimes this requires the involvement of another service that is equipped to address distress, such as psychiatry. Unfortunately, few patients desire referrals to other

**Table 1** Demographic, clinical, and process-related characteristics of patients by distress level and treatment intent

	Patients treated with both palliative and definitive intent				Patients treated with definitive intent			
	All patients	Distress score 0-6	Distress score 7-10	<i>P</i> -value	Definitive patients	Distress score 0-6	Distress score 7-10	<i>P</i> -value
Age, y	60.0 (14.0)	59.9 (13.6)	60.3 (15.7)	.94	58.7 (14.4)	58.1 (18.6)	58.8 (13.9)	.91
Female	27 (44%)	21 (45%)	6 (43%)	1	21 (43%)	2 (29%)	19 (45%)	.68
Race								
African American	17 (28%)	14 (30%)	3 (21%)	.54	13 (27%)	1 (14%)	12 (29%)	.41
White	44 (72%)	33 (70%)	11 (79%)		35 (73%)	6 (86%)	29 (71%)	
Stage								
I*	7 (11%)	6 (13%)	1 (7%)	.018	6 (13%)	1 (14%)	5 (13%)	.24
II	15 (25%)	15 (32%)	0 (0%)		14 (30%)	0 (0%)	14 (35%)	
III	15 (25%)	12 (26%)	3 (21%)		13 (28%)	3 (43%)	10 (25%)	
IV	24 (39%)	14 (30%)	10 (71%)		14 (30%)	3 (43%)	11 (28%)	
Cancer site								
Breast	12 (20%)	11 (23%)	1 (7%)	.43	11 (22%)	0 (0%)	11 (26%)	.18
Central nervous system	1 (2%)	1 (2%)	0 (0%)		1 (2%)	0 (0%)	1 (2%)	
Gastrointestinal	6 (10%)	4 (9%)	2 (14%)		5 (10%)	1 (14%)	4 (10%)	
Genitourinary	9 (15%)	8 (17%)	1 (7%)		7 (14%)	0 (0%)	7 (17%)	
Gynecologic	2 (3%)	1 (2%)	1 (7%)		1 (2%)	0 (0%)	1 (2%)	
Head and neck	19 (31%)	14 (30%)	5 (36%)		19 (39%)	5 (71%)	14 (33%)	
Hematologic	5 (8%)	4 (9%)	1 (7%)		2 (4%)	0 (0%)	2 (5%)	
Sarcoma	1 (2%)	0 (0%)	1 (7%)		1 (2%)	1 (14%)	0 (0%)	
Skin	2 (3%)	1 (2%)	1 (7%)		0 (0%)	0 (0%)	0 (0%)	
Thorax	4 (7%)	3 (6%)	1 (7%)		2 (4%)	0 (0%)	2 (5%)	
Treatment with palliative intent	12 (20%)	5 (11%)	7 (50%)	.003				
Process outcomes								
Days to start radiation therapy from consult	50.7 (51.6)	55.8 (55.6)	33.7 (30.9)	.16	58.6 (54.1)	53.0 (33.4)	59.5 (57.1)	.77
Change in weight (kg)	−2.6 (6.0)	−2.2 (5.3)	−4.5 (9.5)	.37	−2.6 (6.0)	−4.5 (9.5)	−2.2 (5.3)	.37
Any canceled treatments	19 (31%)	12 (26%)	7 (50%)	.11	16 (33%)	5 (71%)	11 (26%)	.030
Social work consultation	20 (33%)	13 (28%)	7 (50%)	.19	16 (33%)	5 (71%)	11 (26%)	.030
Admission during radiation therapy	10 (16%)	5 (11%)	5 (36%)	.041	8 (16%)	4 (57%)	4 (10%)	.009
Number of patients	61	47	14		49	42	7	

\* Includes one patient with Stage 0 breast cancer (ductal carcinoma in situ).

professionals, including <15% of patients who report high distress (score >5), and older patients are even less likely to desire referrals.<sup>20–22</sup> Therefore, focusing on more training for radiation oncology professionals to address patients' psychological needs seems prudent, but this is made more difficult by the stigma that surrounds psychosocial issues for both health care providers and patients, especially in the oncologic setting.<sup>23</sup> Oncologists have been reported to be reluctant to address psychosocial issues in patients for fear of causing pain, harm, or additional distress, and one study showed that as many as 73% of depression cases are underrecognized by oncologists.<sup>24,25</sup>

For radiation oncologists, recognizing a patient's psychological response to the stress of a cancer diagnosis and RT is important. These patients commonly use defense mechanisms, including denial and projection, which are important to identify because they can act as further barriers to understanding and treating patient distress.<sup>26</sup> In

the radiation oncology setting, addressing a patient's distress may be as simple as assuaging fears and misconceptions of RT, including concerns of becoming radioactive or impact on fertility.<sup>27</sup> Understanding racial disparities in factors that drive distress, (eg, practical problems such as transportation) may also help identify mechanisms to alleviate distress.<sup>28</sup> Simple training in communication skills and recognizing empathetic opportunities with a focus on behaviors that express support, sympathy, and compassion, can be effective in improving diagnosis and treatment of psychological issues.<sup>29–31</sup>

Increasing protected time to focus on patient distress is another key element.<sup>32</sup> A guideline written to help psychiatric providers care for radiation oncology patients may also provide useful resources for radiation oncologists.<sup>33</sup> This resource should discuss common causes of anxiety in these patients as well as medical and psychiatric medical techniques for management. Another article reviewed psychosocial interventions, including

**Table 2** Breakdown of problem list responses by distress level

	All patients	Distress score 0-6	Distress score 7-10	P-value
Overall distress score (0-10)	3.8 (3.2)	2.4 (2.1)	8.3 (1.2)	<.001
Number of practical problems	0.6 (0.8)	0.5 (0.7)	1.1 (0.8)	.013
Number of emotional problems	1.0 (1.4)	0.7 (1.2)	2.1 (1.6)	<.001
Number of family problems	0.3 (0.7)	0.1 (0.4)	0.8 (1.1)	.002
Number of physical problems	3.4 (4.6)	2.3 (3.4)	6.9 (6.2)	<.001
Spiritual/religious problems	4 (7%)	2 (4%)	2 (14%)	.22
Number of patients	61	47	14	

psychoeducation, cognitive-behavior therapy, and supportive-expressive therapy, that can be done in the RT clinic by training radiation oncologists and staff to be conductors.<sup>34</sup> In this randomized controlled trial, these interventions were effective in reducing depression and anxiety in patients, although they did not reduce the risk of cancer recurrence or death.

Along with focusing on improving distress management in the RT clinic, it is equally important for radiation oncologists to recognize when patients need a referral to outside psychiatric services, especially in the setting of time constraints in the clinic that do not allow for more intensive care. Very elevated distress scores should raise concerns for the need to refer to outside services, with studies showing that 55% of patients with distress scores of 10 desire a referral. Patients with higher scores who did not desire a referral were more likely reluctant because of fears of negative impacts of psychosocial intervention rather than a true lack of need of support.<sup>20,24</sup> Even with this reluctance, most patients were found to highly value oncologists' opinions of support services, and the majority were willing to accept a referral when offered.<sup>35,36</sup> Radiation oncologists should make outside providers aware of the special needs and issues of their patients undergoing RT and refer them to the aforementioned psychiatric guideline for radiation oncology patients.<sup>33</sup> Outside referral has been shown to be effective, with outpatient psychosocial interventions improving outcomes in patients with cancer.<sup>37,38</sup> The benefits of

outpatient psychotherapy have also been shown in the radiation oncology patient population.<sup>39,40</sup>

Although our study found associations between elevated distress and missed RT appointments and hospitalizations, our analysis has several limitations. First, we looked at a combination of patients who were treated with definitive intent and palliative intent. When we analyzed only patients treated with definitive intent alone, we found a stronger association between distress scores and the likelihood of missing treatment appointments or being admitted to the hospital during treatment. When we analyzed our combined group of both palliatively and definitively treated patients, the likelihood of missing a treatment appointment or being admitted during treatment decreased, especially in our high-distress cohort. This could be attributable to the shorter treatment times of palliative treatment with fewer opportunities to miss an appointment or be admitted during treatment. Our analysis uses a relatively small sample size from a single institution, and a larger group of patients would provide better information about the association distress scores have with our targeted outcomes.

Furthermore, our institution is relatively unique in its role as a safety net hospital, and many patients receiving RT at our institution may have different factors driving their distress compared with the overall oncologic population at large. Along those lines, unobserved factors, such as socioeconomic status (education/income), may explain a portion of the effect of distress on these

**Table 3** Overall distress and problem list responses by missed treatment, hospital admission and palliative treatment intent

Variable	Any canceled or missed treatments			Any hospital admissions during treatment			Palliative treatment intent		
	No	Yes	P-value	No	Yes	P-value	No	Yes	P-value
Overall distress score (0-10)	3.2 (3.1)	4.9 (3.1)	.047	3.4 (3.0)	5.8 (3.5)	.024	3.2 (2.9)	5.9 (3.4)	.007
Number of practical problems	0.5 (0.7)	0.8 (0.9)	.16	0.6 (0.8)	0.8 (0.6)	.47	0.6 (0.7)	1.0 (0.9)	.064
Number of emotional problems	0.8 (1.2)	1.6 (1.6)	.027	0.9 (1.3)	1.6 (1.7)	.17	1.0 (1.4)	1.3 (1.4)	.44
Number of family problems	0.2 (0.7)	0.5 (0.7)	.18	0.3 (0.7)	0.4 (0.7)	.6	0.2 (0.5)	0.6 (1.2)	.11
Number of physical problems	2.4 (4.2)	5.5 (4.9)	.015	3.2 (4.8)	4.3 (3.1)	.49	2.8 (3.9)	5.7 (6.6)	.054
Spiritual/religious problems	1 (2%)	3 (16%)	.085	3 (6%)	1 (10%)	.52	2 (4%)	2 (17%)	.17
Social work consultation	11 (26%)	9 (47%)	.14	14 (27%)	6 (60%)	.066	16 (33%)	4 (33%)	1.00
Number of patients	42	19		51	10		49	12	

outcomes, although these patient characteristics are not typically available to clinicians. At our institution, there is a very large referral base from which head and neck cancer cases are captured, which skews our analysis somewhat to that treatment site compared with an average community radiation clinic, which may have less representation of patients with head and neck cancer.

Additionally, as discussed, our distress measures are collected at a single point in time before simulation or treatment. This single snapshot of patient distress does not give a complete picture of distress across the patients' treatment course compared with more elaborate batteries conducted longitudinally. Finally, multiple patients who were offered a screening questionnaire did not complete it, which makes our analytical sample subject to bias related to selection.

## Conclusions

Elevated distress continues to be a difficult problem to handle for many patients with cancer and for providers. Our study provides evidence that high levels of distress can affect radiation oncology—specific outcomes, leading to missed treatments, which may affect long term outcomes including locoregional control and survival. Overall, our findings underscore the need for radiation oncologists to be proactive in addressing patient distress to help decrease potential barriers to receiving treatment. Radiation oncologists may also need to involve several members of the health care team, such as social work, psychiatry, and other medicine subspecialties, early in the course of treatment to quickly and effectively address sources of distress and additional underlying health problems. However, the benefits of these strategies lack extensive evidence, and more attention and research are needed to develop guidelines on the best methods to reduce and manage patient distress and improve outcomes.

## References

- Holland JC, Bultz BD. National Comprehensive Cancer Network (NCCN). The NCCN guideline for distress management: A case for making distress the sixth vital sign. *J Natl Compr Cancer Netw*. 2007;5:3-7.
- Weisman AD, Worden JW. The existential plight in cancer: Significance of the first 100 days. *Int J Psychiatry Med*. 1977;7:1-15.
- Cassileth BR, Lusk EJ, Brown LL, Cross PA, Walsh WP, Hurwitz S. Factors associated with psychological distress in cancer patients. *Med Pediatr Oncol*. 1986;14:251-254.
- Söllner W, DeVries A, Steixner E, et al. How successful are oncologists in identifying patient distress, perceived social support, and need for psychosocial counselling? *Br J Cancer*. 2001;84:179-185.
- Zabora J, BrintzenhofeSzoc K, Curbow B, Hooker C, Piantadosi S. The prevalence of psychological distress by cancer site. *Psychooncology*. 2001;10:19-28.
- Gilbert MR, Dignam JJ, Armstrong TS, et al. A randomized trial of bevacizumab for newly diagnosed glioblastoma. *N Engl J Med*. 2014;370:699-708.
- Kaasa S, Mastekaasa A, Lund E. Prognostic factors for patients with inoperable non-small cell lung cancer, limited disease. *Radiother Oncol*. 1989;15:235-242.
- Carlson LE, Angen M, Cullum J, et al. High levels of untreated distress and fatigue in cancer patients. *Br J Cancer*. 2004;90:2297-2304.
- Habboush Y, Shannon RP, Niazi SK, et al. Patient-reported distress and survival among patients receiving definitive radiation therapy. *Adv Radiat Oncol*. 2017;2:211-219.
- Fowler JF, Lindstrom MJ. Loss of local control with prolongation in radiotherapy. *Int J Radiat Oncol Biol Phys*. 1992;23:457-467.
- Thames HD, Kuban D, Levy LB, et al. The role of overall treatment time in the outcome of radiotherapy of prostate cancer: An analysis of biochemical failure in 4839 men treated between 1987 and 1995. *Radiother Oncol*. 2010;96:6-12.
- Perez CA, Grigsby PW, Castro-Vita H, Lockett MA. Carcinoma of the uterine cervix. Impact of prolongation of overall treatment time and timing of brachytherapy on outcome of radiation therapy. *Int J Radiat Oncol Biol Phys*. 1995;32:1275-1288.
- Cox JD, Pajak TF, Asbell S, et al. Interruptions of high-dose radiation therapy decrease long-term survival of favorable patients with unresectable non-small cell carcinoma of the lung: Analysis of 1244 cases from 3 Radiation Therapy Oncology Group (RTOG) trials. *Int J Radiat Oncol Biol Phys*. 1993;27:493-498.
- Weber DC, Kurtz JM, Allal AS. The impact of gap duration on local control in anal canal carcinoma treated by split-course radiotherapy and concomitant chemotherapy. *Int J Radiat Oncol Biol Phys*. 2001;50:675-680.
- Maciejewski B, Majewski S. Dose fractionation and tumour repopulation in radiotherapy for bladder cancer. *Radiother Oncol*. 1991;21:163-170.
- Bese NS, Sut PA, Ober A. The effect of treatment interruptions in the postoperative irradiation of breast cancer. *Oncology*. 2005;69:214-223.
- Hollingworth W, Metcalfe C, Mancero S, et al. Are needs assessments cost effective in reducing distress among patients with cancer? A randomized controlled trial using the Distress Thermometer and Problem List. *J Clin Oncol*. 2013;31:3631-3638.
- van der Meulen IC, May AM, Koole R, Ros WJG. A distress thermometer intervention for patients with head and neck cancer. *Oncol Nurs Forum*. 2018;45:E14-E32.
- Turner J, Kelly B, Clarke D, et al. A tiered multidisciplinary approach to the psychosocial care of adult cancer patients integrated into routine care: The PROMPT study (a cluster-randomised controlled trial). *Support Care Cancer*. 2017;25:17-26.
- Graves KD, Arnold SM, Love CL, Kirsh KL, Moore PG, Passik SD. Distress screening in a multidisciplinary lung cancer clinic: Prevalence and predictors of clinically significant distress. *Lung Cancer*. 2007;55:215-224.
- Lynch J, Goodhart F, Saunders Y, O'Connor SJ. Screening for psychological distress in patients with lung cancer: Results of a clinical audit evaluating the use of the patient Distress Thermometer. *Support Care Cancer*. 2011;19:193-202.
- Tuinman MA, Gazendam-Donofrio SM, Hoekstra-Weebers JE. Screening and referral for psychosocial distress in oncologic practice: Use of the Distress Thermometer. *Cancer*. 2008;113:870-878.
- Dilworth S, Higgins I, Parker V, Kelly B, Turner J. Patient and health professional's perceived barriers to the delivery of psychosocial care to adults with cancer: A systematic review. *Psychooncology*. 2014;23:601-612.
- Okuyama T, Endo C, Seto T, et al. Cancer patients' reluctance to disclose their emotional distress to their physicians: A study of Japanese patients with lung cancer. *Psychooncology*. 2008;17:460-465.

25. Turner J, Zapart S, Pedersen K, et al. Clinical practice guidelines for the psychosocial care of adults with cancer. *Psychooncology*. 2005;14:159-173.
26. Porcerelli JH, Cramer P, Porcerelli DJ, Arterbery VE. Defense mechanisms and utilization in cancer patients undergoing radiation therapy: A pilot study. *J Nerv Ment Dis*. 2017;205:466-470.
27. Gillan C, Abrams D, Harnett N, Wiljer D, Catton P. Fears and misperceptions of radiation therapy: Sources and impact on decision-making and anxiety. *J Cancer Educ*. 2014;29:289-295.
28. McDonagh PR, Slade AN, Anderson J, Burton W, Fields EC. Racial differences in responses to the NCCN Distress Thermometer and Problem List: Evidence from a radiation oncology clinic. *Psychooncology*. 2018;27:2513-2516.
29. Fallowfield L, Ratcliffe D, Jenkins V, Saul J. Psychiatric morbidity and its recognition by doctors in patients with cancer. *Br J Cancer*. 2001;84:1011-1015.
30. Fogarty LA, Curbow BA, Wingard JR, McDonnell K, Somerfield MR. Can 40 seconds of compassion reduce patient anxiety? *J Clin Oncol*. 1999;17:371-379.
31. Pollak KI, Arnold RM, Jeffreys AS, et al. Oncologist communication about emotion during visits with patients with advanced cancer. *J Clin Oncol*. 2007;25:5748-5752.
32. Biddle L, Paramasivan S, Harris S, Campbell R, Brennan J, Hollingworth W. Patients' and clinicians' experiences of holistic needs assessment using a cancer distress thermometer and problem list: A qualitative study. *Eur J Oncol Nurs*. 2016;23:59-65.
33. Holmes EG, Holmes JA, Park EM. Psychiatric care of the radiation oncology patient. *Psychosomatics*. 2017;58:457-465.
34. Guo Z, Tang HY, Li H, et al. The benefits of psychosocial interventions for cancer patients undergoing radiotherapy. *Health Qual Life Outcomes*. 2013;11:121.
35. Edgar L, Remmer J, Rosberger Z, Fournier MA. Resource use in women completing treatment for breast cancer. *Psychooncology*. 2000;9:428-438.
36. Roth AJ, Kornblith AB, Batel-Copel L, Peabody E, Scher HI, Holland JC. Rapid screening for psychologic distress in men with prostate carcinoma: A pilot study. *Cancer*. 1998;82:1904-1908.
37. Andersen BL, Yang HC, Farrar WB, et al. Psychologic intervention improves survival for breast cancer patients. *Cancer*. 2008;113:3450-3458.
38. Giese-Davis J, Collie K, Rancourt KM, Neri E, Kraemer HC, Spiegel D. Decrease in depression symptoms is associated with longer survival in patients with metastatic breast cancer: A secondary analysis. *J Clin Oncol*. 2011;29:413-420.
39. Hess CB, Chen AM. Measuring psychosocial functioning in the radiation oncology clinic: A systematic review. *Psychooncology*. 2014;23:841-854.
40. Schnur JB, David D, Kangas M, Green S, Bovbjerg DH, Montgomery GH. A randomized trial of a cognitive-behavioral therapy and hypnosis intervention on positive and negative affect during breast cancer radiotherapy. *J Clin Psychol*. 2009;65:443-455.