



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Letter

Repeated vaccination against SARS-CoV-2 elicits robust polyfunctional T cell response in allogeneic stem cell transplantation recipients

Patrick Harrington,^{1,2} Katie J. Doores,³ Chandan Saha,¹ Jamie Saunders,¹ Fiona Child,¹ Richard Dillon,^{1,4} Sukran Saglam,¹ Kavita Raj,¹ Donal McLornan,¹ Daniele Avenoso,⁵ Shahram Kordasti,^{1,2} Amy O'Reilly,¹ Andreas Espehana,¹ Thomas Lechmere,³ Hataf Khan,³ Michael H. Malim,³ Claire Harrison,^{1,2} Varun Mehra,⁵ and Hugues de Lavallade^{1,2,*}

¹Department of Clinical Haematology, Guy's & St. Thomas' NHS Foundation Trust, London, UK

²School of Cancer and Pharmaceutical Science, King's College London, London, UK

³Department of Infectious Diseases, School of Immunology & Microbial Sciences, King's College London, London, UK

⁴Department of Medicine & Molecular Genetics, King's College London, London, UK

⁵Department of Haematological Medicine, King's College Hospital, London, UK

*Correspondence: h.delavallade@nhs.net

<https://doi.org/10.1016/j.ccell.2021.10.002>

SARS-CoV-2 has led to unprecedented global healthcare challenges, with poor outcomes observed in groups with immune deficiency, including allogeneic stem cell transplantation (allo-SCT) recipients (Bakouny et al., 2020). T cell and B cell responses following vaccination against SARS-CoV-2 are important in reducing the risk of severe COVID-19, but the T cell response has not been extensively investigated in this population. We designed a prospective study to evaluate response to vaccination in patients with hematologic malignancies. Herein we report analysis of T cell and humoral response to sequential dosing of vaccination against SARS-CoV-2 in allo-SCT recipients.

Anti-SARS-CoV-2 Spike protein (S) IgG ELISA and neutralizing antibody testing were performed as described previously. The induction of virus-specific T cell responses by vaccination was assessed by flow-cytometric enumeration of antigen-specific CD8⁺ and CD4⁺ T lymphocytes using an intracellular cytokine assay for IFN γ and TNF α .

A total of 23 patients were analyzed at one or more time point around the two-dose vaccination schedule (Table S1). Median age was 55 years (range 25–74), and 69.6% (16) were male. Median time from allo-SCT was 55 months (19–172), and BNT162b2 vaccine was given to 81% (21) of patients, while others received ChAdOx1-S.

Following a first dose of vaccine, an anti-S IgG response was assessed in 18 patients at a median of 4.2 weeks after

vaccination. Anti-S IgG was detectable in only 38.9% (7), with 4 of these having weak positive results (Figure S1A). A mean anti-S IgG EC₅₀ of 76 (range 0–526) was observed at this time point (Figure S1B). Neutralizing antibody analysis was performed in all 7 patients with detectable anti-S IgG at this time point, with a mean ID50 of 292 observed (32–968) (Figure S1C).

Antibody testing was performed in 16 patients following two doses of vaccine, at a median of 12 weeks after the second dose. A detectable anti-S IgG was observed in 81% (13) of patients ($p \leq 0.017$) (Figure S1A), with a mean anti-S IgG of 1043 (0–5594) ($p = 0.025$) (Figure S1B). Neutralizing antibody testing performed in 13 patients with detectable IgG showed a mean ID50 of 747 (107–4707) (Figure S1C). After two doses of vaccine, antibody testing was performed in 10 patients with chronic graft-versus-host disease (GvHD) receiving extracorporeal photopheresis (ECP) and 6 patients not receiving ECP, with a mean EC₅₀ of 574 in ECP group, compared with 1826 non-ECP ($p = 0.17$). Similarly, mean neutralizing antibody ID50 was 312 in those requiring ECP compared with 719 in non-ECP. There was a significant correlation between anti-S IgG level and neutralizing ability from paired samples, with r value of 0.83 ($p < 0.0001$) (Figure S1D).

T cell analysis was performed in 17 patients after a single dose of vaccine and in 17 patients after two doses. A T cell response was observed in 35.3% (6)

of patients after one dose and in 82.3% (14) of patients after two doses ($p = 0.013$) (Figure S1E). A CD4⁺ T cell response was observed in 29.4% (5) of patients after one dose and 70.6% (12) of patients after two doses ($p = 0.038$), while a CD8⁺ T cell response was only seen in 17.6% (3) after one dose but 52.3% (9) after two doses ($p = 0.07$). Mean CD4⁺/CD8⁺ TNF α expression after a single dose was 0.12%/0.04%, which increased to 0.42%/0.13% after second dose ($p = 0.17/0.3$). Similarly, mean CD4⁺/CD8⁺ IFN γ expression after a single dose was 0.06%/0.03%, which again increased to 0.07%/0.17% ($p = 0.8/0.1$).

A polyfunctional T cell response, with dual expression of more than one proinflammatory cytokine within the same cell, was observed in 29.4% (5) of patients after one dose and 70.6% (12) after two doses ($p = 0.038$) (Figures S1F and S1G). After a single dose, mean CD4⁺ polyfunctional T cell response was 0.009%, with an increase to 0.026% after 2 doses ($p = 0.068$) (Figure S1H). Consistently, more than 90% of reactive T cells expressing pro-inflammatory cytokines showed co-expression of CD45RO, a surface protein marker for memory T cells. After a second dose, patients with chronic GvHD requiring ECP had a mean CD4⁺ TNF α expression of 0.18% compared with 0.86% in those not requiring ECP ($p = 0.09$) (Figure S1I).

Patients with prior allo-SCT who contract COVID-19 infection have poor outcomes, with overall survival reported at 68% at 30 days post diagnosis (Sharma



et al., 2021). Therefore, the development of immunity is particularly important in this patient group. We have previously reported that a single dose of BNT162b2 is sufficient to generate both a humoral and a T cell response in most patients with chronic myeloid malignancies (Harrington et al., 2021). This is in contrast to the response observed in many cancer-patient groups, particularly those with lymphoid malignancies who have received anti-CD20 targeted therapy (Addeo et al., 2021, Greenberger et al., 2021, Thakkar et al., 2021). We demonstrate here how a second dose is required for a significant increase in seroconversion rates and detectable memory T cells in allo-SCT recipients. Through analysis of samples at consecutive time points, including sequential samples from the same patients, we were able to observe the longitudinal response to vaccination and show that a second dose is required for adequate immunogenicity in this population. Our findings are in keeping with that from two studies on isolated antibody responses in allo-SCT patients which reported an anti-S IgG response after a second injection in 83% and 78% of participants, respectively (Redjoul et al., 2021, Le Bourgeois et al., 2021).

Our data report the T cell response to SARS-CoV-2 vaccination in patients with previous allo-SCT. Despite a poor T cell response after a first vaccine injection, a second dose elicited anti-S reactive T cells in most patients. Moreover, a polyfunctional T cell response was also elicited by a second dose, which may have particular functional relevance with regards to anti-viral immunity, with these cells recognized as providing a more effective anti-viral response in the context of COVID-19 infection (Peng et al., 2020). A memory T cell response may play a particularly important role in providing immunity to COVID-19, as studies have shown significant decline in antibody levels in the general population at 3 months post natural infection (Seow et al., 2020).

We have also focused our analysis on patients considered to be particularly

immune suppressed with regards to chronic GvHD and ongoing systemic immune suppression. While these patients did show a reduced T cell and antibody response when compared with patients off immune suppression, this was not significant, and most showed an adequate neutralizing antibody response after a second injection. Our study is, however, limited by small sample size, and further longitudinal data are required to evaluate whether the response generated is adequate to provide anti-viral protection.

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.ccell.2021.10.002>.

ACKNOWLEDGMENTS

The authors acknowledge a Blood Cancer UK award to P.H. and H.d.L. (reference 21012), King's Together Rapid COVID-19 Call awards to M.H.M. and K.J.D., a Huo Family Foundation Award to M.H.M. and K.J.D., Chronic Disease Research Foundation award CDRF-22/2020 to K.J.D. and M.H.M., and Wellcome Trust Investigator Award 106223/Z/14/Z to M.H.M. C.G. was supported by the MRC-KCL Doctoral Training Partnership in Biomedical Sciences (MR/N013700/1). Fondation Dormeur, Vaduz funded equipment to K.J.D.

AUTHOR CONTRIBUTIONS

P.H. and H.d.L. designed the research, performed the research, analyzed the data, and wrote the manuscript. K.J.D., T.L., H.K., and M.M. performed the research and reviewed the manuscript. F.C., R.D., S.S., K.R., D.M., D.A., DA, S.K., C.H., and V.M. assisted with patient recruitment and reviewed the manuscript. C.S., J.S., A.O.R., and A.E. assisted with patient recruitment and patient interviews and reviewed the manuscript. V.M. and H.d.L. share senior authorship.

DECLARATION OF INTERESTS

P.H. received research funding from Bristol Myers Squibb and speaker fees from Incyte. D.M. received speaker fees and advisory boards Novartis, Celgene, and Jazz pharmaceuticals. S.K. received Celgene and Novartis research grant and Alexion speaker honorarium. C.H. received speaker fees from Novartis, Janssen, CTI, Celgene, and Medscape and has served on the advisory board for Incyte, CTI, Sierra Oncology, Novartis, Celgene, Roche, AOP Pharma, Geron, and Astra Zenica. H.d.L. has received grants and speakers fees from Bristol Myers Squibb and Incyte and speaker fees from Novartis.

REFERENCES

- Addeo, A., Shah, P.K., Bordry, N., Hudson, R.D., Albracht, B., Di Marco, M., Kaklamani, V., Dietrich, P.Y., Taylor, B.S., Simand, P.F., et al. (2021). Immunogenicity of SARS-CoV-2 messenger RNA vaccines in patients with cancer. *Cancer Cell* 39, 1091–1098.
- Bakouny, Z., Hawley, J.E., Choueiri, T.K., Peters, S., Rini, B.I., Warner, J.L., and Painter, C.A. (2020). COVID-19 and cancer: current challenges and perspectives. *Cancer Cell* 38, 629–646.
- Greenberger, L.M., Saltzman, L.A., Senefeld, J.W., Johnson, P.W., DeGennaro, L.J., and Nichols, G.L. (2021). Antibody response to SARS-CoV-2 vaccines in patients with hematologic malignancies. *Cancer Cell* 39, 1031–1033.
- Harrington, P., Doores, K.J., Radia, D., O'Reilly, A., Lam, H.P.J., Seow, J., Graham, C., Lechmere, T., McLornan, D., Dillon, R., et al. (2021). Single dose of BNT162b2 mRNA vaccine against severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) induces neutralising antibody and polyfunctional T-cell responses in patients with chronic myeloid leukaemia. *Br. J. Haematol.* 194, 999–1006.
- Le Bourgeois, A., Coste-Burel, M., Guillaume, T., Peterlin, P., Garnier, A., Béné, M.C., and Chevallier, P. (2021). Safety and antibody response after 1 and 2 doses of BNT162b2 mRNA vaccine in recipients of allogeneic hematopoietic stem cell transplant. *JAMA Netw. Open* 4, e2126344.
- Peng, Y., Mentzer, A.J., Liu, G., Yao, X., Yin, Z., Dong, D., Dejnirattisai, W., Rostron, T., Supasa, P., Liu, C., et al. (2020). Broad and strong memory CD4⁺ and CD8⁺ T cells induced by SARS-CoV-2 in UK convalescent COVID-19 patients. *bioRxiv*, 2020.06.05.134551.
- Redjoul, R., Le Bouter, A., Beckerich, F., Fourati, S., and Maury, S. (2021). Antibody response after second BNT162b2 dose in allogeneic HSCT recipients. *Lancet* 398, 298–299.
- Seow, J., Graham, C., Merrick, B., Acors, S., Pickering, S., Steel, K.J.A., Hemmings, O., O'Byrne, A., Kouphou, N., Galao, R.P., et al. (2020). Longitudinal observation and decline of neutralizing antibody responses in the three months following SARS-CoV-2 infection in humans. *Nat. Microbiol.* 5, 1598–1607.
- Sharma, A., Bhatt, N.S., St Martin, A., Abid, M.B., Bloomquist, J., Chemaly, R.F., Dandoy, C., Gauthier, J., Gowda, L., Perales, M.A., et al. (2021). Clinical characteristics and outcomes of COVID-19 in haematopoietic stem-cell transplantation recipients: an observational cohort study. *Lancet Haematol.* 8, e185–e193.
- Thakkar, A., Gonzalez-Lugo, J.D., Goradia, N., Gali, R., Shapiro, L.C., Pradhan, K., Rahman, S., Kim, S.Y., Ko, B., Sica, R.A., et al. (2021). Seroconversion rates following COVID-19 vaccination among patients with cancer. *Cancer Cell* 39, 1081–1090.