

# Phylogenetic-based nomenclatural proposals for *Ophiocordycipitaceae* (*Hypocreales*) with new combinations in *Tolypocladium*

C. Alisha Quandt<sup>1</sup>, Ryan M. Kepler<sup>2</sup>, Walter Gams<sup>3</sup>, João P. M. Araújo<sup>4</sup>, Sayaka Ban<sup>5</sup>, Harry C. Evans<sup>6</sup>, David Hughes<sup>4,7</sup>, Richard Humber<sup>8</sup>, Nigel Hywel-Jones<sup>9</sup>, Zengzhi Li<sup>10</sup>, J. Jennifer Luangsa-ard<sup>11</sup>, Stephen A. Rehner<sup>2</sup>, Tatiana Sanjuan<sup>12</sup>, Hiroki Sato<sup>13</sup>, Bhushan Shrestha<sup>14</sup>, Gi-Ho Sung<sup>15</sup>, Yi-Jian Yao<sup>16</sup>, Rasoul Zare<sup>17</sup>, and Joseph W. Spatafora<sup>1</sup>

<sup>1</sup>Department of Botany and Plant Pathology, Oregon State University, Corvallis, OR 97331, USA; corresponding author e-mail: owensbyc@science.oregonstate.edu

<sup>2</sup>USDA-ARS, Systematic Mycology and Microbiology Laboratory, Beltsville, MD 20705, USA

<sup>3</sup>Formerly CBS-KNAW, Fungal Biodiversity Centre, Utrecht, The Netherlands

<sup>4</sup>Department of Biology, Pennsylvania State University, University Park, State College, PA 16802, USA

<sup>5</sup>Biological Resource Center, National Institute of Technology and Evaluation 2-5-8 Kazusakamatai, Kisarazu, Chiba 292-0818, Japan

<sup>6</sup>CAB International, E-UK Centre, Egham, Surrey TW20 9TY, UK

<sup>7</sup>Department of Entomology, Center for Infectious Disease Dynamics, Pennsylvania State University, University Park, State College, PA 16802, USA

<sup>8</sup>USDA-ARS Biological Integrated Pest Management Research, Robert W. Holley Center for Agriculture and Health, Ithaca, NY14853, USA

<sup>9</sup>Milton Biotech Ltd, Krathum Rai, Nong Chok, Bangkok 10530, Thailand

<sup>10</sup>Department of Forestry, Anhui Agricultural University, Hefei, Anhui 230036, China

<sup>11</sup>Microbe Interaction Laboratory, National Center for Genetic Engineering and Biotechnology (BIOTEC), 113 Thailand Science Park, Phahonyothin Rd, Khlong Nueng, Khlong Luang, Pathum Thani 12120, Thailand

<sup>12</sup>Laboratorio de Taxonomía y Ecología de Hongos, Instituto de Biología, Facultad de Ciencias Naturales, Universidad de Antioquia, Medellín, Colombia and Laboratorio de micología y fitopatología, Departamento Ciencias Biológicas, Universidad de los Andes, Bogotá, Colombia

<sup>13</sup>Department of Forest Entomology, Forestry and Forest Products Research Institute, 1 Matsunosato, Tsukuba, Ibaraki 305, Japan

<sup>14</sup>Institute of Life Science and Biotechnology, Sungkyunkwan University, Suwon 440-746, Korea

<sup>15</sup>Mushroom Research Division, National Institute of Horticultural and Herbal Science, Rural Development Administration, Eumseong 369-873, Korea

<sup>16</sup>State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China

<sup>17</sup>Department of Botany, Iranian Research Institute of Plant Protection, P.O. Box 1454, Tehran 19395, Iran

**Abstract:** *Ophiocordycipitaceae* is a diverse family comprising ecologically, economically, medicinally, and culturally important fungi. The family was recognized due to the polyphyly of the genus *Cordyceps* and the broad diversity of the mostly arthropod-pathogenic lineages of *Hypocreales*. The other two cordyceps-like families, *Cordycipitaceae* and *Clavicipitaceae*, will be revised taxonomically elsewhere. Historically, many species were placed in *Cordyceps*, but other genera have been described in this family as well, including several based on anamorphic features. Currently there are 24 generic names in use across both asexual and sexual life stages for species of *Ophiocordycipitaceae*. To reflect changes in Art. 59 in the *International Code of Nomenclature for algae, fungi, and plants* (ICN), we propose to protect and to suppress names within *Ophiocordycipitaceae*, and to present taxonomic revisions in the genus *Tolypocladium*, based on rigorous and extensively sampled molecular phylogenetic analyses. When approaching this task, we considered the principles of priority, monophyly, minimizing taxonomic revisions, and the practical utility of these fungi within the wider biological research community.

## Key words:

arthropod-pathogens

Article 59

new combinations

nomenclature

*Ophiocordycipitaceae*

*Tolypocladium*

**Article info:** Submitted: 31 March 2014; Accepted: 2 June 2014; Published: 19 June 2014.

## BACKGROUND

The revision of Art. 59 in the *International Code of Nomenclature for algae, fungi, and plants* (ICN; McNeill *et al.* 2012) has created a major task for mycologists, who must now reconcile under one name various possible names existing for different morphs of the same species of fungus

(Hibbett & Taylor 2013). Groups have already begun to propose names which should be protected or suppressed within *Hypocreales* in accordance with the 'one fungus one name' policy (Geiser *et al.* 2013, Rossman *et al.* 2013, Leuchtmann *et al.* 2014, Johnston *et al.* 2014, Kepler *et al.* 2014) and others are in progress. Here, we seek to retain names in *Ophiocordycipitaceae* with the goal of harmonizing

© 2014 International Mycological Association

You are free to share - to copy, distribute and transmit the work, under the following conditions:

**Attribution:** You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).

**Non-commercial:** You may not use this work for commercial purposes.

**No derivative works:** You may not alter, transform, or build upon this work.

For any reuse or distribution, you must make clear to others the license terms of this work, which can be found at <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>. Any of the above conditions can be waived if you get permission from the copyright holder. Nothing in this license impairs or restricts the author's moral rights.

priority, monophyly, simplicity of taxonomic revisions, and minimization of disruption to the research community.

The family *Ophiocordycipitaceae* was described by Sung et al. (2007) to accommodate species that were determined to be phylogenetically distinct from *Cordycipitaceae* and *Clavicipitaceae* s.str. Asexual morphologies in *Ophiocordycipitaceae* show a tremendous range of variation, some of which are restricted in their phylogenetic distribution while others are often found in disparate lineages. For example, *Verticillium* is a common asexual morph of many species in several hypocrealean families, including *Ophiocordycipitaceae*, *Cordycipitaceae* and *Clavicipitaceae* (see Zare et al. 2000, Sung et al. 2001, 2007, and Gams & Zare 2001).

*Ophiocordyceps* is the most speciose genus of the family, and was described originally by Petch (1931a) for species of *Cordyceps* that have septate ascospores that do not disarticulate into part-spores at maturity and asci with inconspicuous apical caps (Petch 1931a, 1933). Kobayasi (1941) later used *Ophiocordyceps* as a subgeneric classification of the genus *Cordyceps*, but Sung et al. (2007) restored *Ophiocordyceps* to the rank of genus to include those *Cordyceps* species within *Ophiocordycipitaceae* forming a sister clade with the genus *Elaphocordyceps* (see below). The type of the genus is *O. blattae*, a rarely collected cockroach pathogen for which no culture or molecular data are available.

Asexual generic names associated with *Ophiocordyceps* include *Sorosporella*, the oldest name still in use for species in the clade, *Hirsutella*, *Hymenostilbe*, *Stilbella*, *Syngliocladium*, and *Paraisaria*. *Hirsutella* species typically produce one to several conidia in a limited mucus droplet borne on basally subulate phialides that taper into slender necks (Gams & Zare 2003). *Hymenostilbe* was proposed by Petch (1931b), and there is some evidence to support restricting its use within the genus *Ophiocordyceps* to the ‘*O. sphecocephala* clade’, most species of which sporulate from adult insects (Sung et al. 2007, Luangsa-ard et al. 2011a). These taxa produce conidia singly from multiple denticles on conidiogenous cells forming a palisade-like layer along the entire outer surface of synnemata (Mains 1950). The *Stilbella* morphology has been applied broadly among species associated with *Ophiocordyceps*, as well as to fungi later reclassified in other genera (Seifert 1985, Gräfenhan et al. 2011). *Stilbella* species often produce aggregate synnemata with a fertile, terminal head of conidia. *Syngliocladium* spp. often have laterally arising conidiophores similar in morphology to the hypocreaceous asexual morph *Gliocladium*, and they may be either synnematous or mononematous on their arthropod hosts (Petch 1932, Hodge et al. 1998). *Sorosporella*, a chlamydospore producing spore state, has been linked as a synasexual morph of *Syngliocladium* (Speare 1917, 1920), but the two morphologies are not always produced by all species (Hodge et al. 1998, Evans & Shah 2002). Species of *Paraisaria* possess feathery synnemata which fruit from arthropod hosts, and several species have been linked via cultural and molecular data to the *O. gracilis* clade (Samson & Brady 1983, Sung et al. 2007, Evans et al. 2010). Names of genera associated with *Ophiocordyceps* whose types are located outside of *Hypocreales* include *Tilachlidiosis*

and *Podonectria*, members of the *Agaricomycetes* and *Dothideomycetes*, respectively (Rossman 1978, Stalpers et al. 1991, Hughes et al. 2001, Boonmee et al. 2011). Despite the large number of taxa associated with *Ophiocordyceps*, a lack of support for internal nodes resulting in equivocal topologies has limited inferences about relationships within the genus in previous studies (Sung et al. 2007).

The most notable species in the *Ophiocordyceps* clade is *O. sinensis*, which is nearly double the price of gold by weight (Stone 2008, Shrestha & Bawa 2013) and the subject of intense research, especially in China (Shrestha et al. 2010, Hu et al. 2013, Ren & Yao 2013, Bushley et al. 2013a, etc.). Almost exclusively found parasitizing the larvae of ghost moths (*Hepialidae: Thitarodes*) in the alpine and sub-alpine pastures of the Tibetan plateau and the Himalayas, this species is undergoing heavy, possibly unsustainable, and destructive harvesting (Cannon et al. 2009, Shrestha & Bawa 2013).

The recently described genus *Elaphocordyceps* is typified by *E. ophioglossoides*, one of the first *Cordyceps* species to be described. Species in *Elaphocordyceps* are mostly parasites of the ectomycorrhizal truffle genus *Elaphomyces* (Ascomycota, Eurotiales). The majority of *Elaphocordyceps* species have no known asexual morph, but where known they produce ones which are verticillium-like or *Tolypocladium* (Sung et al. 2007). There are a few *Elaphocordyceps* species known to be entomopathogens, including three cicada pathogens (*E. inegoensis*, *E. paradoxa*, and *E. toriharamontana*), and one beetle pathogen, *E. subsessilis* (syn. *Tolypocladium inflatum*) (Hodge et al. 1996, Sung et al. 2007). *Tolypocladium inflatum* (a name conserved by the rejection of *Pachybasium niveum*; Dreyfuss & Gams 1994), is a medicinally important fungus and the subject of much research due to its production of the immunosuppressant drug, cyclosporin A (Survase et al. 2011, Bushley et al. 2013b). The other species of *Tolypocladium* have no known sexual morphs and have mainly been isolated from soil (Gams 1971, Bissett 1983) or observed parasitizing rotifers or insects (Barron 1980, 1981, 1983, Samson & Soares 1984, Weiser et al. 1991). The asexually typified genus *Chaunopycnis* is also related to this clade (Bills et al. 2002) and has been isolated mainly from soil samples (Gams 1980, Bills et al. 2002), although one species was isolated from epilithic Antarctic lichens (Möller & Gams 1993). The similarity of conidiogenesis between *Chaunopycnis* and *Tolypocladium* was noted in the original description of *Chaunopycnis* (Gams 1980), and its phialides often taper in a manner similar to those of *Tolypocladium*. Interestingly, these two genera have also been linked by their shared production of cyclosporin A (Traber & Dreyfuss 1996). Two of the described *Chaunopycnis* species produce loosely enclosed conidiomata, a morphology not seen in other members of the clade or within *Ophiocordycipitaceae* as a whole.

The relationships among the species of the *Purpureocillium* clade were recently reviewed by Luangsa-ard et al. (2011b). The genus was proposed to encompass taxa closely related to *Purpureocillium lilacinum* (syn. *Paecilomyces lilacinus*) and consists of species with purple-hued conidia, including *Nomuraea atypicola* and *Isaria*

*takamizusanensis*. The type of *Nomuraea* is *N. rileyi* (syn. *N. prasina*), which has recently been synonymized with *Metarhizium* (Kepler et al. 2014). The type of *Isaria* is a member of *Cordycipitaceae* (Gams et al. 2005, Hodge et al. 2005, Luangsa-ard et al. 2011b). While *N. atypicola* and *I. takamizusanensis* have not been addressed taxonomically, other studies found close relationships between these taxa and *Purpureocillium* (Sung et al. 2007, Perdomo et al. 2013). *Nomuraea atypicola* is the asexual morph of *C. cylindrica* (Hywel-Jones & Sivichai 1995), the only sexual morph described for this clade and one of the “residual” *Cordyceps* s. lat. left without reassignment to any phylogenetically redefined genus by Sung et al. (2007).

Nematode pathogens have been described in many genera throughout *Hypocreales*. The largest and oldest of these is the asexually typified genus *Harposporium*. Most *Harposporium* species produce crescent-shaped or helicoid conidia that are ingested by their hosts and become lodged in the upper portions of the digestive tract (Barron 1977). Conidia are produced on spherical conidiogenous cells, and several species are known to produce hirsutella-like synasexual morphs (Hodge et al. 1997, Chaverri et al. 2005, Li et al. 2005). While the majority of *Harposporium* species are known from nematodes, these fungi are common in the soil and several studies have reported an entomopathogenic ecology as well (e.g., Shimazu & Glockling 1997, Evans & Whitehead 2005). In 2005, Chaverri et al. reported the asexual-sexual morph connection between *Harposporium* and *Podocrella*, an arthropod-pathogenic genus. Several researchers initially described nematophagous taxa in the originally plant-pathogenic genus *Meria* (Vuillemin 1896, Drechsler 1941), but this genus was found to be polyphyletic (Gams & Jansson 1985), and for this reason *Drechmeria* was erected for the nematophagous meria-like taxa in *Hypocreales*. The type of *Drechmeria*, *D. coniospora*, has cone-shaped conidia whose conidiogenous cells are not basally swollen as in *Harposporium*. One protozoan-infecting species of *Drechmeria*, *D. harposporioides*, produces crescent-shaped conidia similar to those of *Harposporium* (Barron & Szijarto 1982). *Haptocillium* was erected for asexual nematode pathogens bearing verticillate phialides and whose conidia are not ingested but adhere to the surface of their hosts (Zare & Gams 2001).

*Polycephalomyces* represents a diverse clade that is currently *incertae sedis* within *Hypocreales*, as its placement has lacked support in previous molecular studies (Kepler et al. 2013). Of particular uncertainty was whether *Polycephalomyces* and its closest related taxon, *C. pleuric平ata*, formed a sister clade to *Ophiocordycipitaceae*, or if it was more closely related to *Clavicipitaceae*. Many morphological characters are shared between *Ophiocordycipitaceae* and *Polycephalomyces*. For example, numerous species in both clades produce hirsutella-like anamorphs with conidia often borne in a slimy mass (Seifert 1985). In addition, sexual sporing structures of *Polycephalomyces* often possess a wiry, tough, carbonaceous stipe which is a common morphology of *Ophiocordyceps* (Kepler et al. 2013). Many species within this genus are known mycoparasites of other hypocrealean entomopathogens and myxomycetes, but there are also several species of

entomopathogens. *Cordyceps pleuric平ata* was deemed a residual species of *Cordyceps* of uncertain placement by Kepler et al. (2013), due to a lack of statistical support joining that species and *Polycephalomyces*.

In this paper we expand the taxon sampling presented in Sung et al. (2007) by 222 hypocrealean isolates. This includes sexual and asexual states which provide the framework for addressing the nomenclatural issues demanded by changes to the most recent ICN.


## MATERIALS AND METHODS

Sequences from five nuclear loci, including the small and large subunits of the rDNA (SSU and LSU), the transcription elongation factor-1α (TEF), and the first and second largest subunits of RNA polymerase II (RPB1 and RPB2) were used for phylogenetic analyses. DNA extraction and PCR amplification were carried out as previously described (Kepler et al. 2013). Sequencing reactions were performed at the University of Washington High-Throughput Genomics Center (Seattle, WA) with the primers used for the initial amplifications. All other sequences were collected from GenBank. Efforts were made for all specimens to have data for at least three of the five genes to be considered in our analyses. However, certain taxa for which only one or two genes were available were included due to the importance in addressing the taxonomic issues at hand (Table 1).

Raw sequences were processed, aligned, and gaps excluded as in Kepler et al. (2013), using the programs MAFFT v. 6 (Katoh et al. 2002, Katoh & Toh 2008), Geneious v. 7.0.6 (Biomatters, available <http://www.geneious.com>), and Gblocks (Talavera & Castresana 2007). The final alignment length was 4570 nucleotides - 1023 for SSU, 879 for LSU, 987 for TEF, 646 for RPB1, and 1035 for RPB2. RAxML v. 7.6.6 (Stamatakis 2006) was used to perform Maximum likelihood (ML) estimation of the phylogeny with 500 bootstrap replicates on the concatenated dataset using eleven data partitions. These included one each for SSU and LSU, and three for each of the three codon positions of the protein coding genes, TEF, RPB1, and RPB2. The GTR-GAMMA model of nucleotide substitution was used.

## RESULTS AND DISCUSSION

Our results are in agreement with the overall phylogenetic structure of the order *Hypocreales* put forth by Sung et al. (2007). Nomenclatural issues for taxa in the other two families of cordyceps-like organisms, *Cordycipitaceae* and *Clavicipitaceae*, will be presented elsewhere or have already been published (Leuchtmann et al. 2014, Kepler et al. 2014). Based on this exhaustive phylogenetic reconstruction (Fig. 1), we recognize six genera within *Ophiocordycipitaceae*: *Ophiocordyceps*, *Tolypocladium*, *Purpureocillium*, *Harposporium*, *Drechmeria*, and *Polycephalomyces* (Table 2). This framework will provide clarity for researchers, ease of communication for instructors, and phylogenetic taxonomy around which to investigate the evolution of life histories (e.g. morphology, ecology).



**Fig. 1.** ML tree of Ophiocordycipitaceae obtained using RAxML to analyze the concatenated five gene dataset (SSU, LSU, TEF, RPB1, and RPB2). Proposed genus level names to protect are delimited, but names of individual species have not been changed on the leaves of the tree, to demonstrate the diversity of taxa sampled. Values above branches represent MLBP proportions greater than or equal to 70 % from 500 replicates. Inset tree shows the larger phylogeny of Hypocreales.

## TAXONOMY

### *Ophiocordyceps* Petch 1931

*Ophiocordyceps sensu* Sung et al. (2007) is resolved as a well-supported (MLBP=77) clade (Fig. 1, Node 3). This clade is speciose, diverse, and almost exclusively comprises insect pathogens. In spite of increased taxon sampling, current reconstructions fail to find strong statistical support at the internal nodes, and therefore we refrain from defining infrageneric groupings (Fig. 1). While *Sorosporella* is the oldest name for any members in this clade, there are only two described species, and Evans & Shah (2002) argued *Sorosporella* should be synonymized with *Syngliocladium* instead of being recognized as a an asexual morph, as *Synnematium* was previously treated with respect to *Hirsutella* (Evans & Samson 1982). We propose, therefore, to suppress the use of *Sorosporella* for this clade. *Hirsutella* is the next oldest name, but the type, *H. entomophila*, which was described growing from adult Coleoptera, has not been sampled and no culture of this species is available. Sung et al. (2007) argued that the *Hirsutella* morphology was phylogenetically informative for the ‘*O. unilateralis* group’ which they resolved as paraphyletic, a topology recovered in the current analyses as well (Fig 1, Nodes 4 and 5). However, the *Hirsutella* morphology is observed in other clades (e.g. *Harposporium*, *Polycephalomyces*, *Clavicipitaceae*), and while it is difficult to place the type species based on morphology alone, it appears from its original description to be morphologically and ecologically (as a parasite of adults) similar to species of *Hymenostilbe* found in the ‘*O. sphecocephala*’ clade and not *Hirsutella* of the ‘*O. unilateralis* group’ (Patouillard 1892). Another reason for suppressing the use of *Hirsutella* for this clade is the larger number of new combinations that would have to be made – 178 for *Ophiocordyceps* vs. 77 for *Hirsutella* – as the vast majority of species encompassed here are currently described as *Ophiocordyceps*. Also, preservation of the name “cordyceps” within the name of *O. sinensis* is considered paramount given its economic, medicinal, and cultural importance in addition to being the most widely known and researched species in the clade (Shrestha et al. 2010).

At this time, we also propose to suppress the use of the other names proposed for taxa in this clade, including *Hymenostilbe*, *Syngliocladium*, and *Paraisaria*, because these names are younger, and they contain fewer associated taxa than either *Ophiocordyceps* or *Hirsutella*. Our results suggest the restriction of *Hymenostilbe* to the ‘*O. sphecocephala* clade’ (Fig. 1, Node 6) which occupies a long branch and has strong support (MLBP=100), however, because the other internal nodes of the clade do not receive support, we refrain from making this distinction now as it would result in a paraphyletic *Ophiocordyceps*. These analyses place one species of *Stilbella*, *S. buquetii*, in this clade, while other studies (Seifert 1985, Gräfenhan et al. 2011) have placed other *Stilbella* species in *Nectriaceae*, *Bionectriaceae*, or *Polycephalomyces*, and the current placement of *Stilbella* remains *Hypocreales incertae sedis* (Kirk et al. 2008). The type of *Stilbella*, a coprophile, has yet to be considered in a phylogenetic context, and for these reasons we do not address that name here, but reject the use of that name for

this clade. Therefore, we propose to protect *Ophiocordyceps* as the genus name for the entire clade, while acknowledging that future studies including more data and taxonomic sampling may provide better resolution of the relationships within the genus and a narrower concept of *Ophiocordyceps*.

### *Tolypocladium* W. Gams 1971

*Tolypocladium* is proposed for protection over the other two generic names in the clade, *Elaphocordyceps* and *Chaunopycnis*. The clade itself is well supported (MLBP=97) in this and other published analyses (Sung et al. 2007, Kepler et al. 2013). However, relationships between species in this clade are very sensitive to taxon sampling, and there is little bootstrap support for internal branches from the current data to justify more than one name for this clade. The asexual-sexual morph connection between *Tolypocladium* and some *Elaphocordyceps* species has been known for several years (Hodge et al. 1996), although where known most *Elaphocordyceps* spp. do not possess the morphology associated with *Tolypocladium* (Sung et al. 2007). While this may cause some short-term confusion, the alternative would be to name the clade *Elaphocordyceps* (which would cause the fewest name changes, 12 vs. 26 for *Tolypocladium*) and suppress *Tolypocladium*, a much more widely known, medicinally important, and older name, and therefore we find this a poor option. In this analysis the *Chaunopycnis* species sampled form a monophyletic clade which is the most divergent group within the clade. However, this may be the result of limited taxon and genetic sampling; only small subunit rDNA data for the sampled *Chaunopycnis* species was available for these analyses.

Here, we present a list of 26 new combinations within the genus *Tolypocladium*, which we emend to include species whose anamorphic forms do not possess inflated phialide bases, but that do form a single monophyletic clade encompassing a large number of truffle parasites, several insect pathogens, rotifer pathogens, and several fungi isolated to date only from soil.

### *Tolypocladium* W. Gams, Persoonia 6: 185 (1971).

*Synonyms:* *Chaunopycnis* W. Gams, Persoonia 11: 75 (1980).

*Elaphocordyceps* G.H. Sung & Spatafora, Stud. Mycol. 57: 36 (2007).

*Circumscription:* The genus *Tolypocladium* is emended here to apply to all descendants of the node defined in the reference phylogeny (Fig. 1) as the terminal *Tolypocladium* clade. It is the least inclusive clade containing *T. album*, *T. capitatum*, *T. cylindrosporum*, *T. fractum*, *T. inflatum*, *T. japonicum*, *T. longisegmentum*, *T. ophioglossoides*, and *T. pustulatum*. No definitive synapomorphies are known for the clade. Morphologies associated with sexual reproductive states include robust stipitate stroma with clavate to capitate clava (e.g. *T. capitatum*) to highly reduced stroma comprising rhizomorphs and aggregated perithecia (e.g. *T. inflatum*); perithecia may be immersed and ordinal to the long axis of the stroma or superficial and produced on a highly reduced stromatic pad; ascii are single-walled, long and cylindrical with a pronounced apical cap; ascospores

are filiform, approximately as long as ascospores, septate and typically disarticulate into part-spores. Where known, asexual states include morphologies described as *Tolypocladium sensu* Gams (1970), *Chaunopycnis sensu* Gams (1979), or verticillium-like. Ecologies include parasites and pathogens of insects, rotifers and fungi, as well as, soil-inhabiting.

Type: *Tolypocladium inflatum* W. Gams 1971.

**Tolypocladium inflatum** W. Gams, *Persoonia* **6**: 185 (1971), nom. cons.

Synonyms: *Cordyceps subsessilis* Petch, *Trans. Brit. Mycol. Soc.* **21**: 39 (1937).

*Elaphocordyceps subsessilis* (Petch) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

*Cordyceps facis* Kobayasi & Shimizu, *Trans. Mycol. Soc. Japan* **23**: 361 (1982); as 'Codyceps'.

**Tolypocladium album** (W. Gams) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808698

Basionym: *Chaunopycnis alba* W. Gams, *Persoonia* **11**: 75 (1979).

**Tolypocladium capitatum** (Holmsk. : Fr.) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808699

Basionym: *Clavaria capitata* Holmsk., *Beata Ruris Otia Fung. Dan.* **1**: 38 (1790).

Synonyms: *Sphaeria capitata* (Holmsk. : Fr.) Pers., *Comm. Fung. Clav.* **13** (1797); Fr., *Syst. Mycol.* **2**: 324 (1822).

*Cordyceps capitata* (Holmsk.: Fr.) Link, *Handb. Erk. Gew.* **3**: 347 (1833).

*Torrubia capitata* (Holmsk. : Fr.) Tul. & C. Tul., *Sel. Fung. Carpol.* **3**: 22 (1865).

*Elaphocordyceps capitata* (Holmsk. : Fr.) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

*Sphaeria agariciformis* Bolton, *Hist. Fung. Halifax*: 130 (1789).

*Cordyceps agariciformis* (Bolton) Seaver, *N. Amer. Fl.* **3**: 53 (1910).

*Cordyceps canadensis* Ellis & Everh., *Bull. Torrey Bot. Club* **25**: 501 (1898).

*Cordyceps capitata* var. *canadensis* (Ellis & Everh.) Lloyd, *Mycol. Writ.* **5**: 609 (1916).

*Cordyceps nigriceps* Peck, *Bull. Torrey Bot. Club* **27**: 21 (1900).

**Tolypocladium delicatistipitatum** (Kobayasi) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808700

Basionym: *Cordyceps delicatistipitata* Kobayasi, *Bull. Natn. Sci. Mus., Tokyo* **5** (2, no. 47): 79 (1960); as 'delicatostipitata'.

Synonym: *Elaphocordyceps delicatistipitata* (Kobayasi) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

**Tolypocladium fractum** (Mains) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808701

Basionym: *Cordyceps fracta* Mains, *Bull. Torrey. Bot. Club* **84**: 250 (1957).

Synonym: *Elaphocordyceps fracta* (Mains) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

**Tolypocladium inegoense** (Kobayasi) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808702

Basionym: *Cordyceps inegoensis* Kobayasi, *Bull. Natn. Sci. Mus., Tokyo* **6**: 292 (1963)

Synonyms: *Elaphocordyceps inegoensis* (Kobayasi) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007); as 'inegoënsis'.

**Tolypocladium intermedium** (S. Imai) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808703

Basionym: *Cordyceps intermedia* S. Imai, *Proc. Imp. Acad. Japan* **10**: 677 (1934).

Synonyms: *Elaphocordyceps intermedia* (S. Imai) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

**Tolypocladium intermedium** f. **michinokuense** (Kobayasi & Shimizu) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808704

Basionym: *Cordyceps intermedia* f. *michinokuensis* Kobayasi & Shimizu, *Bull. Natn. Sci. Mus., Tokyo*, B **8**: 116 (1982).

Synonym: *Elaphocordyceps intermedia* f. *michinokuensis* (Kobayasi & Shimizu) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007); as 'michinokuënsis'.

**Tolypocladium japonicum** (Lloyd) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808705

Basionym: *Cordyceps japonica* Lloyd, *Mycol. Writ.* **6** (Letter 62): 913 (1920).

Synonyms: *Elaphocordyceps japonica* (Lloyd) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

*Cordyceps umemurae* S. Imai, *Trans. Sapporo Nat. Hist. Soc.* **11**: 32 (1930) [1929]; as 'umemurai'.

**Tolypocladium jezoense** (S. Imai) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808706

Basionym: *Cordyceps jezoensis* S. Imai, *Trans. Sapporo Nat. Hist. Soc.* **11**: 33 (1930) [1929]. Synonym: *Elaphocordyceps jezoensis* (S. Imai) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007); as 'jezoënsis'.

**Tolypocladium longisegmentum** (Ginns) Quandt, Kepler & Spatafora, comb. nov.

MycoBank MB808856

Basionym: *Cordyceps longisegmentis* Ginns, *Mycologia* **80**: 219 (1988).

Synonym: *Elaphocordyceps longisegmentis* (Ginns) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

**Tolypocladium minazukiense** (Kobayasi & Shimizu) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808857  
*Basionym:* *Cordyceps minazukiensis* Kobayasi & Shimizu, *Bull. Natn. Sci. Mus., Tokyo, B* **8**: 117 (1982).  
*Synonym:* *Elaphocordyceps minazukiensis* (Kobayasi & Shimizu) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (1982).

**Tolypocladium miomoteanum** (Kobayasi & Shimizu) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808858  
*Basionym:* *Cordyceps miomoteana* Kobayasi & Shimizu, *Bull. Natn. Sci. Mus., Tokyo, B* **8**: 118 (1982).  
*Synonym:* *Elaphocordyceps miomoteana* (Kobayasi & Shimizu) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (1982).

**Tolypocladium ophioglossoides** (Ehrh. ex J.F. Gmel.) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808859  
*Basionym:* *Sphaeria ophioglossoides* Ehrh. ex J.F. Gmel., *Syst. Na.*, 13<sup>th</sup> edn **2**: 1474 (1792).  
*Synonyms:* *Sphaeria ophioglossoides* Ehrh., *Pl. Crypt. Exs. fasc.* 16 no. 160 (1789); nom. inval. (Art. 38.1).  
*Cordyceps ophioglossoides* (Ehrh. ex G.F. Gmel.) Link, *Handb. Erk. Gew.* **3**: 347 (1833) : Fr., *Syst. Mycol.* **2**: 324 (1822).  
*Torrubia ophioglossoides* (Ehrh. ex G.F. Gmel.) Tul. & C. Tul., *Sel. Fung. Carp.* **3**: 20 (1865).  
*Elaphocordyceps ophioglossoides* (Ehrh. ex G.F. Gmel.) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).  
*Clavaria parasitica* Willd., *Fl. Berol. Prodr.*: 405 (1787).  
*Cordyceps parasitica* (Willd.) Henn., *Nerthus* **6**: 4 (1904).

**Tolypocladium ophioglossoides f. album** (Kobayasi & Shimizu ex Y.J. Yao) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808860

*Basionym:* *Cordyceps ophioglossoides f. alba* Kobayasi & Shimizu ex Y.J. Yao, *Acta Mycol. Sin.* **14**: 257 (1995).  
*Synonym:* *Elaphocordyceps ophioglossoides f. alba* (Kobayasi & Shimizu ex Y.J. Yao) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

**Tolypocladium ophioglossoides f. cuboides** (Kobayasi) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808861  
*Basionym:* *Cordyceps ophioglossoides f. cuboides* Kobayasi, *Bull. Natn. Sci. Mus., Tokyo* **5** (2, no. 47): 77 (1960).  
*Synonym:* *Elaphocordyceps ophioglossoides f. cuboides* (Kobayasi) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

**Tolypocladium ovalisporum** (C. Möller & W. Gams) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808862  
*Basionym:* *Chaunopycnis ovalispora* C. Möller & W. Gams, *Mycotaxon* **48**: 442 (1993).

**Tolypocladium paradoxum** (Kobayasi) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808863  
*Basionym:* *Cordyceps paradoxa* Kobayasi, *Bulletin of the Biogeogr. Soc. Jap.* **9**: 156 (1939).  
*Synonym:* *Elaphocordyceps paradoxa* (Kobayasi) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

**Tolypocladium pustulatum** (Bills et al.) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808864  
*Basionym:* *Chaunopycnis pustulata* Bills et al., *Mycol. Progr.* **1**: 8 (2002).

**Tolypocladium ramosum** (Teng) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808865  
*Basionym:* *Cordyceps ramosa* Teng, *Sinensis* **7**: 810 (1936).  
*Synonym:* *Elaphocordyceps ramosa* (Teng) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

**Tolypocladium rouxii** (Cand.) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808866  
*Basionym:* *Cordyceps rouxii* Cand., *Mycotaxon* **4**: 544 (1976).  
*Synonym:* *Elaphocordyceps rouxii* (Cand.) G.H. Sung et al., *Stud. Mycol.* **57**: 37 (2007).

**Tolypocladium szemaoense** (M. Zang) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808867  
*Basionym:* *Cordyceps szemaoensis* M. Zang, *Acta Bot. Yunn.* **23**: 295 (2001).  
*Synonym:* *Elaphocordyceps szemaoensis* (M. Zang) G.H. Sung et al., *Stud. Mycol.* **57**: 38 (2007); as 'szemaoënsis'.

**Tolypocladium tenuisporum** (Mains) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808868  
*Basionym:* *Cordyceps tenuispora* Mains, *Bull. Torrey Bot. Club* **84**: 247 (1957).  
*Synonym:* *Elaphocordyceps tenuispora* (Mains) G.H. Sung et al., *Stud. Mycol.* **57**: 38 (2007).

**Tolypocladium toriharamontanum** (Kobayasi) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808869  
*Basionym:* *Cordyceps toriharamontana* Kobayasi, *Bull. Natn. Sci. Mus., Tokyo* **6**: 305 (1963).  
*Synonym:* *Elaphocordyceps toriharamontana* (Kobayasi) G.H. Sung et al., *Stud. Mycol.* **57**: 38 (2007).

**Tolypocladium valliforme** (Mains) Quandt, Kepler & Spatafora, **comb. nov.**  
MycoBank MB808870  
*Basionym:* *Cordyceps valliformis* Mains, *Bull. Torrey Bot. Club* **84**: 250 (1957).  
*Synonym:* *Elaphocordyceps valliformis* (Mains) G.H. Sung et al., *Stud. Mycol.* **57**: 38 (2007).

**Tolypocladium valvatistipitatum** (Kobayasi) Quandt, Kepler & Spatafora, **comb. nov.**  
Mycobank MB808871  
*Basionym:* *Cordyceps valvatistipitata* Kobayasi, *Bull. Natn. Sci. Mus., Tokyo* **5**(2, no. 47): 81 (1960); as ‘*valvatositipitata*’.  
*Synonym:* *Elaphocordyceps valvatistipitata* (Kobayasi) G.H. Sung et al., *Stud. Mycol.* **57**: 38 (2007).

**Tolypocladium virens** (Kobayasi) Quandt, Kepler & Spatafora, **comb. nov.**  
Mycobank MB808872  
*Basionym:* *Cordyceps virens* Kobayasi, *J. Jap. Bot.* **58**: 222 (1983).  
*Synonym:* *Elaphocordyceps virens* (Kobayasi) G.H. Sung et al., *Stud. Mycol.* **57**: 38 (2007).

### Purpureocillium Luangsa-ard et al. 2011

Our findings support those reported by Luangsa-ard et al. (2011b) for the *Purpureocillium* clade, and the change in Art. 59 allows for the inclusion of *N. atypicola* (syn. *Cordyceps cylindrica*) and *Isaria takamizusanensis* within this genus. Shared characters for this clade include purple-hued conidia and pathogenesis of arthropods, although *P. lilacinum* and *P. lavendulum* have been cultured from various substrates (Perdomo et al. 2013), and *P. lilacinum* can cause keratitis and other mycoses in humans and other vertebrates (Pastor & Guarro 2006, Rodríguez et al. 2010). Because this genus is well supported (MLBP=76) as sister to the nematode pathogen clade (Fig. 1), it is important to mention that *P. lilacinum* is frequently collected from nematodes (Luangsa-ard et al. 2011b), and has been used in the biocontrol of plant pathogenic nematodes (Kalele et al. 2006, Castillo et al. 2013).

### Harposporium Lohde 1874 and Drechmeria W. Gams & H.-B. Jansson 1985

Our analyses reconstruct a well-supported (MLBP=76) monophyletic origin of the mostly nematophagous clade of *Ophiocordycipitaceae* (Fig. 1 Node 2). Within this clade, there is strong phylogenetic support for two clades: one containing *Harposporium* and *Podocrella*, and the other consisting of *Drechmeria*, *Haptocillium*, and *Cordyceps gunnii*. The relationship between *Harposporium* and *Podocrella* has already been described (Chaverri et al. 2005), but the revision of Art. 59 requires that one name be chosen for this genus. *Harposporium* is an older name, and the morphology of at least somewhat crescent-shaped conidia is a shared character for this clade. Suppression of *Podocrella* also requires the fewest taxonomic revisions (3 vs 30). For these reasons, we propose to protect *Harposporium* over *Podocrella* (Table 2).

Within the other nematophagous subclade, *Drechmeria* is an older name than *Haptocillium*, and the isolate included in these analyses is nested within the *Haptocillium* isolates sampled. For this reason, we propose to protect *Drechmeria* over *Haptocillium*. The inclusion of *C. gunnii* in this clade also provides a name for this residual taxon of *Cordyceps*. Most species however, are nematophagous (*C. gunnii* being the exception), and conidia may be cone-shaped, formed on conidiogenous cells in rosettes or verticils, or in the case

of *C. gunnii*, paecilomyces-like. We did not have access to molecular data from *D. harposporioides*, but given our finding that the two nematophagous clades in *Ophiocordycipitaceae* are monophyletic in origin, it will be interesting to see if this species, a protozoan pathogen with helical conidia, is truly a member of the *Drechmeria* clade or in fact a species within *Harposporium* that simply lacks the basally swollen conidiogenous cells.

### Polycephalomyces Kobayasi 1941

This study is the first to have definitive ML support (MLBP=82) for the sister relationship between the *Polycephalomyces* clade and *Ophiocordycipitaceae* (Fig. 1 Node 1). Support for this relationship remains even with the exclusion of *C. pleuric capitata*, which is on an early-diverging, long branch within the clade. Two options remain to deal with this finding. Either a new family must be erected to account for this clade, or *Polycephalomyces* and related taxa must be moved into *Ophiocordycipitaceae*. We propose to accept *Polycephalomyces* and *C. pleuric capitata* in *Ophiocordycipitaceae*, where it will be the earliest diverging lineage of the family. The taxonomy of *C. pleuric capitata* will be addressed elsewhere.

## CONCLUSIONS

We present a concise, thorough, phylogenetically relevant, and taxonomically accurate revision of the family *Ophiocordycipitaceae* with the aim of complying with the changes to Art. 59 of the ICN. With the criteria of naming monophyletic taxa, and where possible, of adhering to priority while avoiding changes that would be disruptive to the wider community of researchers, we have proposed to protect six genera within *Ophiocordycipitaceae*, including incorporation of the genus *Polycephalomyces* within the family. We have also formally revised the genus *Tolypocladium*, to reflect the nomenclature suggested by our results.

## ACKNOWLEDGEMENTS

We would like to acknowledge the following individuals for their contribution to this manuscript: Priscila Chaverri, Mingjun Chen, Tsuyoshi Hosoya, Jae-Mo Sung, and James White. An NSF Graduate Research Fellowship supported C.A.Q.

## REFERENCES

- Barron GL (1977) *The Nematode-destroying Fungi*. [Topics in Mycobiology no. 1.] Guelph, ON: Canadian Biological Publications.
- Barron GL (1980) Fungal parasites of rotifers: a new *Tolypocladium* with underwater conidiation. *Canadian Journal of Botany* **58**: 439–442.
- Barron GL (1981) Two new fungal parasites of bdelloid rotifers. *Canadian Journal of Botany* **59**: 1449–1455.
- Barron GL (1983) Structure and biology of a new *Tolypocladium* attacking bdelloid rotifers. *Canadian Journal of Botany* **61**: 2566–2569.

- Barron GL, Szjarto E (1982) A new hyphomycete parasitic on the ciliated protozoans *Vorticella* and *Opercularia*. *Canadian Journal of Botany* **60**: 1031–1034.
- Bills GF, Polishook JD, Goetz MA, Sullivan RF, White Jr. JF (2002) *Chaunopycnis pustulata* sp. nov., a new clavicipitalean anamorph producing metabolites that modulate potassium ion channels. *Mycological Progress* **1**: 3–17.
- Bissett J (1983) Notes on *Tolypocladium* and related genera. *Canadian Journal of Botany* **61**: 1311–1329.
- Boonmee S, Zhang Y, Chomnunti P, Chukeatirote E, Tsui CKM, Bahkali AH, Hyde KD (2011) Revision of lignicolous *Tubeufiaceae* based on morphological reexamination and phylogenetic analysis. *Fungal Diversity* **51**: 63–102.
- Bushley KE, Li Y, Wang WJ, Wang XL, Jiao L, Spatafora JW, Yao YJ (2013a) Isolation of the MAT1-1 mating type idiomorph and evidence for selfing in the Chinese medicinal fungus *Ophiocordyceps sinensis*. *Fungal Biology* **117**: 599–610.
- Bushley KE, Raja R, Jaiswal P, Cumbie JS, Nonogaki M, Boyd AE, Owensby CA, Knaus BJ, Elser J, Miller D, Di Y, McPhail KL, Spatafora JW (2013b) Draft genome sequence of the Cyclosporin producing fungus *Tolypocladium inflatum* reveals complex patterns of secondary metabolite evolution and expression. *PLoS Genetics* **9**: e1003496.
- Cannon PF, Hywel-Jones NL, Maczey N, Norbu L, Samdup T, Lhendup P (2009) Steps towards sustainable harvest of *Ophiocordyceps sinensis* in Bhutan. *Biodiversity and Conservation* **18**: 2263–2281.
- Castillo JD, Lawrence KS, Kloepper JW (2013) Biocontrol of the reniform nematode by *Bacillus firmus* GB-126 and *Paecilomyces lilacinus* 251 on cotton. *Plant Disease* **97**: 967–976.
- Chaverri P, Samuels GJ, Hodge KT (2005) The genus *Podocrella* and its nematode-killing anamorph *Harposporium*. *Mycologia* **97**: 433–443.
- Drechsler C (1941) Some hyphomycetes parasitic on free-living tericolous nematodes. *Phytopathology* **31**: 773–802.
- Dreyfuss M, Gams W (1994) Proposal to reject *Pachybasium niveum* Rostr. in order to retain the name *Tolypocladium inflatum* W. Gams for the fungus that produces cyclosporin. *Taxon* **43**: 660–661.
- Evans HC, Samson RA (1982) *Cordyceps* species and their anamorphs pathogenic on ants (*Formicidae*) in tropical forest ecosystems 1. The *Cephalotes* (*Myrmicinae*) complex. *Transactions of the British Mycological Society* **79**: 431–453.
- Evans HC, Shah PA (2002) Taxonomic status of the genera *Sorosporella* and *Syngliocladium* associated with grasshoppers and locusts (*Orthoptera: Acridoidea*) in Africa. *Mycological Research* **106**: 737–744.
- Evans HC, Whitehead PF (2005) Entomogenous fungi of arboreal *Coleoptera* from Worcestershire, England, including the new species *Harposporium bredonense*. *Mycological Progress* **4**: 91–99.
- Evans HC, Groden E, Bischoff JF (2010) New fungal pathogens of the red ant *Myrmica rubra* from the UK and implications for ant invasions in the USA. *Fungal Biology* **114**: 451–466.
- Gams W (1971) *Tolypocladium*, eine Hyphomycetengattung mit geschwollenen Phialiden. *Persoonia* **6**: 185–191.
- Gams W (1980) *Chaunopycnis alba*, gen. et sp. nov., a soil fungus intermediate between *Moniliales* and *Sphaeropsidales*. *Persoonia* **11**: 75–79.
- Gams W, Jansson H-B (1985) The nematode parasite *Meria coniospora* Drechsler in pure culture and its classification. *Mycotaxon* **22**: 33–38.
- Gams W, Hodge KT, Samson RA, Korf RP, Seifert KA (2005) (1684) Proposal to conserve the name *Isaria* (anamorphic fungi) with a conserved type. *Taxon* **54**: 537.
- Gams W, Zare R (2001) A revision of *Verticillium* section *Prostrata*. III. Generic classification. *Nova Hedwigia* **72**: 329–337.
- Gams W, Zare R (2003) A taxonomic review of the clavicipitaceous anamorphs parasitizing nematodes and other microinvertebrates. In: *Clavicipitalean Fungi: evolutionary biology, chemistry, biocontrol and cultural Impacts* (White JF jr., Bacon CW, Hywel-Jones NL, Spatafora JW, eds): 17–73. New York: Marcel Dekker.
- Geiser DM, Aoki T, Bacon CW, Baker SE, Bhattacharyya MB, et al. (2013) One Fungus, One Name: defining the genus *Fusarium* in a scientifically robust way that preserves longstanding use. *Phytopathology* **103**: 400–408.
- Gräfenhan T, Schroers H-J, Nirenberg HI, Seifert KA (2011) An overview of the taxonomy, phylogeny, and typification of nectriaceous fungi in *Cosmospora*, *Acremonium*, *Fusarium*, *Stilbella*, and *Volutella*. *Studies in Mycology* **68**: 79–113.
- Hibbett DS, Taylor JW (2013) Fungal systematics: is a new age of enlightenment at hand? *Nature Reviews Microbiology* **11**: 129–133.
- Hodge KT, Gams W, Samson RA, Korf RP, Seifert KA (2005) Lectotypification and status of *Isaria* Pers. : Fr. *Taxon* **54**: 485–489.
- Hodge KT, Humber RA, Wozniak CA (1998) *Cordyceps variabilis* and the genus *Syngliocladium*. *Mycologia* **90**: 743–753.
- Hodge KT, Krasnoff SB, Humber RA (1996) *Tolypocladium inflatum* is the anamorph of *Cordyceps subsessilis*. *Mycologia* **88**: 715–719.
- Hodge KT, Viaene NM, Gams W (1997) Two *Harposporium* species with *Hirsutella* synanamorphs. *Mycological Research* **101**: 1377–1382.
- Hu X, Zhang Y, Xiao G, Zheng P, Xia Y, Zhang X, St. Leger RJ, Liu X, Wang C (2013) Genome survey uncovers the secrets of sex and lifestyle in caterpillar fungus. *Chinese Science Bulletin* **58**: 2846–2854.
- Hughes KW, Petersen RH, Johnson JE, Moncalvo JM, Vilgalys R, Redhead SA, Thomas T, McGhee LL (2001) Infrageneric phylogeny of *Collybia* s. str. based on sequences of ribosomal ITS and LSU regions. *Mycological Research* **105**: 164–172.
- Hywel-Jones NL, Sivichai S (1995) *Cordyceps cylindrica* and its association with *Nomuraea atypicola* in Thailand. *Mycological Research* **99**: 809–812.
- Kalele DN, Affokpon A, Coosemans J (2006) Efficacy of *Paecilomyces lilacinus* strain 251 against root knot nematodes in tomato under greenhouse conditions. *Communications in Agricultural and Applied Biological Sciences* **72**: 209–213.
- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* **30**: 3059–3066.
- Katoh K, Toh H (2008) Recent developments in the MAFFT multiple sequence alignment program. *Briefings in Bioinformatics* **9**: 286–298.
- Kepler RM, Ban S, Nakagiri A, Bischoff J, Hywel-Jones N, Owensby CA, Spatafora JW (2013) The phylogenetic placement of hypocrealean insect pathogens in the genus *Polycephalomyces*: an application of One Fungus One Name. *Fungal Biology* **117**: 611–622.
- Kepler RM, Humber RA, Bischoff JF, Rehner SA (2014) Clarification of generic and species boundaries for *Metarhizium* and related fungi through multigene phylogenetics. *Mycologia* doi:10.3852/13-319.

- Kirk PM, Cannon PF, Minter DW, Stalpers JA (eds) (2008) *Ainsworth & Bisby's Dictionary of the Fungi*. 10<sup>th</sup> edn. Wallingford: CAB International.
- Kobayasi Y (1941) The genus *Cordyceps* and its allies. *Science Reports of the Tokyo Bunrika Daigaku*, sect. B, **84** (5): 53–260.
- Leuchtman A, Bacon CW, Schardl CL, White JF jr, Tadych M (2014) Nomenclatural realignment of *Neotyphodium* species with genus *Epichloë*. *Mycologia* doi:10.3852/13-251.
- Li X, Luo H, Zhang K (2005) A new species of *Harposporium* parasitic on nematodes. *Canadian Journal of Botany* **83**: 558–562.
- Luangsa-ard JJ, Houbraken J, van Doorn T, Hong S-B, Borman AM, Hywel-Jones NL, Samson RA (2011b) *Purpureocillium*, a new genus for the medically important *Paecilomyces lilacinus*. *FEMS Microbiology Letters* **321**: 141–149.
- Luangsa-ard JJ, Ridkaew R, Tasanathai K, Thanakitpipattana D, Hywel-Jones NL (2011a) *Ophiocordyceps halabalaensis*: a new species of *Ophiocordyceps* pathogenic to *Camponotus gigas* in Hala Bala Wildlife Sanctuary, Southern Thailand. *Fungal Biology* **115**: 608–614.
- Mains EB (1950) Entomogenous species of *Akanthomyces*, *Hymenostilbe* and *Insecticola* in North America. *Mycologia* **42**: 566–588.
- McNeill J (2012) Guidelines for requests for binding decisions on application of the Code. *Taxon* **61**: 477–478.
- Möller C, Gams W (1993) Two new hyphomycetes isolated from Antarctic lichens. *Mycotaxon* **48**: 441–450.
- Pastor FJ, Guarro J (2006) Clinical manifestations, treatment and outcome of *Paecilomyces lilacinus* infections. *Clinical Microbiology and Infection* **12**: 948–960.
- Patouillard NT (1892) Une Clavariée entomogène. *Revue Mycologique* **14**: 67–70.
- Perdomo H, Cano J, Gené J, García D, Hernández M, Guarro J (2013) Polyphasic analysis of *Purpureocillium lilacinum* isolates from different origins and proposal of the new species *Purpureocillium lavendulum*. *Mycologia* **105**: 151–161.
- Petch T (1931a) Notes on entomogenous fungi. *Transactions of the British Mycological Society* **16**: 55–75.
- Petch T (1931b) New species of *Cordyceps*, collected during the Whitby foray. *The Naturalist, Hull* **1931**: 101–103.
- Petch T (1932) A list of the entomogenous fungi of Great Britain. *Transactions of the British Mycological Society* **17**: 170–178.
- Petch T (1933) Notes on entomogenous fungi. *Transactions of the British Mycological Society* **18**: 48–75.
- Ren SY, Yao YJ (2013) Evaluation of nutritional and physical stress conditions during vegetative growth on conidial production and germination in *Ophiocordyceps sinensis*. *FEMS Microbiology Letters* **346**: 29–35.
- Rodríguez MM, Pastor FJ, Serena C, Guarro J (2010) Efficacy of voriconazole in a murine model of invasive paecilomycosis. *International Journal of Antimicrobial Agents* **35**: 362–365.
- Rossman AY (1978) *Podonectria*, a genus in the *Pleosporales* on scale insects. *Mycotaxon* **7**: 163–182.
- Rossman AY, Seifert KA, Samuels GJ, Minnis AM, Schroers H-J, Lombard L, Crous PW, Pöldmaa K, Cannon PF, Summerbell RC, Geiser DM, Zhuang W, Hirooka Y, Herrera C, Salgado-Salazar C, Chaverri P (2013) Genera in *Bionectriaceae*, *Hypocreaceae*, and *Nectriaceae* (*Hypocreales*) proposed for acceptance or rejection. *IMA Fungus* **4**: 41–51.
- Samson RA, Brady BL (1983) *Paraisaria*, a new genus for *Isaria dubia*, the anamorph of *Cordyceps gracilis*. *Transactions of the British Mycological Society* **81**: 285–290.
- Samson RA, Soares GG jr (1984) Entomopathogenic species of the hyphomycete genus *Tolypocladium*. *Journal of Invertebrate Pathology* **43**: 133–139.
- Seifert KA (1985) A monograph of *Stilbella* and some allied hyphomycetes. *Studies in Mycology* **27**: 1–234.
- Shimazu M, Glockling SL (1997) A new species of *Harposporium* with two spore types isolated from the larva of a cerambycid beetle. *Mycological Research* **101**: 1371–1376.
- Shrestha B, Bawa KS (2013) Trade, harvest, and conservation of caterpillar fungus (*Ophiocordyceps sinensis*) in the Himalayas. *Biological Conservation* **159**: 514–520.
- Shrestha B, Zhang WM, Zhang YJ, Liu XZ (2010) What is the Chinese caterpillar fungus *Ophiocordyceps sinensis* (*Ophiocordycipitaceae*)? *Mycology* **1**: 228–236.
- Speare AT (1917) *Sorosporella uvella* and its occurrence in cutworms in America. *Journal of Agricultural Research* **8**: 189–194.
- Speare AT (1920) Further studies of *Sorosporella uvella*, a fungous parasite of noctuid larvae. *Journal of Agricultural Research* **18**: 399–439.
- Stalpers JA, Seifert KA, Samson RA (1991) A revision of the genera *Antromycopsis*, *Sclerostilbum*, and *Tilachlidiosis* (hyphomycetes). *Canadian Journal of Botany* **69**: 6–15.
- Stamatakis A (2006) RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**: 2688–2690.
- Stone R (2008) Last stand for the body snatcher of the Himalayas? *Science* **322**: 1182.
- Sung G-H, Spatafora JW, Zare R, Hodge KT, Gams W (2001) A revision of *Verticillium* sect. *Prostrata*. II. Phylogenetic analyses of SSU and LSU nuclear rDNA sequences from anamorphs and teleomorphs of the *Clavicipitaceae*. *Nova Hedwigia* **72**: 311–328.
- Sung G-H, Hywel-Jones NL, Sung J-M, Luangsa-ard JJ, Shrestha B, Spatafora JW (2007) Phylogenetic classification of *Cordyceps* and the clavicipitaceous fungi. *Studies in Mycology* **57**: 5–59.
- Survase SA, Kagliwal LD, Annapure US, Singhal RS (2011) Cyclosporin A - A review on fermentative production, downstream processing and pharmacological applications. *Biotechnology Advances* **29**: 418–435.
- Talavera G, Castresana J (2007) Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Systematic Biology* **56**: 564–577.
- Traber R, Dreyfuss MM (1996) Occurrence of cyclosporins and cyclosporin-like peptolides in fungi. *Journal of Industrial Microbiology* **17**: 397–401.
- VUILLEMIN P (1896) Les Hypostomacées, nouvelle famille de champignons parasites. *Bulletin de la Société des Sciences de Nancy*, sér. 2 **14**: 15–67.
- Weiser J, Matha V, Jegorov A (1991) *Tolypocladium terricola* sp. n., a new mosquito-killing species of the genus *Tolypocladium* Gams (hyphomycetes). *Folia Parasitologica* **38**: 363–369.
- Zare R, Gams W, Culham A (2000) A revision of *Verticillium* sect. *Prostrata*. I. Phylogenetic studies using ITS sequences. *Nova Hedwigia* **71**: 465–480.
- Zare R, Gams W (2001) A revision of *Verticillium* section *Prostrata*. VI. The genus *Haptocillium*. *Nova Hedwigia* **73**: 271–292.

**Table 1.** Specimen information and GenBank accession numbers for sequences used in this study.

Species	Voucher Information	SSU	LSU	TEF	RPB1	RPB2
<i>Chaunopycnis alba</i>	MRL GB5502 MRL MF6799		AF245297 AF373284			
<i>Chaunopycnis pustulata</i>	MRL GB6597 MRL MF5368LR		AF389190 AF373282			
<i>Cordyceps cylindrica</i>	CEM 1185	KJ878907	KJ878872	KJ878955		
<i>Cordyceps formosana</i>	TNM F13893	KJ878908		KJ878956	KJ878988	KJ878943
<i>Cordyceps gunnii</i>	OSC 76404	AF339572	AF339522	AY489616	AY489650	DQ522426
<i>Cordyceps irangiensis</i>	OSC 128579	EF469123	EF469076	EF469060	EF469089	EF469107
<i>Cordyceps nipponica</i>	BCC 18108	KF049608	KF049626	KF049681	KF049644	
<i>Cordyceps pleuric capitata</i>	NBRC 100745	KF049606	KF049624	KF049679	KF049642	KF049667
<i>Cordyceps pleuric capitata</i>	NBRC 100746	KF049607	KF049625	KF049680	KF049643	KF049668
<i>Cordyceps</i> sp.	EFCC 12075	KJ878909	KJ878873	KJ878957	KJ878989	
<i>Drechmeria coniospora</i>	CBS 596.92	AF106012				
<i>Elaphocordyceps capitata</i>	OSC 71233	AY489689	AY489721	AY489615	AY489649	DQ522421
<i>Elaphocordyceps fracta</i>	OSC 110990	DQ522545	DQ518759	DQ522328	DQ522373	DQ522425
<i>Elaphocordyceps japonica</i>	OSC 110991	DQ522547	DQ518761	DQ522330	DQ522375	DQ522428
<i>Elaphocordyceps longisegmentis</i>	OSC 110992		EF468816		EF468864	EF468919
<i>Elaphocordyceps ophioglossoides</i>	CBS 100239 OSC 106405	KJ878910 AY489691	KJ878874 AY489723	KJ878958 AY489618	KJ878990 AY489652	KJ878944 DQ522429
<i>Elaphocordyceps subsessilis</i>	OSC 71235	EF469124	EF469077	EF469061	EF469090	EF469108
<i>Haptocillium balanooides</i>	CBS 250.82	AF339588	AF339539	DQ522342	DQ522388	DQ522442
<i>Haptocillium sinense</i>	CBS 567.95	AF339594	AF339545	DQ522343	DQ522389	DQ522443
<i>Haptocillium zeosporum</i>	CBS 335.8	AF339589	AF339540	EF469062	EF469091	EF469109
<i>Harposporium anguillulae</i>	ARSEF 5407 ARSEF 5593		AY636080 AY636081			
<i>Harposporium helicooides</i>	ARSEF 5354	AF339577	AF339527			
<i>Hirsutella crinalis</i>	TNS F18550	KJ878911	KJ878875	KJ878959		
<i>Hirsutella</i> sp.	OSC 128575	EF469126	EF469079	EF469064	EF469093	EF469110
<i>Hirsutella</i> sp.	NHJ 12525	EF469125	EF469078	EF469063	EF469092	EF469111
<i>Hymenostilbe aurantiaca</i>	OSC 128578	DQ522556	DQ518770	DQ522345	DQ522391	DQ522445
<i>Hymenostilbe muscaria</i>	OSC 151902	KJ878912	KJ878876		KJ878991	KJ878945
<i>Hymenostilbe odonatae</i>	TNS F18563 TNS F27117		KJ878877 KJ878878		KJ878992	
<i>Isaria takamizuensis</i>	NHJ 3582	EU369097	EU369034	EU369015		
<i>Isaria takamizusanensis</i>	NHJ 3497	EU369096	EU369033	EU369014	EU369053	EU369074
<i>Nomuraea atypicola</i>	RCEF 3833 OSC 151901	KJ878913 KJ878914	KJ878879 KJ878880	KJ878960 KJ878961	KJ878993 KJ878994	
<i>Ophiocordyceps aciculalis</i>	CBS 744.73 OSC 110987 OSC 110988	EF468987 EF468950 EF468951	EF468841 EF468805 EF468804	EF468786 EF468744 EF468745	EF468892 EF468852 EF468853	
<i>Ophiocordyceps agriotidis</i>	ARSEF 5692	DQ522540	DQ518754	DQ522322	DQ522368	DQ522418
<i>Ophiocordyceps annulata</i>	CEM 303	KJ878915	KJ878881	KJ878962	KJ878995	
<i>Ophiocordyceps aphodii</i>	ARSEF 5498	DQ522541	DQ518755	DQ522323		DQ522419
<i>Ophiocordyceps brunneipunctata</i>	OSC 128576	DQ522542	DQ518756	DQ522324	DQ522369	DQ522420
<i>Ophiocordyceps clavata</i>	CEM 1762 NBRC 106961	KJ878916 JN941727	KJ878882 JN941726	KJ878963 JN941414	KJ878996 JN992461	
	CEM 1763 NBRC 106962		KJ878883 JN941415	KJ878964 JN992460	KJ878997	

**Table 1.** (Continued).

Species	Voucher Information	SSU	LSU	TEF	RPB1	RPB2
<i>Ophiocordyceps communis</i>	NHJ 12581	EF468973	EF468831	EF468775		EF468930
	NHJ 12582	EF468975	EF468830	EF468771		EF468926
<i>Ophiocordyceps curculionum</i>	OSC 151910	KJ878918	KJ878885		KJ878999	
<i>Ophiocordyceps dipterigena</i>	OSC 151911	KJ878919	KJ878886	KJ878966	KJ879000	
	OSC 151912	KJ878920	KJ878887	KJ878967	KJ879001	
<i>Ophiocordyceps elongata</i>	OSC 110989		EF468808	EF468748	EF468856	
<i>Ophiocordyceps entomorrhiza</i>	KEW 53484	EF468954	EF468809	EF468749	EF468857	EF468911
<i>Ophiocordyceps formicarum</i>	TNS F18565	KJ878921	KJ878888	KJ878968	KJ879002	KJ878946
<i>Ophiocordyceps forquignonii</i>	OSC 151908	KJ878922	KJ878889		KJ879003	KJ878947
<i>Ophiocordyceps gracilis</i>	EFCC 3101	EF468955	EF468810	EF468750	EF468858	EF468913
	EFCC 8572	EF468956	EF468811	EF468751	EF468859	EF468912
	OSC 151906	KJ878923	KJ878890	KJ878969		
<i>Ophiocordyceps heteropoda</i>	EFCC 10125	EF468957	EF468812	EF468752	EF468860	EF468914
	OSC 106404	AY489690	AY489722	AY489617	AY489651	
<i>Ophiocordyceps irangiensis</i>	OSC 128577	DQ522546	DQ518760	DQ522329	DQ522374	DQ522427
<i>Ophiocordyceps konnoana</i>	EFCC 7295	EF468958			EF468862	EF468915
<i>Ophiocordyceps konnoana</i>	EFCC 7315	EF468959		EF468753	EF468861	EF468916
<i>Ophiocordyceps lloydii</i>	OSC 151913	KJ878924	KJ878891	KJ878970	KJ879004	KJ878948
<i>Ophiocordyceps longissima</i>	EFCC 6814		EF468817	EF468757	EF468865	
	TNS F18448	KJ878925	KJ878892	KJ878971	KJ879005	
<i>Ophiocordyceps longissima</i>	HMAS_199600	KJ878926		KJ878972	KJ879006	KJ878949
<i>Ophiocordyceps melolonthae</i>	OSC 110993	DQ522548	DQ518762	DQ522331	DQ522376	
<i>Ophiocordyceps myrmecophila</i>	HMAS_199620	KJ878929	KJ878895	KJ878975	KJ879009	
	CEM 1710	KJ878927	KJ878893	KJ878973	KJ879007	
	TNS 27120	KJ878928	KJ878894	KJ878974	KJ879008	
<i>Ophiocordyceps neovolkiana</i>	OSC 151903	KJ878930	KJ878896	KJ878976	KJ879010	
<i>Ophiocordyceps nigrella</i>	EFCC 9247	EF468963	EF468818	EF468758	EF468866	EF468920
<i>Ophiocordyceps nutans</i>	OSC 110994	DQ522549	DQ518763	DQ522333	DQ522378	
<i>Ophiocordyceps pruinosa</i>	NHJ 12994	EU369106	EU369041	EU369024	EU369063	EU369084
<i>Ophiocordyceps pulvinata</i>	TNS-F 30044	GU904208		GU904209	GU904210	
<i>Ophiocordyceps purpureostromata</i>	TNS F18430	KJ878931	KJ878897	KJ878977	KJ879011	
<i>Ophiocordyceps ravenelii</i>	OSC 110995	DQ522550	DQ518764	DQ522334	DQ522379	DQ522430
	OSC 151914	KJ878932		KJ878978	KJ879012	KJ878950
<i>Ophiocordyceps rhizoidea</i>	NHJ 12522	EF468970	EF468825	EF468764	EF468873	EF468923
	NHJ 12529	EF468969	EF468824	EF468765	EF468872	EF468922
<i>Ophiocordyceps ryogamiensis</i>	NBRC 101751	KF049614	KF049633	KF049688	KF049650	
<i>Ophiocordyceps sinensis</i>	EFCC 7287	EF468971	EF468827	EF468767	EF468874	EF468924
<i>Ophiocordyceps sobolifera</i>	KEW 78842	EF468972	EF468828		EF468875	EF468925
	TNS F18521	KJ878933	KJ878898	KJ878979	KJ879013	
<i>Ophiocordyceps</i> sp.	TNS F18495	KJ878937	KJ878901		KJ879017	
<i>Ophiocordyceps</i> sp.	OSC 110997	EF468976		EF468774	EF468879	EF468929
<i>Ophiocordyceps</i> sp.	OSC 151904	KJ878934	KJ878899	KJ878980	KJ879014	
<i>Ophiocordyceps</i> sp.	OSC 151905	KJ878935		KJ878981	KJ879015	KJ878951
<i>Ophiocordyceps</i> sp.	OSC 151909	KJ878936	KJ878900	KJ878982	KJ879016	KJ878952
<i>Ophiocordyceps</i> sphecocephala	OSC 110998	DQ522551	DQ518765	DQ522336	DQ522381	DQ522432
<i>Ophiocordyceps</i> stylophora	OSC 110999	EF468982	EF468837	EF468777	EF468882	EF468931
	OSC 111000	DQ522552	DQ518766	DQ522337	DQ522382	DQ522433
<i>Ophiocordyceps</i> tricentri	CEM 160	AB027330	AB027376			
<i>Ophiocordyceps</i> unilateralis	OSC 128574	DQ522554	DQ518768	DQ522339	DQ522385	DQ522436

**Table 1.** (Continued).

Species	Voucher Information	SSU	LSU	TEF	RPB1	RPB2
<i>Ophiocordyceps variabilis</i>	OSC 111003 ARSEF 5365	EF468985 DQ522555	EF468839 DQ518769	EF468779 DQ522340	EF468885 DQ522386	EF468933 DQ522437
<i>Ophiocordyceps yakusimensis</i>	HMAS_199604	KJ878938	KJ878902		KJ879018	KJ878953
<i>Paecilomyces lilacinus</i>	ARSEF 2181 CBS 431.87 CBS 284.36	AF339583 AY624188 AY624189	AF339534 EF468844 AY624227	EF468790 EF468791 EF468792	EF468896 EF468897 EF468898	EF468940 EF468941
<i>Podocrella harposporifera</i>	ARSEF 5472	AF339569	AF339519	DQ118747	DQ127238	
<i>Podonectria citrina</i>	TNS F18537		KJ878903	KJ878983		KJ878954
<i>Polycephalomyces cuboideus</i>	TNS F18487	KF049609	KF049628	KF049683		
<i>Polycephalomyces cuboideus</i>	NBRC 101740	KF049610	KF049629	KF049684	KF049646	
<i>Polycephalomyces formosus</i>	ARSEF 1424	KF049615	AY259544	DQ118754	DQ127245	KF049671
<i>Polycephalomyces nipponicus</i>	BCC 1881 BCC 1682 NHJ4286 BCC2325	KF049618 KF049620 KF049621 KF049622	KF049636 KF049638 KF049639 KF049640	KF049692 KF049694 KF049695 KF049696		KF049674 KF049676 KF049655 KF049677
<i>Polycephalomyces paracuboideus</i>	NBRC 101742	KF049611	KF049630	KF049685	KF049647	KF049669
<i>Polycephalomyces prolificus</i>	TNS F18481 TNS F18547	KF049612 KF0496613	KF049631 KF049632	KF049686 KF049687	KF049648 KF049649	KF049670
<i>Polycephalomyces ramosopulvinatus</i>	SU-65 EFCC 5566		DQ118742 KF049627	DQ118753 KF049682	DQ127244 KF049645	
<i>Polycephalomyces</i> sp.	JB07.08.16_08	KF049616	KF049635	KF049690	KF049652	KF049672
<i>Polycephalomyces</i> sp.	JB07.08.17_07b	KF049617		KF049691	KF049653	KF049673
<i>Polycephalomyces</i> sp.	BBC 2637	KF049619	KF049637	KF049693		KF049675
<i>Polycephalomyces tomentosus</i>	BL4	KF049623	AY259545	KF049697	KF049656	KF049678
<i>Stilbella buquetii</i>	HMAS_199613 HMAS_199617	KJ878939 KJ878940	KJ878904 KJ878905	KJ878984 KJ878985	KJ879019 KJ879020	
<i>Tilachlidiopsis nigra</i>	TNS 16252 TNS 16250	KJ878941 KJ878942		KJ878986 KJ878987		KJ879021
<i>Tolypocladium cylindrosporum</i>	NRRL 28025	AF049153	AF049173			

**Table 2.** Proposed list of generic names in Ophiocordycipitaceae to be protected and their competing synonyms. Names to be protected are in bold type, and names previously synonymized are in blue.

Proposed to protect	Proposed to suppress
<b><i>Ophiocordyceps</i></b> Patch, Trans. Br. Mycol. Soc. <b>16</b> : 74 (1931). Type: <i>O. blattae</i> Patch 1931.	<i>Sorosporella</i> Sorokin Zentbl. Bakt. Parasitkde., Abt. II <b>4</b> : 644 (1888). Type: <i>S. agrotidis</i> Sorokin 1888.
	Hirsutella Pat., Revue Mycol. <b>14</b> : 67 (1892). Type: <i>H. entomophila</i> Pat. 1892.
	Didymobotryopsis Henn., Hedwigia <b>41</b> : 149 (1902). Type: <i>D. parasitica</i> Henn. 1902.
	Mahevia Lagarde, Archs Zool. Exp. Gen. <b>56</b> : 292 (1917). Type: <i>M. guignardii</i> (Maheu) Lagarde 1917.
	Synnematium Speare, Mycologia <b>12</b> : 74 (1920). Type: <i>S. jonesii</i> Speare 1920.

**Table 2.** (Continued).

Proposed to protect	Proposed to suppress
	<i>Trichosterigma</i> Petch, <i>Trans. Br. Mycol. Soc.</i> <b>8</b> : 215 (1923). Type: <i>T. clavisporum</i> Petch 1923.
	<i>Didymobotrys</i> Clem. & Shear, <i>Gen. Fungi</i> : 228 (1931). Type: <i>D. parasitica</i> (Henn.) Clem. & Shear 1931.
	<i>Troglobiomyces</i> Pacioni, <i>Trans. Br. Mycol. Soc.</i> <b>74</b> : 244 (1980). Type: <i>T. guignardii</i> (Maheu) Pacioni 1980.
	<i>Hymenostilbe</i> Petch, <i>Naturalist (Hull)</i> , ser. 3, <b>1931</b> : 101 (1931). Type: <i>H. muscaria</i> Petch 1931.
	<i>Syngliocladium</i> Petch, <i>Trans. Br. Mycol. Soc.</i> <b>17</b> : 177 (1932). Type: <i>S. aranearium</i> Petch 1932.
	<i>Cordycepioideus</i> Stifler, <i>Mycologia</i> <b>33</b> : 83 (1941). Type: <i>C. bisporus</i> Stifler 1941.
	<i>Paraisaria</i> Samson & B.L. Brady, <i>Trans. Br. Mycol. Soc.</i> <b>81</b> : 285 (1983). Type: <i>P. dubia</i> (Delacr.) Samson & B.L. Brady 1983.
<b><i>Purpureocillium</i></b> Luangsa-ard et al., <i>FEMS Microbiol Lett</i> <b>321</b> : 144 (2011). Type: <i>P. lilacinum</i> (Thom) Luangsa-ard et al. 2011 (syn. <i>Penicillium lilacinum</i> Thom 1920).	
<b><i>Tolypocladium</i></b> W. Gams, <i>Persoonia</i> <b>6</b> : 185 (1971). Type: <i>T. inflatum</i> W. Gams 1971.	<i>Chaunopycnis</i> W. Gams, <i>Persoonia</i> <b>11</b> : 75 (1980). Type: <i>C. alba</i> W. Gams 1980. <i>Elaphocordyceps</i> G.H. Sung & Spatafora, <i>Stud. Mycol.</i> <b>57</b> : 36 (2007). Type: <i>E. ophioglossoides</i> (Ehrh. ex J.F. Gmel. : Fr.) G.H. Sung et al. 2007.
<b><i>Harposporium</i></b> Lohde, <i>Tagbl. Versamml. Ges. Deutsch. Naturf.</i> <b>47</b> : 206 (1874). Type: <i>H. anguillulae</i> Lohde 1874.	<i>Polyrhina</i> Sorokin, <i>Annls Sci. Nat., Bot., sér 6, 4</i> : 65 (1876). Type: <i>P. multiformis</i> Sorokin 1876.
	<i>Podocrella</i> Seaver, <i>Mycologia</i> <b>20</b> : 57 (1928). Type: <i>P. poronioides</i> Seaver 1928.
	<i>Atricordyceps</i> Samuels, <i>N.Z. Jl. Bot.</i> <b>21</b> : 174 (1983). Type <i>A. harposporifera</i> Samuels 1983.
<b><i>Drechmeria</i></b> W. Gams & H.-B. Jansson, <i>Mycotaxon</i> <b>22</b> : 36 (1985). Type: <i>D. coniospora</i> (Drechsler) W. Gams & H.-B. Jansson 1985 (syn. <i>Meria coniospora</i> Drechsler 1941).	<i>Haptocillium</i> W. Gams & Zare, <i>Nova Hedwigia</i> <b>73</b> : 334 (2001). Type: <i>H. balanoides</i> (Drechsler) Zare & W. Gams 2001.
<b><i>Polycephalomyces</i></b> Kobayasi, <i>Sci. Rep. Tokyo Bunrika Daig.</i> , sect. B <b>5</b> : 245 (1941). Type: <i>P. formosus</i> Kobayasi 1941.	<i>Blistum</i> B. Sutton, <i>Mycol. Pap.</i> <b>132</b> : 16 (1973). Type: <i>B. tomentosum</i> (Schrad.) B. Sutton 1973.