

Septin 2/6/7 complexes tune microtubule plus-end growth and EB1 binding in a concentration- and filament-dependent manner

Konstantinos Nakos, Megan R. Radler, and Elias T. Spiliotis*

Department of Biology, Drexel University, Philadelphia, PA 19104

ABSTRACT Septins (SEPTs) are filamentous guanosine-5'-triphosphate (GTP)-binding proteins, which affect microtubule (MT)-dependent functions including membrane trafficking and cell division, but their precise role in MT dynamics is poorly understood. Here, in vitro reconstitution of MT dynamics with SEPT2/6/7, the minimal subunits of septin heteromers, shows that SEPT2/6/7 has a biphasic concentration-dependent effect on MT growth. Lower concentrations of SEPT2/6/7 enhance MT plus-end growth and elongation, while higher and intermediate concentrations inhibit and pause plus-end growth, respectively. We show that SEPT2/6/7 has a modest preference for GTP- over guanosine diphosphate (GDP)-bound MT lattice and competes with end-binding protein 1 (EB1) for binding to guanosine 5'-O-[γ -thio] triphosphate (GTP γ S)-stabilized MTs, which mimic the EB1-preferred GDP-Pi state of polymerized tubulin. Strikingly, SEPT2/6/7 triggers EB1 dissociation from plus-end tips in *cis* by binding to the MT lattice and in *trans* when MT plus ends collide with SEPT2/6/7 filaments. At these intersections, SEPT2/6/7 filaments were more potent barriers than actin filaments in pausing MT growth and dissociating EB1 in vitro and in live cells. These data demonstrate that SEPT2/6/7 complexes and filaments can directly impact MT plus-end growth and the tracking of plus end-binding proteins and thereby may facilitate the capture of MT plus ends at intracellular sites of septin enrichment.

Monitoring Editor

Manuel Théry
CEA, Hôpital Saint-Louis

Received: Jul 8, 2019

Revised: Sep 23, 2019

Accepted: Sep 25, 2019

INTRODUCTION

Septins (SEPTs) are guanosine-5'-triphosphate (GTP)-binding proteins that assemble into filamentous higher-order oligomers and polymers comprising a major component of the mammalian cytoskeleton alongside actin, microtubules (MTs), and intermediate filaments (Mostowy and Cossart, 2012; Spiliotis, 2018). Septins associate with MTs and affect MT organization and dynamics in various

cell types (Kremer *et al.*, 2005; Bowen *et al.*, 2011; Ageta-Ishihara *et al.*, 2013; Bai *et al.*, 2013; Shindo *et al.*, 2018). Notably, septins are functionally involved in MT-dependent functions such as cell division and polarized membrane traffic (Spiliotis *et al.*, 2005, 2008; Estey *et al.*, 2010; Menon *et al.*, 2014; Karasmanis *et al.*, 2018). A growing number of studies suggests that septins are involved in the entry and growth of MTs in membrane protrusions (e.g., filopodia, cilia, and microtentacles) (Ghossoub *et al.*, 2013; Nolke *et al.*, 2016; Ostevold *et al.*, 2017), but how septins affect the dynamic instability of MTs is not well understood.

Mammalian septins are a family of 13 paralogous genes, which are classified in four groups named after SEPT2, SEPT3, SEPT6, and SEPT7 (Kinoshita, 2003b; Weirich *et al.*, 2008; Russell and Hall, 2011). The minimal building block of septin heteromers is a palindromic hetero-hexamers that contains septins from the SEPT2, SEPT6, and SEPT7 groups in a stoichiometry of 2:2:2, which can further expand into a hetero-octamer on the addition of a septin paralogue from the SEPT3 group (Kinoshita, 2003a; Sheffield *et al.*, 2003; Sirajuddin *et al.*, 2007; Kim *et al.*, 2011; Sellin *et al.*, 2011). Depending on the relative expression of septin paralogues, which differs among cell types, subunits of septin heteromers have been

This article was published online ahead of print in MBoC in Press (<http://www.molbiolcell.org/cgi/doi/10.1091/mbc.E19-07-0362>) on October 2, 2019.

The authors have no conflicts of interest to declare.

*Address correspondence to: Elias Spiliotis (ets33@drexel.edu).

Abbreviations used: BSA, bovine serum albumin; DIV, days in vitro; EB1, end-binding protein 1; EB3, end-binding protein 3; GDP, guanosine diphosphate; GMPCPP, guanosine-5'-[(α,β)-methylene]triphosphate; GTP, guanosine-5'-triphosphate; GTP γ S, guanosine 5'-O-[γ -thio]triphosphate; MAP, microtubule-associated protein; MAP4, microtubule-associated protein 4; MT, microtubule; SEPT, septin; TIRF, total internal reflection fluorescence.

© 2019 Nakos *et al.* This article is distributed by The American Society for Cell Biology under license from the author(s). Two months after publication it is available to the public under an Attribution-NonCommercial-Share Alike 3.0 Unported Creative Commons License (<http://creativecommons.org/licenses/by-nc-sa/3.0>).

"ASCB®," "The American Society for Cell Biology®," and "Molecular Biology of the Cell®" are registered trademarks of The American Society for Cell Biology.

reported to colocalize with MTs (Spiliotis *et al.*, 2008, 2010; Sellin *et al.*, 2012, 2014; Spiliotis, 2018; Targa *et al.*, 2019). SEPT9 binds MTs directly through N-terminal repeat motifs (K/R-R/x-x-D/E), which associate with the C-terminal tails of β -tubulin (Bai *et al.*, 2013), but it is unclear whether these motifs function similarly for septins of the SEPT6 and SEPT7 groups, which also associate with MTs; SEPT2 does not appear to bind MTs directly (Bai *et al.*, 2013).

How septins, as individual subunits or higher-order complexes, impact the dynamic properties of MTs has been difficult to determine owing to disparate results from septin knockdown approaches. SEPT7 was initially posited to enhance MT dynamics, because SEPT7 knockdown increased MT acetylation, a marker of MT stability (Kremer *et al.*, 2005; Janke and Montagnac, 2017). However, this effect was attributed to cytosolic SEPT7, which sequestered the MT-associated protein 4 (MAP4) away from MTs (Kremer *et al.*, 2005) and promoted the interaction of histone deacetylase HDAC6 with cytosolic free tubulin (Ageta-Ishihara *et al.*, 2013). Comprehensive analysis of MT dynamics in SEPT2-depleted Madin–Darby canine kidney cells revealed a distinct increase in MT catastrophe, indicating that septins promote persistent MT growth and thereby may promote MT growth (Bowen *et al.*, 2011). More recently, septins were proposed to pro-

mote the formation of MT-driven membrane protrusions through an interplay with the MT plus-end tracking protein end-binding protein 1 (EB1) in cells that are infected with a *Clostridium difficile* toxin (Nolke *et al.*, 2016). Despite these findings, it remains unclear how precisely septins regulate MT dynamics. To further our understanding, we reconstituted MT plus-end and EB1 dynamics in vitro with soluble septin 2/6/7 (SEPT2/6/7) complexes and SEPT2/6/7 filaments.

RESULTS

SEPT2/6/7 tunes MT plus-end dynamics in a concentration-dependent manner

Using a TIRF microscopy assay that visualizes the polymerization and depolymerization of Rhodamine-labeled tubulin onto stable HiLyte 488-labeled MT seeds, we tested how MT dynamics are impacted by different concentrations of recombinant SEPT2/6/7 complexes, which were purified from bacteria. Quantification of the parameters of MT dynamics revealed a biphasic concentration-dependent response for catastrophe frequencies and MT lengths (Figure 1, A and B; Supplemental Movies S1–S3). Lower concentrations of SEPT2/6/7 (10–400 nM) reduced catastrophe frequencies by 20–34% and increased MT lengths by 1.2- to 1.8-fold, whereas




FIGURE 1: SEPT2/6/7 tunes MT dynamics in a concentration-dependent manner. (A–D) Dot plots show mean (\pm SEM) frequency of MT plus-end catastrophe (A) and length (B), and rates of MT plus-end polymerization (C) and depolymerization (D) with SEPT2/6/7 at concentrations of 0 nM ($n = 69$), 10 nM ($n = 65$), 100 nM ($n = 51$), 200 nM ($n = 61$), 400 nM ($n = 66$), 800 nM ($n = 50$), 1 μ M ($n = 51$), 2 μ M ($n = 49$), and 4 μ M ($n = 44$). (E–M) Kymographs show representative MT plus-end dynamics (red) on nucleation from MT seeds (green) in the presence of 0 nM (E), 10 nM (F), 100 nM (G), 200 nM (H), 400 nM (I), 800 nM (J), 1 μ M (K), 2 μ M (L), and 4 μ M (M) of SEPT2/6/7; ns, nonsignificant ($p > 0.05$). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

higher concentrations of SEPT2/6/7 (>1 μM) had the opposite effect, enhancing catastrophe frequencies by 1.2- to 1.6-fold and decreasing MT lengths by 30–48%. Rates of MT growth remained similar up to 800 nM of SEPT2/6/7 and began to decrease at higher concentrations (Figure 1C). Depolymerization rates trended toward reduction with increased concentrations of SEPT2/6/7 and decreased significantly at SEPT2/6/7 concentrations >400 nM (Figure 1D). As also evidenced by the kymographs of MT dynamics (Figure 1, E–M), these data show that lower concentrations of SEPT2/6/7 induce persistent MT growth by decreasing the frequency of catastrophe events and rates of depolymerization. However, micromolar concentrations of SEPT2/6/7 dampen MT dynamics and reduce MT lengths by decreasing rates of MT growth and enhancing catastrophe.

Because SEPT2/6/7 complexes assemble into higher-order filamentous structures (Supplemental Figure S1A), which depend on concentration and ionic conditions, we reasoned that the biphasic effect might be due to concentration-dependent differences in the assembly state of SEPT2/6/7 complexes. To assess the higher-order state of SEPT2/6/7, we performed a sedimentation assay after diluting and incubating SEPT2/6/7 complexes in the buffer of the MT dynamics assay. Quantifications of SEPT2/6/7 in the pellet and supernatant fractions showed that the levels of sedimenting higher-order septins were fairly equivalent up to concentrations of 400 nM, but increased in a stepwise concentration-dependent manner above 800 nM (Supplemental Figure S1, B and C). This increase in the amount of higher-order SEPT2/6/7 correlated with the dampening of MT dynamics observed at concentrations >800 nM (Figure 1). Hence, higher-order fibrillar SEPT2/6/7 does not appear to favor MT growth and elongation, which are enhanced by lower concentrations of SEPT2/6/7.

Given that MT growth and dynamics are inhibited by higher concentrations of SEPT2/6/7 (> 800 nM), we reasoned that higher-order SEPT2/6/7 may associate with free unpolymerized tubulin, lowering the effective concentration of tubulin that is available to polymerize. This is plausible as we recently found that SEPT9 binds and recruits free tubulin to the MT lattice (Nakos *et al.*, 2019). To test this possibility, SEPT2/6/7 and SEPT9 complexes were immobilized on functionalized glass chambers and fluorescent tubulin was added in the buffer conditions of the MT dynamics assay (Supplemental Figure S2A). Binding of soluble tubulin to septin complexes was visualized by TIRF imaging similar to a previous assay of MT nucleation (Lazarus *et al.*, 2013). While tubulin was increasingly bound to chambers with submicromolar concentrations of SEPT9, no binding was observed for a similar concentration range of SEPT2/6/7 (Supplemental Figure S2, B and C). These data indicate that SEPT2/6/7 does not sequester unpolymerized tubulin and the effects of SEPT2/6/7 on MT dynamics depend on the septin concentration and assembly state (oligomers vs. higher-order polymers).

SEPT2/6/7 pauses MT plus-end growth

In kymographs of MT dynamics, we noticed that MT plus ends exhibited a variety of pausing events in the presence of SEPT2/6/7 complexes. In several kymographs, MT plus ends paused transiently while growing and subsequently depolymerized or continued to grow before undergoing catastrophe (arrowheads in Figure 2, A, B, and D; Supplemental Movie S4). Although much less frequent, MT plus ends also paused transiently during catastrophe before depolymerizing to the MT seed (Figure 2, C, D, and G). Additionally, the slope of depolymerization events was shallower, which was consistent with decreased rates of depolymerization (Figure 1).

Given the concentration dependence of SEPT2/6/7 effects on MT dynamics, we analyzed the frequency of pausing events that occurred in growth phases. Strikingly, MT pausing increased with increasing concentrations of SEPT2/6/7 peaking at 400 nM, where the percentage of MTs with pausing events doubled from 23% to 48% (Figure 2E). This pausing effect was unique to SEPT2/6/7 as SEPT9 did not cause a similar effect (Figure 2F). However, at SEPT2/6/7 concentrations higher than 400 nM, MT pausing began to decrease and micromolar concentrations of SEPT2/6/7 did not increase pausing above the levels observed in the absence of SEPT2/6/7. Hence, MT pausing appears to be an intermediate phenotype that occurs by SEPT2/6/7 complexes in between concentrations that promote and inhibit MT growth and elongation.

SEPT2/6/7 associates with the MT lattice and plus ends

On the basis of the effects of SEPT2/6/7 on MT plus-end growth and pausing, we sought to examine whether SEPT2/6/7 complexes associate with the lattice and/or tips of dynamic MT plus ends. At low concentrations (10–100 nM), mCherry-SEPT2/6/7 localized sparsely and in patches on the lattice of MT seeds and growing plus ends and occasionally at the tips of plus ends (Figure 3A). In kymographs of time-lapse imaging, we observed binding of mCherry-SEPT2/6/7 to the lattice of the MT plus-end segments, which persisted for several seconds to a few minutes, and to the tips of polymerizing and depolymerizing plus ends (Figure 3A). Because of the sparse localization, we quantified MTs categorically based on the distribution of mCherry-SEPT2/6/7 (Figure 3B). At 10 nM, mCherry-SEPT2/6/7 localized to the lattice of the seed and/or plus-end segments of most MTs (~80%) and the tips of plus ends on 27% of total MTs. At 100 nM, however, binding of mCherry-SEPT2/6/7 to plus-end tips was rare, and SEPT2/6/7 associated exclusively with the lattice of MT seeds (65%) or with both the lattice of MT seeds and plus-end segments (Figure 3B). At concentrations >100 nM, mCherry-SEPT2/6/7 decorated MTs more extensively (Figure 3C). Quantification of mCherry-SEPT2/6/7 fluorescence on the lattice of MT seeds versus plus ends showed a 1.5-fold preference for the guanosine-5'-[(α,β)-methylene]triphosphate (GMPCPP)-stabilized lattice (Figure 3, C and D), which is significantly less than the ~4-fold difference that was previously observed for SEPT9, which in similar concentrations (400–800 nM) did not fully decorate the lattice segments of MT plus ends (Nakos *et al.*, 2019). Collectively, our results show that SEPT2/6/7 has a higher affinity than SEPT9 for the guanosine diphosphate (GDP)-bound lattice of MT plus-end segments.

SEPT2/6/7 complexes inhibit MT plus-end binding and tracking of EB1

Recent reports of an interaction between EB1 and septins (Nolke *et al.*, 2016) raises the question of whether EB1 and SEPT2/6/7 cooperate or compete with one another on MT plus ends. Given the effects of SEPT2/6/7 on MT plus ends, we sought to test whether SEPT2/6/7 affects the binding and dynamics of the plus end-binding protein EB1, which can also provide more insight into how SEPT2/6/7 impacts MT plus-end dynamics.

We reconstituted MT dynamics in the presence of recombinant EB1-GFP and mCherry-SEPT2/6/7 (Figure 4A). Tracking of EB1-GFP on MT plus ends required buffer conditions (50 mM KCl), which limit diffusive interactions of EB1-GFP with the MT lattice (Zanic *et al.*, 2009). In these ionic conditions, the MT-binding affinity of mCherry-SEPT2/6/7 was reduced, but mCherry-SEPT2/6/7 exhibited the same concentration-dependent pattern of decoration with a modest preference for GMPCPP-bound MT seeds (Figure 4A).





FIGURE 2: SEPT2/6/7 pauses MT plus-end growth. (A–C) Kymographs show pausing events during MT growth or shrinkage in the presence of 0 nM (A), 400 nM (B and C, left) and 800 nM (B and C, right) of SEPT2/6/7. (B) Arrowheads point to pause events in the presence of 400 and 800 nM of SEPT2/6/7. (C) Arrows point to pause events observed during depolymerization of MT plus ends in the presence of 400 and 800 nM of SEPT2/6/7. (D) Images show still frames of MT seeds (green) and dynamic MTs (red) undergoing a pause event during growth in the presence of 800 nM SEPT2/6/7 (D, left) or a pause event during depolymerization in the presence of 400 nM SEPT2/6/7 (D, right). Dashed lines indicate MT length during pause events. Time is in minutes:seconds. (E, F) Bar graphs show the percentage of MTs with continuous growth (no pause, blue) or pause while growing (pause, red) in the presence of 0 nM ($n = 69$), 10 nM ($n = 65$), 100 nM ($n = 51$), 200 nM ($n = 61$), 400 nM ($n = 66$), 800 nM ($n = 50$), 1 μ M ($n = 51$), 2 μ M ($n = 49$), and 4 μ M ($n = 44$) of SEPT2/6/7, or in the presence of 0 nM ($n = 21$), 10 nM ($n = 44$), 100 nM ($n = 31$), 200 nM ($n = 38$), 400 nM ($n = 40$) and 800 nM ($n = 34$) of SEPT9_i1. (G) Bar graph shows the percentage of MTs with continuous depolymerization (no pause, gray) or pause during shrinkage (pause, orange) in the presence of 0 nM ($n = 69$), 10 nM ($n = 65$), 100 nM ($n = 51$), 200 nM ($n = 61$), 400 nM ($n = 66$), 800 nM ($n = 50$), 1 μ M ($n = 51$), 2 μ M ($n = 49$), and 4 μ M ($n = 44$) of SEPT2/6/7.

In the absence of mCherry-SEPT2/6/7, EB1-GFP (75 nM) tracked on growing MT plus ends and was absent from MT tips during catastrophe as previously reported (Bieling *et al.*, 2007, 2008; Dixit *et al.*, 2009). In kymographs, EB1-GFP appeared as diagonal lines, outlining the trajectory of the growing MT plus ends (Figure 4A). Strikingly, the fluorescence intensity of EB1-GFP decreased with increasing concentrations of mCherry-SEPT2/6/7, which was indicative of a reduction in the binding of EB1-GFP to growing MT plus ends (Figure 4A). Quantification of EB1-GFP fluorescence showed a statistically significant reduction from the plus ends of MTs in the

presence of 400–800 nM mCherry-SEPT2/6/7 (Figure 4B). Additionally, enhancement of MT plus-end growth rates by EB1-GFP is reversed in the presence of mCherry-SEPT2/6/7 (Figure 4C); an increase in rates of plus-end growth by EB1 is consistent with previous studies (Vitre *et al.*, 2008; Zanic *et al.*, 2009, 2013). Similar results were obtained with purified nonfluorescent SEPT2/6/7, which inhibited EB1-GFP association with MT plus ends and its effect on MT growth rates (Supplemental Figure S3, A–C).

Inhibition of EB1-GFP binding and tracking by SEPT2/6/7 pointed to a few possibilities. First, EB1-GFP might be sequestered






FIGURE 4: SEPT2/6/7 inhibits EB1 binding to MT plus ends in a concentration-dependent manner. (A) Kymographs show examples of polymerizing MTs (magenta; GMPCPP-stable seeds in bright magenta) with EB1-GFP (75 nM; green) in the absence or presence of increasing concentrations of mCherry-SEPT2/6/7 (red; 100–800 nM). Note that the intensity of EB1-GFP decreases with increasing concentrations of mCherry-SEPT2/6/7. Insets show in higher magnification the binding of mCherry-SEPT2/6/7 to plus-end sites proximal to EB1-GFP and the persistence of mCherry-SEPT2/6/7 to these sites (arrows) while plus ends continue to grow with newly added EB1-GFP. The fluorescence intensity of mCherry-SEPT2/6/7 at 800 nM was adjusted as it is saturated when shown in the same dynamic range and exposure times with concentrations below 800 nM. (B) Mean (\pm SEM) fluorescence intensity of EB1-GFP at MT plus-end tips in the absence ($n = 52$) or presence of 100 nM ($n = 39$), 200 nM ($n = 39$), 400 nM ($n = 33$), and 800 nM ($n = 54$) mCherry-SEPT2/6/7. (C) Graph shows mean (\pm SEM) growth rates of tubulin in the absence of

by soluble SEPT2/6/7 complexes as has been shown before for tau (Ramirez-Rios *et al.*, 2016). Second, SEPT2/6/7 may compete transiently with EB1 for binding to tips of MT plus ends that transition from tubulin-GTP to tubulin-GDP-Pi, which are, respectively, the low- and high-affinity binding sites of EB1 (Zanic *et al.*, 2009; Maurer *et al.*, 2011, 2012). Third, binding of SEPT2/6/7 to the GDP-bound MT lattice may cause long-range conformational changes along MT protofilaments, affecting allosterically the binding of EB1 to plus ends (Zanic *et al.*, 2013; Brouhard and Rice, 2018).

To examine whether EB1-MT binding is inhibited through a direct interaction between SEPT2/6/7 and EB1, we performed protein-binding assays with EB1 and SEPT2/6/7 in buffers of two different ionic conditions including those of the TIRF *in vitro* assay. In neither of these conditions, SEPT2/6/7 interacted with EB1 (Supplemental Figure S4A). In contrast, we were able to detect an association between EB1 and SEPT5 (Supplemental Figure S4, B and C), a septin that possesses the EB1-interaction motif SxIP (Honnappa *et al.*, 2009; Jiang *et al.*, 2012). Hence, SEPT2/6/7 is unlikely to inhibit EB1-MT binding through a direct interaction.

Next, we tested the possibility of a competition between EB1 and SEPT2/6/7 for binding to MTs using GMPCPP- and guanosine 5'-O-[γ -thio]triphosphate (GTP γ S)-stabilized MTs, which, respectively, mimic the GTP-bound (low EB1 affinity) and GDP-Pi-bound (high EB1 affinity) conformations of polymerized tubulin (Zanic *et al.*, 2009; Maurer *et al.*, 2011, 2012; Roth *et al.*, 2018). Unlike EB1, SEPT2/6/7 showed no preference between GMPCPP- and GTP γ S-stabilized MTs (Figure 4D), indicating that SEPT2/6/7 interacts with GTP γ S-bound MT lattices as optimally as observed with the GMPCPP-stabilized MT seeds (Figure 3, B–D). In a competition experiment, EB1-GFP (75 nM) was incubated together with increasing concentrations of mCherry-SEPT2/6/7 in the presence of GTP γ S-stabilized MTs. Decoration of MTs with EB1-GFP was reduced and mCherry-SEPT2/6/7 accumulated to MTs with concomitant displacement of EB1-GFP (Figure 4, E and F); nonfluorescent SEPT2/6/7 had a similar effect on MT-binding of EB1-GFP (Supplemental Figure S3, D and E). These data suggest that association of SEPT2/6/7 with the GDP-Pi lattice of MT plus ends could prevent binding and tracking of EB1.

Despite the competition between EB1 and SEPT2/6/7 for binding to the lattice of GTP γ S-stabilized MTs, which mimic the GDP-Pi segments of MT plus-end tips (high-affinity state of EB1) (Maurer *et al.*, 2011), we were not able to resolve any exchange between mCherry-SEPT2/6/7 and EB1-GFP on the tips of growing plus ends—mCherry-SEPT2/6/7 and EB1-GFP did not alternate tracking on plus-end tips (Figure 4A). Occasionally, mCherry-SEPT2/6/7 localized near the tips of plus ends transiently or remained at the same position while the plus ends continued to grow with EB1-GFP (arrows, insets in Figure 4A). This might be due to the higher affinity

of EB1 for GDP-Pi than SEPT2/6/7; note that full decoration of GTP γ S MTs by EB1 is achieved in much lower concentrations than SEPT2/6/7 (Figure 4E). However, even at high concentrations (800 nM) of SEPT2/6/7, which strongly diminish EB1 binding to plus ends, we did not observe a distinct accumulation of SEPT2/6/7 to plus-end tips—SEPT2/6/7 presence near the plus-end tips was indistinguishable from its localization along the MT lattice (Figure 4A). Thus, while a direct competition at the plus-end tips cannot be ruled out, we posit that MT-bound SEPT2/6/7 may also affect EB1 binding by causing long-range conformational changes on MT protofilaments (Brouhard and Rice, 2018).

End-on collisions of MT plus ends with SEPT2/6/7 filaments results in pausing and EB1 dissociation

Although SEPT2/6/7 does not track with growing MT plus ends in cells or *in vitro*, MT plus ends have been observed to collide end-on with septins (Bowen *et al.*, 2011; Nolke *et al.*, 2016). We, therefore, sought to determine whether septin filaments can directly impact MT plus-end dynamics during end-on encounters, which do not involve association of septin complexes with the MT lattice.

We reconstituted MT dynamics in the presence of septin filaments by immobilizing mCherry-SEPT2/6/7 and MT seeds on glass substrata. On introduction of soluble tubulin, we were able to visualize the polymerizing plus ends of MTs intersecting with filamentous septins. During these encounters, MT plus ends paused, continued to grow, or depolymerized (Figure 5A). Quantification showed that ~46% of encounters with immobilized mCherry-SEPT2/6/7 filaments resulted in MT plus pausing (Supplemental Movie S5), while continuation of growth and depolymerization were observed at ~43% and ~9% of intersections, respectively. Notably, MT pausing events were significantly higher (~2-fold) compared to plus-end collisions with immobilized actin filaments (Figure 5B). On encountering F-actin, continuation of MT plus-end growth was much more frequent than pausing—continued growth (73%) was threefold more frequent than pausing (~23%) (Figure 5B; Supplemental Movie S6).

Next, we asked whether MT plus ends exhibited similar behavior on encountering septin and actin filaments in the presence of EB1-GFP. Interestingly, EB1-bound MT plus ends paused less than free MT plus ends at both septin and F-actin intersections (Figure 5, C and E). However, EB1-bound plus ends still paused more frequently at intersections with septins than actin filaments (26% vs. 16%) and dissociation of EB1 from plus ends was more frequent at collisions with SEPT2/6/7 than actin filaments (Figure 5, D and E). These data show that pausing of MT plus ends and EB1 dissociation is more likely to occur at intersections with septin than actin filaments.

To test whether septin filaments are a more potent barrier to MT plus-end growth than actin filaments in living cells, we imaged and quantified the dynamics of end-binding protein 3 (EB3)-tdTomato at

mCherry-SEPT2/6/7 or EB1-GFP (green; $n = 75$), and in the presence of 75 nM EB1-GFP (magenta; $n = 80$) without or with 800 nM mCherry-SEPT2/6/7 (red; $n = 52$). Note that the enhancement of MT growth rate by EB1-GFP is inhibited by mCherry-SEPT2/6/7. (D) Dot plots show the mean (\pm SEM) fluorescent intensity of mCherry-SEPT2/6/7 on the lattice of GMPCPP- and GTP γ S-stabilized MTs. Quantification was performed from static images of MTs after 15 min of incubation in the presence of 10 nM (GMPCPP, $n = 587$; GTP γ S, $n = 482$), 100 nM (GMPCPP, $n = 597$; GTP γ S, $n = 511$), 200 nM (GMPCPP, $n = 730$; GTP γ S, $n = 537$), 400 nM (GMPCPP, $n = 614$; GTP γ S, $n = 457$), and 800 nM (GMPCPP, $n = 454$; GTP γ S, $n = 322$) of mCherry-SEPT2/6/7. (E) Images show examples of GTP γ S seeds (magenta) incubated with 75 nM EB1-GFP (green) and increasing concentrations of mCherry-SEPT2/6/7 (red; 100–800 nM). Note that EB1-GFP binding decreases with increasing concentrations of mCherry-SEPT2/6/7. (F) Mean (\pm SEM) fluorescent intensity of EB1-GFP on GTP γ S-stabilized MTs in the absence or presence of mCherry-SEPT2/6/7. Quantification was performed from static images of MTs after 15 min of incubation without ($n = 528$) or with mCherry-SEPT2/6/7 at concentrations of 100 nM ($n = 626$), 200 nM ($n = 559$), 400 nM ($n = 581$), and 800 nM ($n = 626$); ns, nonsignificant ($p > 0.05$). * $p < 0.05$, *** $p < 0.001$, **** $p < 0.0001$.




FIGURE 5: MT plus-end collisions with SEPT2/6/7 filaments enhance pausing and EB1 dissociation. (A) Kymographs show examples of end-on collisions of growing MT plus ends (green; GMPCPP-stable MT seeds in magenta) with mCherry-SEPT2/6/7 or Rhodamine-actin filaments (red), which were immobilized on the glass matrix. Arrowheads point to pauses of MT plus ends at intersections with mCherry-SEPT2/6/7 or Rhodamine-labeled actin filaments. (B) Bar graph shows the percentage of intersection events in which MT plus ends paused (red) on encountering SEPT2/6/7 ($n = 207$ events) or actin filaments ($n = 178$ events), continued to grow (blue) or depolymerized (green). Note that the percentage of pause events increased by twofold on encountering SEPT2/6/7 compared with actin. (C) Kymographs show examples of dynamic MTs (magenta) assembled with 75 nM EB1-GFP (green) in the

sites of intersection with GFP-SEPT7 or GFP-F-tractin in primary rat hippocampal neurons (Figure 5, F–H). In agreement with the *in vitro* data, EB3 comets paused more frequently on intersecting with GFP-SEPT7 than GFP-F-tractin (~46% vs. ~33%; Figure 5I). Kymographs showed that EB3 comets became dimmer or disappeared on contact with SEPT7 filaments (Figure 5, G and H). EB3 comets were also found to resume growth with reduced (deceleration) or enhanced (acceleration) rates compared with their velocity prior to pausing at SEPT7 or F-tractin. Notably, deceleration of MT plus-end growth was more frequent after collision with SEPT7 than F-tractin (Figure 5J) and the reduction in velocity was twofold higher on encountering septin ($\Delta V = 0.14 \pm 0.02 \mu\text{m/s}$) than actin ($\Delta V = 0.07 \pm 0.006 \mu\text{m/s}$) (Figure 5K); frequency and rates of accelerated EB3 movement were not different between septin and actin. Thus, septin filaments not only are more potent barriers than actin filaments in pausing MT plus ends but they also slow down MT regrowth after pausing.

DISCUSSION

A growing number of studies implicate septins in MT organization and dynamics as well as MT-dependent functions, but how septins impact the dynamic instability of MTs is not well understood. Results from cell biological experiments have been difficult to interpret as septins are multifunctional proteins with various binding partners, several of which associate with the cytoskeleton. Hence, *in vitro* reconstitution experiments can provide valuable insights by elucidating the direct effects of septins on MT dynamics.

In this study, MT dynamics were reconstituted with SEPT2/6/7 complexes and filaments. Our results show that the effects of SEPT2/6/7 on MT dynamics are dependent on their concentration and state of assembly, which may account for the disparity of previous findings obtained from septin knockdown approaches. We found a biphasic graded effect on MT growth that correlated with a concentration-dependent assembly of SEPT2/6/7 into higher-order filaments (Figure 6A). At lower concentrations (10–100 nM), where assembly of SEPT2/6/7 complexes into higher-order filaments is limited, MT growth and elongation were enhanced. However, at higher concentrations (low μM), which enhance the filamentous state of SEPT2/6/7, MT growth was stunted. Interestingly, in the intermediate range of concentrations, SEPT2/6/7 functions as a pausing factor, causing transient arrests of MT plus-end growth.

The effects of SEPT2/6/7 on MT dynamics have striking similarities and differences with SEPT9. While both SEPT2/6/7 heteromers and SEPT9 homomers promote the growing phase of MTs, SEPT2/6/7 achieves this effect at lower concentrations than SEPT9 (Nakos *et al.*, 2019). Moreover, micromolar concentrations of SEPT9 stabilize elongated MTs, which seize to grow and shrink, while micromolar SEPT2/6/7 dampens MT dynamics without stabilizing MTs, which undergo more catastrophe. Additionally, SEPT9 binds and recruits unpolymerized tubulin to the lattice of MTs (Nakos *et al.*, 2019), while SEPT2/6/7 does not bind unpolymerized tubulin. Conversely, submicromolar concentrations of SEPT2/6/7 can pause MT plus-end growth, which was not observed with SEPT9. Taken together, these data indicate that there are septin subunit- and paralogue-specific differences in the modulation of MT dynamics, which might be physiologically relevant for the modulation of MT dynamics in cell types and pathological conditions with distinct patterns of septin expression.

Our results show that SEPT2/6/7 can trigger dissociation of EB1 from MT plus ends *in cis* by binding to the lattice of plus ends or *in trans* by coming in contact as a filamentous structure with on-coming MT plus ends (Figure 6, B and C). A similar effect has been shown for tau, which antagonizes EB1 binding to plus ends (Ramirez-Rios *et al.*, 2016). However, unlike tau, which inhibits EB1-MT binding by forming a soluble complex with EB1 off MTs, SEPT2/6/7 does not appear to sequester EB1 off MTs. Interestingly, SEPT2/6/7 competes with EB1 for binding to GTP γ S-stabilized MTs, which resemble the EB1-preferred tubulin-GDP-Pi conformation of plus-end tips. In addition, SEPT2/6/7 is able to reduce the enhanced MT growth rates that we observed in the presence of EB1. Although EB1 dissociation on MT plus-end contact with SEPT2/6/7 filaments indicates a direct competition between EB1 and septins at the tips of MT plus ends, SEPT2/6/7 tracking on MT plus ends is rare. Moreover, we have not been able to observe an exchange of EB1 and SEPT2/6/7 on MT plus ends. If SEPT2/6/7 does not affect EB1 binding at the tips of plus ends, binding of SEPT2/6/7 to the MT-GDP lattice may result in conformational changes, which are propagated to the tips of plus ends allosterically along the protofilaments of MTs (Brouhard and Rice, 2018). Evidence for this mechanism comes from studies of the potent MT polymerase XMAP215 and EB1 (Zanic *et al.*, 2013). Even though XMAP215 and EB1 do not interact, a synergistic effect is posited to occur through conformational

presence of immobilized mCherry-SEPT2/6/7 or Rhodamine-labeled actin filaments (red). Arrowheads point to loss or diminution of EB1-GFP from plus-end tips at intersections with septin and actin filaments. Note that EB1-GFP binding at the MT plus-end tips is reduced or lost at points of intersection with mCherry-SEPT2/6/7 filaments. (D) Percentage of collision events during which EB1-GFP was reduced or lost from MT plus ends (red) or remained unaffected (yellow) on encountering mCherry-SEPT2/6/7 ($n = 132$ events) or actin filaments ($n = 213$ events). (E) Bar graph shows the percentage of events of MT plus ends with EB1-GFP (75 nM), which paused (red) on encountering mCherry-SEPT2/6/7 ($n = 132$ events) or actin filaments ($n = 213$ events), continued to grow (blue) or depolymerized (green). Note that the percentage of pause events is higher on encountering SEPT2/6/7 (26%) than actin filaments (16%). (F–H) Still images (F) show a rat hippocampal neuron (DIV3) cotransfected with plasmids expressing GFP-SEPT7 and EB3-tdTomato. An outlined neurite region (dashed rectangle) in which EB3-tdTomato movement was analyzed with respect to GFP-SEPT7 is shown in higher magnification and corresponds to the depicted kymographs (G, H). Kymographs show the trajectories of EB3-tdTomato comets (red) that intersect with GFP-SEPT7 (green) (G, H). Outlined regions with diminution of EB3-tdTomato intensity (dashed rectangles 1 and 2) during overlap with GFP-SEPT7 and thereafter resumed movement (dashed rectangle 2) are shown in higher magnification (H). (I) Bar graph shows the percentage of EB3 comets that paused (red) or continued to move (blue) on encountering SEPT7 ($n = 342$ comets; 27 neurites) or actin filaments ($n = 399$ comets; 22 neurites). Note that the percentage of pause events is higher on encountering SEPT7 (46%) than actin filaments (33%). (J) Bar graph shows the percentage of EB3 comets that resumed movement with reduced (deceleration; gray) or enhanced velocity (acceleration; orange) compared with their velocity prior to pausing at SEPT7 ($n = 124$) or actin filaments ($n = 168$). Deceleration was more frequent after pausing at SEPT7 (61%) than actin filaments (46%). (K) Mean (\pm SEM) change of velocity in the events of deceleration of EB3 comets after pausing at SEPT7 ($n = 76$) or actin filaments ($n = 77$); ns, nonsignificant ($p > 0.05$). * $p < 0.05$.




FIGURE 6: SEPT2/6/7 effects on MT plus-end dynamics and EB1 plus-end tracking. (A) SEPT2/6/7 associates with GMPCPP-bound seeds with higher affinity than the GDP-bound lattice, and affects plus-end dynamics in a manner that depends on septin concentration and state of assembly (oligomers, polymers). At low concentrations (10–100s nanomolar), oligomeric SEPT2/6/7 induces persistent growth and elongation of MT plus ends. At higher concentrations (micromolar), however, higher-order filamentous SEPT2/6/7 results in stunted MT growth and enhanced catastrophe. At the intermediate nanomolar-to-micromolar concentration range, SEPT2/6/7 pause MT plus ends in their growth phase. (B) SEPT2/6/7 inhibits EB1 association and tracking on MT plus ends. This effect could be due to either transient competition for binding on MT plus ends or through long-range conformational effects, which are propagated along the MT lattice to the tips of plus ends. (C) MT plus ends pause and EB1 dissociates during MT end-on collisions with SEPT2/6/7 filaments.

changes (Zanic *et al.*, 2013; Brouhard and Rice, 2018). Functionally, a septin-mediated dissociation of EB1 from MT plus ends could provide a mechanism for pausing and stable capture of MT plus ends. Given that EB1 appears to protect MT plus ends from pausing during end-on collisions with septin and actin filaments (Figure 5), dissociation of EB1 would further enhance pausing of MT plus ends on contact with septin filaments.

How are the concentration-dependent effects that septins exert on MT plus ends *in vitro* relevant to the concentration, organization, and localization of septins in living cells? MT growth and elongation are enhanced at septin concentrations of hundreds of nanomolar, which reportedly are the estimated concentrations of septins such as SEPT2 (~800 nM), SEPT6 (~200 nM), SEPT7 (~500 nM), and SEPT9 (~400 nM) in HeLa cells (Hein *et al.*, 2015). By contrast, micromolar septin concentrations dampen MT dynamics, resulting in stunted growth or complete MT stabilization. Given that septins are not uniformly distributed in cells but rather enriched in specific subcellular regions (e.g., membrane domains of micron-scale curvature), it is plausible that septins tune MT dynamics in a region-specific manner according to their local concentration. Therefore, persistent MT growth could occur in cytoplasmic areas of moderate septin concentration, while MT pausing and stabilization could occur at cytoplasmic and membrane regions of septin enrichment. The latter could be key for the capture of MT plus ends at the base of membrane protrusions, where septins accumulate at domains of micron-

scale curvature (Zhang *et al.*, 1999; Steels *et al.*, 2007; Akil *et al.*, 2016; Bridges *et al.*, 2016; Dolat and Spiliotis, 2016; Cannon *et al.*, 2019), or the stabilization and dampening of MT plus-end dynamics at the tips of primary cilia, in which septins are also highly enriched (Hu *et al.*, 2010; Ghossoub *et al.*, 2013; Fliegauf *et al.*, 2014).

The concentration- and assembly-dependent effects of septins on MT dynamics suggest that previous observations from septin knockdown approaches were sensitive to the expression levels and organization of the targeted septins and their complexes. Of note, septin turnover is slow and knockdown levels can alter residual septin assembly and localization. Therefore, interpretations of previous phenotypes are prone to confounding variables and may not be directly comparable to results from *in vitro* approaches, which conversely lack the protein complexity of plus-end tips and MT surfaces that include MAPs and motors. In epithelial cells, previous work indicated that MT-associated septins guide plus-end growth during MT-MT collision (Bowen *et al.*, 2011). While the effects of SEPT2 knockdown on MT plus-end dynamics are consistent with enhanced MT growth by nanomolar concentrations of SEPT2/6/7 *in vitro*, SEPT2 depletion increases plus-end pausing at MT-MT intersections rather than reducing it as the *in vitro* data would predict (Bowen *et al.*, 2011). Note, however, that this study examined MT plus-end collisions with septin and actin filaments and not septin-coated MTs. Nevertheless, we postulate that in cells, in which plus ends and MT lattices associate with MT motors, pausing of MT plus ends by

SEPT2/6/7 might be an intermediate step in allowing motors to engage with opposing MTs and thereby direct their growth. Interestingly, we recently discovered that MT-associated septins modulate kinesin attachment to MTs and their motility (Karasmans et al., 2018). Thus, it is plausible that motor-bound plus ends behave differently when they encounter septin-coated MTs and in the absence of septins, MT plus-end pausing might be mediated by MAPs. Overall, our findings are in general agreement with septin roles in the intracellular capture and guidance of MTs (Bowen et al., 2011; Hu et al., 2012; Nolke et al., 2016), but may not fully recapitulate in vivo phenomena due to the minimal components and reductionist nature of the in vitro assays.

Despite these caveats, modulation of MT plus-end growth at sites of contact with septin filaments suggests that septins may function as cytoplasmic barriers, which gate spatially and temporally MT growth and invasion. Recent studies have shown that actin filaments act as physical barriers that regulate MT growth in vitro and in cultured cells (Colin et al., 2018; Inoue et al., 2019). Our results show that septins are more potent barriers in pausing and dampening MT regrowth than actin filaments. As septins associate preferentially with transverse arcs of migrating cells (Dolat et al., 2014), they may be critical for the stable capture and regrowth of MT plus ends against the retrograde flow of lamellar actin that sweeps MTs centripetally. Hence, septins might be part of the mechanisms that enable MTs to reach the front of migrating cells.

In summary, our findings demonstrate that SEPT2/6/7 can directly impact MT dynamics in a manner that depends on their concentration and state of assembly into higher-order filamentous structures. Thus, septins could play important roles in the modulation of MT dynamics in health and disease. As septins have been identified in the neurofibrillary tangles and Lewy bodies of Alzheimer's and Parkinson's patient brains (Kinoshita et al., 1998; Ihara et al., 2003, 2007; Shehadeh et al., 2009), filamentous aggregates of septins could have adverse effects on MT dynamics exacerbating the cytotoxicity and pathological progression of neurodegenerative disorders.

MATERIALS AND METHODS

Plasmids

The pET-15b plasmid encoding for His-SEPT2-mCherry was constructed and gifted by Shae Padrick (College of Medicine, Drexel University), and the pNEA-vH and pNCS vectors encoding respectively for His-SEPT2 and SEPT6/7-strep (Mavrakakis et al., 2014) have been previously described and provided by Amy Gladfelter (Department of Biology, University of North Carolina). In the pNCS vector, the N-terminal 19 amino acids (1–57 nucleotides) of human SEPT7 were added using a four-step Kapa Biosystems Site-directed mutagenesis protocol. Briefly, SEPT6/7-strep tagged was amplified with KAPA HiFi HotStart DNA polymerase (KK2502; Kapa Biosystems) using the primers 5'-GTAATAATTTTGTAACTTTAAGAAGGAGATATACATATGTCGGTCAGTATGG TAGCTCAACAGAAGAA-3' and 5'-TTCTTCTGTTGAGCTACCATACTGACCGACATATGTA TATCTCTTCTAAAGTTAAACAAAATTATTAC-3' to insert the first 12 base pairs (1–12 base pairs) at the N-terminus of SEPT7. The amplified PCR product was treated with *DpnI* (R0176L; New England Biolabs) for 1 h at 37°C. *DpnI*-treated product was heat inactivated at 80°C for 20 min and subsequently was transformed in *Escherichia coli* DH5a cells. The new construct was used as a template and amplified using the primers 5'-GATATACATATGTCGGTCAGTGCAGATCCG CTGCTATGGTAGCTCAACAGAAGAAC-3' and 5'-GTTCTTCTGTTGAGCTACCATAGC AGCGGATCTCGCACTGACCGACATATGTATATC-3' in order to insert the next 15 base

pairs (13–27 base pairs) of SEPT7. Subsequently, the PCR product was treated as above and used as a template to introduce the next 18 base pairs (28–45 base pairs) of SEPT7 using the primers 5'-TGC-GAGATCCGCTGCTGCTGAGGAGAGGAGCGTCATGGTAGCTCAACAGA-3' and 5'-TCTGTTGAGCTACCATGACGCTCCTCTCTCAGCAGCAGCGGATCTCGCA-3'. Finally, the resulting construct was used as a template and amplified using the primers 5'-TGAGGAGAGGAGCGTCAACAGCAGCACCATGGTAGCTCAACAGA-3' and 5'-TCTGTTGAGCTACCATGGTGTGCTGTTGAGGCTCCTCTCCTCA-3' to insert the last 12 base pairs (46–57 base pairs) of SEPT7. The pGEX-3x plasmid encoding for GST-tagged EB1 was a gift from Anna Akhmanova (Department of Biology, Utrecht University). The pET-21a plasmid encoding for recombinant His-tagged EB1-GFP was a gift from Antonina Roll-Mecak (Cell Biology and Biophysics, National Institutes of Health [NIH]) (Vemu et al., 2018). EB3-tdTomato was purchased from Addgene (tdTomato-N1-EB3; Addgene plasmid # 50708; <http://n2t.net/addgene:50708>; RRID: Addgene_50708) (Merriam et al., 2013). Plasmid encoding rat EGFP-SEPT7 was a kind gift from Smita Yadav (University of Washington) (Yadav et al., 2017). Plasmid encoding GFP-F-tractin was a kind gift from Tatyana Svitkina (Department of Biology, University of Pennsylvania) and was cloned into the pEGFP-C1 vector (Johnson and Schell, 2009; Yang and Svitkina, 2019). His-tagged SEPT5 plasmid was constructed by PCR amplifying SEPT5 from pCMV-Myc-tagged SEPT5 (Addgene plasmid #27272; <http://n2t.net/addgene:27272>; RRID: Addgene_27272) (Amin et al., 2008) using the primers 5'-ATG GGT CGC GGA TCC GAA ATG AGC ACA GGC CTG-3' and 5'-CTC GAG TGC GGC CGC ATC ACT GGT CCT GCA TC-3' and inserting the fragment into the pET28a(+) vector.

Protein expression and purification

Recombinant SEPT2/6/7 complex was prepared as described before (Mavrakakis et al., 2014). The plasmids encoding for His-SEPT2 or His-SEPT2-mCherry and SEPT6/7-strep were cotransformed into *E. coli* BL21 (DE3) (Invitrogen). Bacterial cultures were grown to OD₆₀₀ of 2–3 and induced with 1 mM IPTG (isopropyl β-D-1-thiogalactopyranoside) for 1 h at 37°C (His-SEPT2, SEPT6/7-strep) or to OD₆₀₀ of 0.5 and induced with 0.2 mM IPTG for 16 h at 18°C (His-SEPT2-mCherry, SEPT6/7-strep). Cultures were centrifuged at 4000 rpm for 20 min at 4°C. Pellets were resuspended in lysis buffer containing 50 mM Tris-HCl, pH 8.0, 500 mM KCl, 10 mM imidazole, 5 mM MgCl₂, 1 mg/ml lysozyme, 1 mM PMSF, and 1× Bacterial Protease Arrest cocktail (G-Biosciences; 786–330) and lysed by sonication (10 sets of 15 pulses on ice with a 30-s interval between each set). Cell lysates were centrifuged at 13,000 rpm for 30 min at 4°C and passed through a 0.45-μm-pore filter. Supernatants were loaded onto gravity flow columns with Ni-NTA agarose beads (745400.25; Macherey-Nagel), which were equilibrated with 10 ml of buffer containing 50 mM Tris-HCl, pH 8.0, 500 mM KCl, 10 mM imidazole, 5 mM MgCl₂, 1 mM PMSF, and 1× of the Bacterial Protease Arrest cocktail. Protein complexes were eluted in elution buffer containing 50 mM Tris-HCl, pH 8.0, 500 mM KCl, 250 mM imidazole, and 5 mM MgCl₂ and loaded to a StrepTrap HP column (GE Healthcare) equilibrated with buffer containing 50 mM Tris-HCl, pH 8, 300 mM KCl, and 5 mM MgCl₂. Protein complexes were eluted in elution buffer containing 50 mM Tris-HCl, pH 8.0, 300 mM KCl, 5 mM MgCl₂, and 2.5 mM D-Desthiobiotin (Sigma; D1411) and dialyzed overnight at 4°C in BRB80 (80 mM PIPES, pH 6.9, 2 mM MgCl₂, 1 mM EGTA [ethylene glycol-bis(2-aminoethyl ether)-N,N,N',N'-tetraacetic acid]). His-SEPT2-mCherry-SEPT6/7-strep was further purified using an AKTA FPLC system (GE Healthcare) with a Superdex 200 10/300 GL (Amersham Biosciences) gel filtration column.

Recombinant His-tagged EB1-GFP and His-tagged SEPT5 were transformed into *E. coli* BL21 (DE3) (Invitrogen). Bacterial cultures were grown to OD₆₀₀ of 0.6–0.8 and induced with 0.2 mM IPTG for 16 h at 18°C for His-tagged EB1-GFP or cultures were grown to OD₆₀₀ of 0.4–0.6 and induced with 1 mM IPTG for 5 h at 23°C for His-tagged SEPT5. Cultures were centrifuged at 4000 rpm for 20 min at 4°C. Pellets were resuspended in lysis buffer containing 50 mM Tris, pH 8.0, 150 mM NaCl, 10% glycerol, 1 mM PMSF, 1 mg/ml lysozyme, 10 mM imidazole, and 1× Bacterial Protease Arrest protease inhibitor cocktail (G-Biosciences; 786-330) and lysed by sonication (10 sets of 15 pulses on ice with a 30-s interval between each set). Cell lysates were centrifuged at 13,000 rpm for 30 min at 4°C and passed through a 0.45- μ m-pore filter. Supernatants were loaded onto a gravity flow column with Ni-NTA agarose beads (745400.25; Macherey-Nagel), which were equilibrated with 10 ml of buffer containing 50 mM Tris, pH 8.0, 150 mM NaCl, 10% glycerol, 1 mM PMSF, 10 mM imidazole, and 1× Bacterial Protease Arrest protease inhibitor cocktail. Columns were washed with 30 ml washing buffer (50 mM Tris, pH 8.0, 300 mM NaCl, 10% glycerol, 10 mM imidazole). Proteins were eluted in elution buffer containing 50 mM Tris, pH 8.0, 150 mM NaCl, 10% glycerol, and 250 mM imidazole. His-tagged EB1-GFP was dialyzed overnight at 4°C in BRB80 (80 mM PIPES, pH 6.9, 2 mM MgCl₂, 1 mM EGTA) supplemented with 50 mM KCl and was further purified using an AKTA FPLC system (GE Healthcare) with a Superdex 200 10/300 GL (Amersham Biosciences) gel filtration column. His-tagged SEPT5 was dialyzed overnight at 4°C in buffer containing 50 mM Tris, pH 8.0, 150 mM NaCl, and 10% glycerol.

Recombinant GST and GST-tagged EB1 were transformed into *E. coli* BL21 (DE3) (Invitrogen). Bacterial cultures were grown to OD₆₀₀ of 0.7 and induced with 1 mM IPTG for 5.5 h at 25°C. Cultures were centrifuged at 4000 rpm for 20 min at 4°C. Cell pellets were resuspended in lysis buffer containing 50 mM Tris, pH 8.0, 150 mM NaCl, 2 mM MgCl₂, 5 mM DTT (dithiothreitol), 10% glycerol, 0.1% Triton X-100, 1 mM PMSF, 1 mg/ml lysozyme, and 1× Bacterial Protease Arrest cocktail (G-Biosciences; 786-330) and lysed by sonication (10 sets of 15 pulses on ice with a 30-s interval between each set). Cell lysates were centrifuged at 13,000 rpm for 30 min at 4°C and passed through a 0.45- μ m-pore filter. Supernatants were loaded onto gravity flow columns with glutathione agarose beads (16100; Thermo Scientific) equilibrated with 10 ml of buffer containing 50 mM Tris, pH 8.0, 150 mM NaCl, 2 mM MgCl₂, 5 mM DTT, 10% glycerol, 0.1% Triton X-100, 1 mM PMSF, and 1× Bacterial Protease Arrest protease inhibitor cocktail. Subsequently, columns were washed with 30 ml washing buffer (50 mM Tris, pH 8.0, 150 mM NaCl, 2 mM MgCl₂, 5 mM DTT, 10% glycerol, and 0.1% Triton X-100) and proteins were eluted in elution buffer containing 50 mM Tris, pH 8.0, 150 mM NaCl, 2 mM MgCl₂, 5 mM DTT, 10% glycerol, and 10 mM glutathione. All proteins were dialyzed overnight at 4°C in buffer containing 50 mM Tris, pH 8.0, 150 mM NaCl and 10% glycerol. Recombinant His-tagged SEPT9_{i1} was expressed and purified as described before (Nakos et al., 2019).

GST-pull-down assays

Protein-protein interaction assays were performed between GST or GST-tagged EB1 and recombinant His-tagged SEPT2/6/7 or SEPT5 proteins. GST or GST-EB1 (5 μ g) were incubated with 20 μ l glutathione agarose beads (16100; Thermo Scientific) for 2 h at 4°C. Subsequently, beads were washed twice with pull-down buffer (50 mM HEPES, pH 7.4, 150 mM NaCl, 0.1% Triton X-100, 1 mM PMSF, 5 mM DTT, 2 mM EGTA, and 10% glycerol) or BRB80 supplemented with 0.1% Triton X-100, 1 mM PMSF, and 50 mM KCl. Beads were incu-

bated with His-tagged SEPT2/6/7 or SEPT5 (5 μ g) for 2 h at 4°C. Beads were washed five times with pull-down buffer or BRB80 (with 0.1% Triton X-100, 1 mM PMSF, and 50 mM KCl) before they were resuspended with loading buffer and boiled. Eluted complexes loaded onto 10% SDS-PAGE gels and transferred to a nitrocellulose membrane. Membranes were blocked with 5% nonfat dry milk and 1% bovine serum albumin (BSA) for 1 h at room temperature. Membranes were washed with PBS-T (phosphate-buffered saline [PBS] 1×/0.1% Tween 20) and incubated with mouse antibody against 6xHis-tag (1:2000; Qiagen), rabbit antibody against GST-tag (1:10,000; Santa Cruz), or rabbit antibody against SEPT7 (1:10,000; IBL) diluted in PBS-T/2% BSA over night at 4°C. Subsequently, membranes were washed with PBS-T and incubated with anti-mouse or anti-rabbit secondary antibodies (LiCor) for 1 h at room temperature before scanning with an imaging system (Odyssey; LICOR).

In vitro polymerization of recombinant SEPT2/6/7 complexes

To test the ability of SEPT2/6/7 complexes to form filaments in vitro (Supplemental Figure S1A), purified nonfluorescent and fluorescent mCherry-SEPT2/6/7 complexes were incubated overnight (~23 h) at 4°C in BRB80 (80 mM PIPES, pH 6.9, 2 mM MgCl₂, 1 mM EGTA) in the presence of 500 mM KCl. 100 nM of SEPT2/6/7 proteins were diluted in BRB80 supplemented with 50 mM KCl, 1 mg/ml BSA and 1 mM GTP. Proteins were incubated at room temperature for 4 h. Subsequently, SEPT2/6/7 (fluorescent and nonfluorescent) complexes were placed on top of poly-L-lysine coated coverslips. mCherry-SEPT2/6/7 complexes were mounted on a slide, sealed with nail polish and imaged with TIRF. Nonfluorescent SEPT2/6/7 complexes were fixed with 3% PFA (paraformaldehyde; EM Sciences) for 10 min, quenched twice with 75 mM NH₄Cl (A9434; Sigma) and blocked with 2% BSA. SEPT2/6/7 complexes were incubated with a mouse antibody against 6xHis-tag (1:100; Qiagen) for 30 min. Subsequently, SEPT2/6/7 complexes were washed with BRB80 and incubated with a secondary donkey anti-mouse antibody conjugated with Alexa 594 (1:5000; Jackson ImmunoResearch Laboratories) for 30 min. SEPT2/6/7 complexes were washed with BRB80 and mounted on a slide, sealed with nail polish, and imaged with TIRF.

Pelleting assay of recombinant SEPT2/6/7

In pelleting assays of recombinant SEPT2/6/7 complexes, increasing concentrations of SEPT2/6/7 (10 nM–4 μ M) were incubated in BRB80 (80 mM PIPES, pH 6.9, 2 mM MgCl₂, 1 mM EGTA) supplemented with 1 mM GTP for 15 min at room temperature. Reactions were placed on top of a cushion buffer and centrifuged at 15,600 \times g (Optima TL100; Beckman Coulter) for 10 min at 25°C. Equal volumes of supernatant and pellet fractions were loaded onto 8% SDS-PAGE gels, which were stained with Coomassie Brilliant Blue. SDS-PAGE gels were scanned and protein bands were quantified with the Odyssey scanning system (LICOR).

TIRF assays of dynamic MTs

Imaging chambers were prepared and constructed as before (Tanenbaum et al., 2013; Reid et al., 2016; Nakos et al., 2019). TIRF imaging was performed as described before (Nakos et al., 2019). Imaging chambers were prepared by sequential treatment with the following buffers: 1% Pluronic F-127 for 5 min, 5 mg/ml Biotin-BSA (A8549; Sigma-Aldrich) for 5 min, and 0.5 mg/ml Neutravidin (A2666; Invitrogen) for 5 min. GMPCPP-stabilized MT seeds (~60–80 nM) were floated into the chamber for 15 min and then washed with blocking buffer (BRB80, 1% Pluronic F-127, 1 mg/ml BSA) for

5 min. GMPCPP MT seeds were prepared by incubating 77% unlabeled tubulin (T240; Cytoskeleton) with 11.5% biotin-tubulin (T333P; Cytoskeleton) and 11.5% HiLyte-488-tubulin (TL488M; Cytoskeleton) or 11.5% HiLyte-647-tubulin (TL670M; Cytoskeleton). The tubulin mix (20 μ M) with 1 mM GMPCPP (NU-405L; Jena Bioscience) in BRB80 (80 mM PIPES, pH 6.9, 2 mM MgCl₂, 1 mM EGTA) was incubated at 37°C for 30 min. After incubation, seeds were diluted in BRB80 and span for 15 min at 100,000 \times g (Optima TL100; Beckman Coulter). Sedimented MT seeds were resuspended in 60 μ l of BRB80. In chambers, MT polymerization was induced by floating a mix of 10 μ M tubulin, which contained 6.6% Rhodamine-tubulin (TL590M; Cytoskeleton) or HiLyte-488-tubulin (TL488M; Cytoskeleton) in BRB80 with 1 mM GTP, 0.1 mg/ml BSA, 0.1% Pluronic F-127, 0.1% κ -casein, 0.2% methyl cellulose, oxygen scavenging system (0.5 mg/ml glucose oxidase, 0.1 mg/ml catalase, 4.5 mg/ml D-glucose, 70 mM β -mercaptoethanol), and recombinant SEPT2/6/7 or mCherry-SEPT2/6/7.

For TIRF assays with recombinant EB1-GFP and mCherry-SEPT2/6/7 or nonfluorescent SEPT2/6/7, GMPCPP MT bright seeds were prepared as described above by incubating 66.7% unlabeled tubulin (T240; Cytoskeleton) with 11.5% biotin-tubulin (T333P; Cytoskeleton) and 21.8% HiLyte-647-tubulin (TL670M; Cytoskeleton). MT polymerization was induced by introducing a mix of 10 μ M tubulin, which contained 4.67% HiLyte-647-tubulin (TL670M; Cytoskeleton) in BRB80 supplemented with 1 mM GTP, 0.1 mg/ml BSA, 0.1% Pluronic F-127, 0.1% κ -casein, 0.2% methyl cellulose, oxygen scavenging system (0.5 mg/ml glucose oxidase, 0.1 mg/ml catalase, 4.5 mg/ml D-glucose, 70 mM β -mercaptoethanol), 50 mM KCl and recombinant mCherry-SEPT2/6/7 or nonfluorescent SEPT2/6/7 and EB1-GFP proteins. Imaging chambers were sealed with vacuum grease prior to imaging at 37°C. Imaging was performed every 2 s for 15 min.

Visual septin-tubulin binding assays

Imaging of binding of soluble unpolymerized tubulin to septins by TIRF (Figure S2) was done by adapting a previous MT nucleation assay (Lazarus *et al.*, 2013). Imaging chambers were constructed using acid washed slides and silanized glass cover slips. Acid washed glass coverslips were incubated for 10 min with 20% 3-Aminopropyltriethoxysilane (430941000; ACROS Organics) diluted in acetone and subsequently washed with ddH₂O for 15 min. Imaging chambers were first incubated with 1% Pluronic F-127 for 5 min. Chambers were washed with BRB80 and then were coated with mouse antibody against 6xHis-tag (1:50; Qiagen) for 10 min. Imaging chambers were washed with BRB80 and blocked with 5% Pluronic F-127 for 5 min. Flow chambers were washed with BRB80-1 mg/ml BSA and subsequently incubated for 15 min with increasing concentrations of recombinant His-tagged SEPT9 or SEPT2/6/7. Unbound proteins were removed from chambers with multiple washes of BRB80-1 mg/ml BSA. Subsequently, we floated into the chambers a tubulin mix (3.5 μ M), which contained 8.6% Rhodamine-tubulin (TL590M; Cytoskeleton) in BRB80 supplemented with 2.5 mM GTP, 1 mg/ml BSA, 0.25% Pluronic F-127, 0.1% κ -casein, and oxygen scavenging system (0.5 mg/ml glucose oxidase, 0.1 mg/ml catalase, 4.5 mg/ml D-glucose, 70 mM β -mercaptoethanol). Chambers were sealed with vacuum grease and imaging was performed every 2 s for 15 min at 37°C.

TIRF assays of dynamic MTs with SEPT2/6/7 or actin filaments

GMPCPP-stabilized HiLyte-647-labeled MT seeds were made as described above. Imaging chambers were incubated first with 1%

Pluronic F-127 for 5 min. Chambers were washed with BRB80 and then were coated with rabbit antibody against SEPT7 (1:10; IBL) for 10 min. Imaging chambers were washed with BRB80 and blocked with 5% Pluronic F-127 for 5 min. Flow chambers were washed with BRB80-1 mg/ml BSA and subsequently incubated for 15 min with 800 nM of recombinant mCherry-SEPT2/6/7. Unbound proteins were removed from chambers with multiple washes of BRB80-1 mg/ml BSA. Imaging chambers were then incubated with 5 mg/ml Biotin-BSA (A8549; Sigma-Aldrich) for 5 min. Chambers were washed with BRB80-1 mg/ml BSA and incubated with 0.5 mg/ml Neutravidin (A2666; Invitrogen) for 5 min. After wash with BRB80-1 mg/ml BSA, chambers were incubated with blocking buffer (BRB80, 1% Pluronic F-127, 1 mg/ml BSA) for 5 min. Imaging chambers were washed again with BRB80 containing 1 mg/ml BSA and incubated with GMPCPP-stabilized HiLyte-647-labeled seeds diluted in BRB80-1 mg/ml BSA for 15 min. Finally, MT polymerization was initiated by introducing a mix of 14 μ M tubulin, which contained 9% HiLyte-488-tubulin (TL488M; Cytoskeleton) in BRB80 supplemented with 1 mM GTP, 1 mg/ml BSA, 0.25% Pluronic F-127, 0.1% κ -casein, 0.2% methyl cellulose, and oxygen scavenging system (0.5 mg/ml glucose oxidase, 0.1 mg/ml catalase, 4.5 mg/ml D-glucose, 70 mM β -mercaptoethanol). For TIRF assays with EB1-GFP, GMPCPP-stabilized MT seeds and dynamic MT plus ends were distinguished respectively by brighter and dimmer HiLyte-647 intensity. MT polymerization reaction was additionally supplemented with 75 nM EB1-GFP and 50 mM KCl. Chambers were sealed with vacuum grease and imaging was performed at 37°C every 2 s for 15 min.

As a control, TIRF assays were performed in the presence of actin filaments. Actin filaments (28 μ M) were made by incubating 83% unlabeled G-actin (APHL99; Cytoskeleton) with 8.5% biotin-G-actin (AB07; Cytoskeleton) and 8.5% Rhodamine-G-actin (APHR; Cytoskeleton) in the presence of F-actin buffer (20 mM HEPES, pH 7.4, 100 mM KCl, 1 mM MgCl₂, 0.5 mM ATP, and 4 mM DTT) for 1 h at room temperature. Imaging chambers were treated as above but with the following differences: 1) chambers were not coated with an antibody, and 2) 50 nM of Rhodamine F-actin filaments were mixed with GMPCPP MT seeds, HiLyte-647-labeled, and incubated for 25 min. MT polymerization reaction was initiated as described above.

TIRF assays of septin binding to GMPCPP- versus GTP γ S-bound MTs

GMPCPP- and GTP γ S-stabilized MTs were prepared similarly to previous studies (Roth *et al.*, 2018). MTs were made by incubating 77% unlabeled tubulin (T240; Cytoskeleton) with 11.5% biotin-tubulin (T333P; Cytoskeleton) and 11.5% HiLyte-647-tubulin (TL670M; Cytoskeleton) or 11.5% HiLyte-488-tubulin (TL488M; Cytoskeleton) on ice for 30 min in the presence of 1 mM GMPCPP or GTP γ S (371545; EMD Millipore). Subsequently, tubulin mixes were placed at 37°C for 30 min. Then, 2 μ M of taxol was added to each mix and seeds were further incubated for an additional 5 min at 37°C. MT seeds were diluted in BRB80 containing 2 μ M taxol and centrifuged at 100,000 \times g (Optima TL100; Beckman Coulter) for 15 min. Pelleted MT seeds were resuspended in 60 μ l of BRB80-2 μ M taxol. Imaging chambers were incubated sequentially with 1% Pluronic F-127 for 5 min, 5 mg/ml Biotin-BSA (A8549; Sigma-Aldrich) for 5 min, 0.5 mg/ml Neutravidin (A2666; Invitrogen) for 5 min, ~60 nM of both GMPCPP and GTP γ S (mix 1:1) or ~60 nM of GTP γ S MT seeds supplemented with 2 μ M taxol for 15 min, and blocking buffer (BRB80, 1% Pluronic F-127, 1 mg/ml BSA, 2 μ M taxol) for 5 min. To test mCherry-SEPT2/6/7 binding to GMPCPP- and GTP γ S-stabilized MTs, increasing concentrations of

recombinant mCherry-SEPT2/6/7 were incubated for 15 min with GMPCPP and GTP γ S MTs in buffer containing BRB80, 0.1 mg/ml BSA, 0.1% Pluronic F-127, 0.1% κ -casein, oxygen scavenging system (0.5 mg/ml glucose oxidase, 0.1 mg/ml catalase, 4.5 mg/ml D-glucose, 70 mM β -mercaptoethanol), and 2 μ M taxol. Imaging chambers were sealed with vacuum grease and still images were taken after 15 min of incubation.

For the competition assay, 75 nM of EB1-GFP was incubated for 15 min with GTP γ S-stabilized MTs in the presence of increasing concentrations of recombinant mCherry-SEPT2/6/7 or nonfluorescent SEPT2/6/7 in a buffer containing BRB80, 0.1 mg/ml BSA, 0.1% Pluronic F-127, 0.1% κ -casein, 50 mM KCl, oxygen scavenging system (0.5 mg/ml glucose oxidase, 0.1 mg/ml catalase, 4.5 mg/ml D-glucose, 70 mM β -mercaptoethanol), and 2 μ M taxol. Imaging chambers were sealed with vacuum grease and still images were taken at 15 min of incubation.

Primary cultures and transfection

Primary rat hippocampal cultures were obtained from the Neuron Culture Service Center (University of Pennsylvania). Hippocampi were dissected from embryonic day 18 pups of mixed genders, obtained from timed pregnant Sprague Dawley rats. Cells were dissociated by incubation in trypsin-containing media. Cells were plated in Neurobasal medium supplemented with 2% B27 supplement (17504044; ThermoFisher). Cells were cultured in 24-well plates on 12-mm round glass coverslips (Bellco Glass) coated with 1 mg/ml poly-L-lysine (OKK-3056; Peptides International). After 1 d in culture, cytosine β -D-arabinofuranoside (C6645; Sigma) was added at a final concentration of 1.5 μ M to inhibit glial proliferation. Cells were maintained at 37°C in a 5% CO₂ incubator. Neurons were cultured for 1–4 d in vitro (DIV) and transfected with Lipofectamine 3000 (L3000; Invitrogen) for 48 h (DIV 3–6) before live imaging.

Imaging of rat hippocampal neurons

Live-cell imaging of hippocampal neurons was performed on DIV 3–6, 48 h after transfection. Coverslips were mounted on bottomless 35-mm dishes with a 7-mm microwell using silicone vacuum grease (Beckman Coulter). Conditioned media from the coverslip's well, along with 2 ml of phenol red-free Neurobasal media, were supplemented with 2% B27 (17504044; ThermoFisher) and 30 mM HEPES (H0887; Sigma). Dishes were sealed with parafilm. Cells were selected for moderate expression of GFP-SEPT7 or GFP-F-tractin. Neurons were imaged at 30 frames/min for 3 min using total internal reflection fluorescence (TIRF) imaging at 37°C with the TIRF module of the DeltaVision OMX V4 imaging platform.

All TIRF microscopy assays were performed with the DeltaVision OMX V4 inverted microscope (GE Healthcare) equipped with 60 \times /1.49 NA TIRF objective lens (Olympus), motorized stage, sCMOS pco.edge cameras (PCO), stage top incubator with temperature controller, and the softWoRx software.

Image and statistical analyses

Quantifications of MT dynamics and fluorescence intensities were performed in ImageJ/Fiji. Time-lapse image series were corrected for bleaching using the bleach correction plug-in. Kymograph analysis was used to quantify MT dynamics with the KymographBuilder plug-in. Growth and depolymerization rates were derived by manual segmentation of the trajectories of growing and shrinking MT plus ends, respectively. The velocity measurement tool macro was used to calculate rates of polymerization and depolymerization, which excluded the durations of any pausing events. Catastrophe frequencies were measured by dividing the total number of catastrophe events by

the total time spent in growing and pausing. Pauses were defined as events without change in the MT length for at least 40 s or longer.

Still images at 15 min of incubation were used to quantify the number of tubulin puncta bound to immobilized recombinant SEPT2/6/7 or SEPT9 (Supplemental Figure S2). Background fluorescence was removed using the subtract background tool (rolling ball radius, 50 pixels) in Fiji. Manual fluorescence thresholding was applied and the analyze particles tool was used to count the number of tubulin puncta.

Quantification of the fluorescence intensity of EB1-GFP (Figure 4B and Supplemental Figure S3B) was performed as previously described (Ramirez-Rios et al., 2016). Background fluorescence was removed with the subtract background tool (rolling ball radius, 20 pixels) and fluorescence intensities of EB1-GFP were measured from kymographs. Segmented lines of a 5-pixel width were drawn manually at the MT plus-end tips and maximum fluorescence intensities were measured.

Fluorescence intensities of mCherry-SEPT2/6/7 were measured from still images at 15 min (Figures 3D and 4D). Segmented lines of equal width (5 pixels) were drawn manually along the entire length of GMPCPP MT seeds and the entire length of the lattice of plus-end segments (Figure 3D) or along the GMPCPP and GTP γ S MT seeds (Figure 4D). The average pixel intensity was derived after subtraction of the average pixel intensity of background fluorescence, which was quantified by taking the mean value from three regions that did not contain any MTs. The same method was used to measure the average pixel intensity of EB1-GFP on GTP γ S MT seeds (Figure 4F and Supplemental Figure S3E) in the presence of increasing concentrations of mCherry-SEPT2/6/7 or nonfluorescent SEPT2/6/7. Line scans (intensity vs. distance plots) in Figure 3C were generated using the plot profile function in Fiji.

Analysis of EB3-tdTomato comet interaction with GFP-SEPT7 or GFP-F-tractin filaments in rat hippocampal neurons was done by making kymographs of neurites from time-lapse TIRF imaging movies using the Segmented Line (15-pixel width) and Kymograph-Builder tools in Fiji. Kymographs were treated as regions of interest and filaments to track comet pausing or procession were selected from these regions. All comets passing at GFP-F-tractin or GFP-SEPT7 filament were tracked and the number of comets that passed or paused is recorded. A pause in these movies is defined as a halt in comet movement lasting longer than 3 frames (6 s). The difference in the rates of EB3-tdTomato movement (Δ Velocity) between before and after pausing at GFP-SEPT7 or GFP-F-tractin was calculated by measuring the rate of movement in the frames immediately before pausing and after resuming growth. For each event of acceleration ($V_{\text{after}} > V_{\text{before}}$) and deceleration ($V_{\text{after}} < V_{\text{before}}$), the slower rate was subtracted from the faster and mean Δ Velocity (\pm SEM) was derived (Figure 5K).

Statistical analysis was performed using GraphPad Prism 6 software. Each data set was tested for normal distribution using the Kolmogorov–Smirnov test. Mean, SEM, SD, and *p* values were derived using Student's *t* test for normally distributed data and the Mann–Whitney *U* test for nonnormally distributed data. Analysis of variance (ANOVA) was used for multiple comparison of groups. For normally distributed data, ordinary one-way ANOVA was used followed by the Tukey's multiple comparison test. For nonnormally distributed data, the Kruskal–Wallis *H* test was used followed by the Dunn's multiple comparison test.

ACKNOWLEDGMENTS

We thank Shae Padrick (College of Medicine, Drexel University), Amy Gladfelter (University of North Carolina), Antonina Roll-Mecak

(Cell Biology and Biophysics, NIH), Anna Akhmanova (Utrecht University), Smita Yadav (University of Washington), and Tatyana Svitkina (Department of Biology, University of Pennsylvania) for plasmids. All imaging was performed at Drexel University's Cell Imaging Center. This work was supported by NIH/National Institute of General Medical Sciences grant GM097664 to E.T.S.

REFERENCES

- Ageta-Ishihara N, Miyata T, Ohshima C, Watanabe M, Sato Y, Hamamura Y, Higashiyama T, Mazitschek R, Bito H, Kinoshita M (2013). Septins promote dendrite and axon development by negatively regulating microtubule stability via HDAC6-mediated deacetylation. *Nat Commun* 4, 2532.
- Akil A, Peng J, Omrane M, Gondeau C, Desterke C, Marin M, Tronchere H, Taveneau C, Sar S, Briolotti P, et al. (2016). Septin 9 induces lipid droplets growth by a phosphatidylinositol-5-phosphate and microtubule-dependent mechanism hijacked by HCV. *Nat Commun* 7, 12203.
- Amin ND, Zheng YL, Kesavapany S, Kanungo J, Guszczynski T, Sihag RK, Rudrabhatla P, Albers W, Grant P, Pant HC (2008). Cyclin-dependent kinase 5 phosphorylation of human septin SEPT5 (hCDCrel-1) modulates exocytosis. *J Neurosci* 28, 3631–3643.
- Bai X, Bowen JR, Knox TK, Zhou K, Pendziwiat M, Kuhlenbaumer G, Sindelar CV, Spiliotis ET (2013). Novel septin 9 repeat motifs altered in neurogenic amyotrophy bind and bundle microtubules. *J Cell Biol* 203, 895–905.
- Bieling P, Kandels-Lewis S, Telley IA, van Dijk J, Janke C, Surrey T (2008). CLIP-170 tracks growing microtubule ends by dynamically recognizing composite EB1/tubulin-binding sites. *J Cell Biol* 183, 1223–1233.
- Bieling P, Laan L, Schek H, Munteanu EL, Sandblad L, Dogterom M, Brunner D, Surrey T (2007). Reconstitution of a microtubule plus-end tracking system in vitro. *Nature* 450, 1100–1105.
- Bowen JR, Hwang D, Bai X, Roy D, Spiliotis ET (2011). Septin GTPases spatially guide microtubule organization and plus end dynamics in polarizing epithelia. *J Cell Biol* 194, 187–197.
- Bridges AA, Jentsch MS, Oakes PW, Occhipinti P, Gladfelter AS (2016). Micron-scale plasma membrane curvature is recognized by the septin cytoskeleton. *J Cell Biol* 213, 23–32.
- Brouhard GJ, Rice LM (2018). Microtubule dynamics: an interplay of biochemistry and mechanics. *Nat Rev Mol Cell Biol* 19, 451–463.
- Cannon KS, Woods BL, Crutchley JM, Gladfelter AS (2019). An amphipathic helix enables septins to sense micrometer-scale membrane curvature. *J Cell Biol* 218, 1128–1137.
- Colin A, Singaravelu P, Thery M, Blanchoin L, Guerou Z (2018). Actin-network architecture regulates microtubule dynamics. *Curr Biol* 28, 2647–2656.e2644.
- Dixit R, Barnett B, Lazarus JE, Tokito M, Goldman YE, Holzbaur EL (2009). Microtubule plus-end tracking by CLIP-170 requires EB1. *Proc Natl Acad Sci USA* 106, 492–497.
- Dolat L, Hunyara JL, Bowen JR, Karasmanis EP, Elgawly M, Galkin VE, Spiliotis ET (2014). Septins promote stress fiber-mediated maturation of focal adhesions and renal epithelial motility. *J Cell Biol* 207, 225–235.
- Dolat L, Spiliotis ET (2016). Septins promote macropinosome maturation and traffic to the lysosome by facilitating membrane fusion. *J Cell Biol* 214, 517–527.
- Estey MP, Di Ciano-Oliveira C, Froese CD, Bejide MT, Trimble WS (2010). Distinct roles of septins in cytokinesis: SEPT9 mediates midbody abscission. *J Cell Biol* 191, 741–749.
- Fliege M, Kahle A, Haffner K, Zieger B (2014). Distinct localization of septin proteins to ciliary sub-compartments in airway epithelial cells. *Biol Chem* 395, 151–156.
- Ghossoub R, Hu Q, Failler M, Rouyez MC, Spitzbarth B, Mostowy S, Wolfrum U, Saunier S, Cossart P, Jamesnelson W, Benmerah A (2013). Septins 2, 7 and 9 and MAP4 colocalize along the axoneme in the primary cilium and control ciliary length. *J Cell Sci* 126, 2583–2594.
- Hein MY, Hubner NC, Poser I, Cox J, Nagaraj N, Toyoda Y, Gak IA, Weisswange I, Mansfeld J, Buchholz F, et al. (2015). A human interactome in three quantitative dimensions organized by stoichiometries and abundances. *Cell* 163, 712–723.
- Honnappa S, Gouveia SM, Weisbrich A, Damberger FF, Bhavesh NS, Jawhari H, Grigoriev I, van Rijssel FJ, Buey RM, Lawera A, et al. (2009). An EB1-binding motif acts as a microtubule tip localization signal. *Cell* 138, 366–376.
- Hu J, Bai X, Bowen JR, Dolat L, Korobova F, Yu W, Baas PW, Svitkina T, Gallo G, Spiliotis ET (2012). Septin-driven coordination of actin and microtubule remodeling regulates the collateral branching of axons. *Curr Biol* 22, 1109–1115.
- Hu Q, Milenkovic L, Jin H, Scott MP, Nachury MV, Spiliotis ET, Nelson WJ (2010). A septin diffusion barrier at the base of the primary cilium maintains ciliary membrane protein distribution. *Science* 329, 436–439.
- Ihara M, Tomimoto H, Kitayama H, Morioka Y, Akiguchi I, Shibasaki H, Noda M, Kinoshita M (2003). Association of the cytoskeletal GTP-binding protein Sept4/H5 with cytoplasmic inclusions found in Parkinson's disease and other synucleinopathies. *J Biol Chem* 278, 24095–24102.
- Ihara M, Yamasaki N, Hagiwara A, Tanigaki A, Kitano A, Hikawa R, Tomimoto H, Noda M, Takanashi M, Mori H, et al. (2007). Sept4, a component of presynaptic scaffold and Lewy bodies, is required for the suppression of alpha-synuclein neurotoxicity. *Neuron* 53, 519–533.
- Inoue D, Obino D, Pineau J, Farina F, Gaillard J, Guerin C, Blanchoin L, Lennon-Dumenil AM, Thery M (2019). Actin filaments regulate microtubule growth at the centrosome. *EMBO J* 38, e99630.
- Janke C, Montagnac G (2017). Causes and consequences of microtubule acetylation. *Curr Biol* 27, R1287–R1292.
- Jiang K, Toedt G, Montenegro Gouveia S, Davey NE, Hua S, van der Vaart B, Grigoriev I, Larsen J, Pedersen LB, Bezstarosti K, et al. (2012). A Proteome-wide screen for mammalian SxIP motif-containing microtubule plus-end tracking proteins. *Curr Biol* 22, 1800–1807.
- Johnson HW, Schell MJ (2009). Neuronal IP3 3-kinase is an F-actin-bundling protein: role in dendritic targeting and regulation of spine morphology. *Mol Biol Cell* 20, 5166–5180.
- Karasmanis EP, Phan CT, Angelis D, Kesisova IA, Hoogenraad CC, McKenney RJ, Spiliotis ET (2018). Polarity of neuronal membrane traffic requires sorting of kinesin motor cargo during entry into dendrites by a microtubule-associated septin. *Dev Cell* 46, 204–218 e207.
- Kim MS, Froese CD, Estey MP, Trimble WS (2011). SEPT9 occupies the terminal positions in septin octamers and mediates polymerization-dependent functions in abscission. *J Cell Biol* 195, 815–826.
- Kinoshita M (2003a). Assembly of mammalian septins. *J Biochem* 134, 491–496.
- Kinoshita M (2003b). The septins. *Genome Biol* 4, 236.
- Kinoshita A, Kinoshita M, Akiyama H, Tomimoto H, Akiguchi I, Kumar S, Noda M, Kimura J (1998). Identification of septins in neurofibrillary tangles in Alzheimer's disease. *Am J Pathol* 153, 1551–1560.
- Kremer BE, Haystead T, Macara IG (2005). Mammalian septins regulate microtubule stability through interaction with the microtubule-binding protein MAP4. *Mol Biol Cell* 16, 4648–4659.
- Lazarus JE, Moughamian AJ, Tokito MK, Holzbaur EL (2013). Dynactin subunit p150(Glued) is a neuron-specific anti-catastrophe factor. *PLoS Biol* 11, e1001611.
- Maurer SP, Bieling P, Cope J, Hoenger A, Surrey T (2011). GTPgammaS microtubules mimic the growing microtubule end structure recognized by end-binding proteins (EBs). *Proc Natl Acad Sci USA* 108, 3988–3993.
- Maurer SP, Fourniol FJ, Bohner G, Moores CA, Surrey T (2012). EBs recognize a nucleotide-dependent structural cap at growing microtubule ends. *Cell* 149, 371–382.
- Mavrakakis M, Azou-Gros Y, Tsai FC, Alvarado J, Bertin A, Iv F, Kress A, Bras-selet S, Koenderink GH, Lecuit T (2014). Septins promote F-actin ring formation by crosslinking actin filaments into curved bundles. *Nat Cell Biol* 16, 322–334.
- Menon MB, Sawada A, Chaturvedi A, Mishra P, Schuster-Gossler K, Galla M, Schambach A, Gossler A, Forster R, Heuser M, et al. (2014). Genetic deletion of SEPT7 reveals a cell type-specific role of septins in microtubule destabilization for the completion of cytokinesis. *PLoS Genet* 10, e1004558.
- Merrim EB, Millette M, Lombard DC, Saengsawang W, Fothergill T, Hu X, Ferhat L, Dent EW (2013). Synaptic regulation of microtubule dynamics in dendritic spines by calcium, F-actin, and drebrin. *J Neurosci* 33, 16471–16482.
- Mostowy S, Cossart P (2012). Septins: the fourth component of the cytoskeleton. *Nat Rev Mol Cell Biol* 13, 183–194.
- Nakos K, Rosenberg M, Spiliotis ET (2019). Regulation of microtubule plus end dynamics by septin 9. *Cytoskeleton (Hoboken)* 76, 83–91.
- Nolke T, Schwan C, Lehmann F, Ostevold K, Pertz O, Aktories K (2016). Septins guide microtubule protrusions induced by actin-depolymerizing toxins like Clostridium difficile transferase (CDT). *Proc Natl Acad Sci USA* 113, 7870–7875.
- Ostevold K, Melendez AV, Lehmann F, Schmidt G, Aktories K, Schwan C (2017). Septin remodeling is essential for the formation of cell

- membrane protrusions (microtentacles) in detached tumor cells. *Oncotarget* 8, 76686–76698.
- Ramirez-Rios S, Denarier E, Prezel E, Vinit A, Stoppin-Mellet V, Devred F, Barbier P, Peyrot V, Sayas CL, Avila J, et al. (2016). Tau antagonizes end-binding protein tracking at microtubule ends through a phosphorylation-dependent mechanism. *Mol Biol Cell* 27, 2924–2934.
- Reid TA, Schuster BM, Mann BJ, Balchand SK, Plooster M, McClellan M, Coombes CE, Wadsworth P, Gardner MK (2016). Suppression of microtubule assembly kinetics by the mitotic protein TPX2. *J Cell Sci* 129, 1319–1328.
- Roth D, Fitton BP, Chmel NP, Wasiluk N, Straube A (2018). Spatial positioning of EB family proteins at microtubule tips involves distinct nucleotide-dependent binding properties. *J Cell Sci* 132, jcs219550.
- Russell SE, Hall PA (2011). Septin genomics: a road less travelled. *Biol Chem* 392, 763–767.
- Sellin ME, Sandblad L, Stenmark S, Gullberg M (2011). Deciphering the rules governing assembly order of mammalian septin complexes. *Mol Biol Cell* 22, 3152–3164.
- Sellin ME, Stenmark S, Gullberg M (2012). Mammalian SEPT9 isoforms direct microtubule-dependent arrangements of septin core heteromers. *Mol Biol Cell* 23, 4242–4255.
- Sellin ME, Stenmark S, Gullberg M (2014). Cell type-specific expression of SEPT3-homology subgroup members controls the subunit number of heteromeric septin complexes. *Mol Biol Cell* 25, 1594–1607.
- Sheffield PJ, Oliver CJ, Kremer BE, Sheng S, Shao Z, Macara IG (2003). Borg/septin interactions and the assembly of mammalian septin heterodimers, trimers, and filaments. *J Biol Chem* 278, 3483–3488.
- Shehadeh L, Mitsi G, Adi N, Bishopric N, Papapetropoulos S (2009). Expression of Lewy body protein septin 4 in postmortem brain of Parkinson's disease and control subjects. *Mov Disord* 24, 204–210.
- Shindo A, Audrey A, Takagishi M, Takahashi M, Wallingford JB, Kinoshita M (2018). Septin-dependent remodeling of cortical microtubule drives cell reshaping during epithelial wound healing. *J Cell Sci* 131, jcs212647.
- Sirajuddin M, Farkasovsky M, Hauer F, Kuhlmann D, Macara IG, Weyand M, Stark H, Wittinghofer A (2007). Structural insight into filament formation by mammalian septins. *Nature* 449, 311–315.
- Spiliotis ET (2010). Regulation of microtubule organization and functions by septin GTPases. *Cytoskeleton (Hoboken)* 67, 339–345.
- Spiliotis ET (2018). Spatial effects—site-specific regulation of actin and microtubule organization by septin GTPases. *J Cell Sci* 131, jcs207555.
- Spiliotis ET, Hunt SJ, Hu Q, Kinoshita M, Nelson WJ (2008). Epithelial polarity requires septin coupling of vesicle transport to polyglutamylated microtubules. *J Cell Biol* 180, 295–303.
- Spiliotis ET, Kinoshita M, Nelson WJ (2005). A mitotic septin scaffold required for Mammalian chromosome congression and segregation. *Science* 307, 1781–1785.
- Steels JD, Estey MP, Froese CD, Reynaud D, Pace-Asciak C, Trimble WS (2007). Sept12 is a component of the mammalian sperm tail annulus. *Cell Motil Cytoskeleton* 64, 794–807.
- Tanenbaum ME, Vale RD, McKenney RJ (2013). Cytoplasmic dynein cross-links and slides anti-parallel microtubules using its two motor domains. *Elife* 2, e00943.
- Targa B, Klipfel L, Cantaloube I, Salameh J, Benoit B, Pous C, Baillet A (2019). Septin filament coalignment with microtubules depends on SEPT9_i1 and tubulin polyglutamylation, and is an early feature of acquired cell resistance to paclitaxel. *Cell Death Dis* 10, 54.
- Vemu A, Szczesna E, Zehr EA, Spector JO, Grigorieff N, Deaconescu AM, Roll-Mecak A (2018). Severing enzymes amplify microtubule arrays through lattice GTP-tubulin incorporation. *Science* 361, eaau1504.
- Vitre B, Coquelle FM, Heichette C, Garnier C, Chretien D, Arnal I (2008). EB1 regulates microtubule dynamics and tubulin sheet closure in vitro. *Nat Cell Biol* 10, 415–421.
- Weirich CS, Erzberger JP, Barral Y (2008). The septin family of GTPases: architecture and dynamics. *Nat Rev Mol Cell Biol* 9, 478–489.
- Yadav S, Osos-Prieto JA, Peters CJ, Zhou J, Pleasure SJ, Burlingame AL, Jan LY, Jan YN (2017). TAOK2 kinase mediates PSD95 stability and dendritic spine maturation through Septin7 phosphorylation. *Neuron* 93, 379–393.
- Yang C, Svitkina TM (2019). Ultrastructure and dynamics of the actin-myosin II cytoskeleton during mitochondrial fission. *Nat Cell Biol* 21, 603–613.
- Zanic M, Stear JH, Hyman AA, Howard J (2009). EB1 recognizes the nucleotide state of tubulin in the microtubule lattice. *PLoS One* 4, e7585.
- Zanic M, Widlund PO, Hyman AA, Howard J (2013). Synergy between XMAP215 and EB1 increases microtubule growth rates to physiological levels. *Nat Cell Biol* 15, 688–693.
- Zhang J, Kong C, Xie H, McPherson PS, Grinstein S, Trimble WS (1999). Phosphatidylinositol polyphosphate binding to the mammalian septin H5 is modulated by GTP. *Curr Biol* 9, 1458–1467.