

Corrigendum

Corrigendum to “Factors Associated with the Incidence and Severity of New-Onset Atrial Fibrillation in Adult Critically Ill Patients”

Péricles A. D. Duarte ¹, **Gustavo Elias Leichtweis**², **Luiza Andriolo**³, **Yasmim A. Delevatti**³, **Amaury C. Jorge**⁴, **Andreia C. Fumagalli**⁵, **Luiz Claudio Santos**⁶, **Cecilia K. Miura**⁷, and **Sergio K. Saito**⁵

¹Hospital do Câncer (UOPECCAN) and Hospital Universitário and Hospital São Lucas, Cascavel, PR, Brazil

²Hospital São Lucas, Cascavel, PR, Brazil

³Curso de Medicina, Faculdade Assis Gurgacz, Cascavel, PR, Brazil

⁴Hospital Universitário, Cascavel, PR, Brazil

⁵Hospital Bom Jesus, Toledo, PR, Brazil

⁶Hospital Nossa Senhora Salete, Cascavel, PR, Brazil

⁷Hospital Costa Cavalcanti, Foz do Iguaçu, PR, Brazil

Correspondence should be addressed to Péricles A. D. Duarte; pericles.duarte@unioeste.br

Received 24 October 2018; Accepted 4 November 2018; Published 29 January 2019

Copyright © 2019 Péricles A. D. Duarte et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Factors Associated with the Incidence and Severity of New-Onset Atrial Fibrillation in Adult Critically Ill Patients” [1], the sentence “However, the AF in non-cardiac surgery patients was also high (10.2%)” in the Discussion should be rephrased to “However, the incidence of AF in all patients (medical or surgical) after excluding those who were not submitted to cardiac surgery was also high (10.2%)” to avoid any misinterpretation.

References

- [1] P. A. D. Duarte, G. E. Leichtweis, L. Andriolo et al., “Factors associated with the incidence and severity of new-onset atrial fibrillation in adult critically ill patients,” *Critical Care Research and Practice*, vol. 2017, Article ID 8046240, 8 pages, 2017.