

RESEARCH ARTICLE

Open Access

Differentiation in MALDI-TOF MS and FTIR spectra between two closely related species *Acidovorax oryzae* and *Acidovorax citrulli*

Yanli Wang^{1,2}, Qing Zhou¹, Bin Li^{1*}, Baoping Liu¹, Guoxing Wu³, Muhammad Ibrahim¹, Guanlin Xie¹, Hongye Li¹ and Guochang Sun²

Abstract

Background: Two important plant pathogenic bacteria *Acidovorax oryzae* and *Acidovorax citrulli* are closely related and often not easy to be differentiated from each other, which often resulted in a false identification between them based on traditional methods such as carbon source utilization profile, fatty acid methyl esters, and ELISA detection tests. MALDI-TOF MS and Fourier transform infrared (FTIR) spectra have recently been successfully applied in bacterial identification and classification, which provide an alternate method for differentiating the two species.

Results: Characterization and comparison of the 10 *A. oryzae* strains and 10 *A. citrulli* strains were performed based on traditional bacteriological methods, MALDI-TOF MS, and FTIR spectroscopy. Our results showed that the identity of the two closely related plant pathogenic bacteria *A. oryzae* and *A. citrulli* was able to be confirmed by both pathogenicity tests and species-specific PCR, but the two species were difficult to be differentiated based on Biolog and FAME profile as well as 16 S rRNA sequence analysis. However, there were significant differences in MALDI-TOF MS and FTIR spectra between the two species of *Acidovorax*. MALDI-TOF MS revealed that 22 and 18 peaks were specific to *A. oryzae* and *A. citrulli*, respectively, while FTIR spectra of the two species of *Acidovorax* have the specific peaks at 1738, 1311, 1128, 1078, 989 cm^{-1} and at 1337, 968, 933, 916, 786 cm^{-1} , respectively.

Conclusions: This study indicated that MALDI-TOF MS and FTIR spectra may give a new strategy for rapid bacterial identification and differentiation of the two closely related species of *Acidovorax*.

Background

Many researches have focused on the specific detection of the two important plant pathogenic bacteria *Acidovorax oryzae* (formerly *Acidovorax avenae* subsp. *avenae*) and *Acidovorax citrulli* (formerly *Acidovorax avenae* subsp. *citrulli*) [1,2]. However, the two species are closely related and often not easy to be differentiated from each other [3,4], which often resulted in a false identification between them based on traditional methods such as carbon source utilization profile, fatty acid methyl esters, PCR and ELISA detection tests [1,5]. Therefore, it is necessary to develop an alternate method for differentiating the two species.

Recently, MALDI-TOF MS and Fourier transform infrared (FTIR) spectra have been successfully applied in bacterial identification and classification [6-11]. MALDI-TOF MS allows bacterial identification at the species level by measuring molecular masses of proteins and other bacterial components obtained from whole bacterial extracts, while FTIR spectroscopy allows the analysis of small quantities of biomass, simultaneous characterization of different functional groups such as lipids, proteins, nucleic acids and polysaccharides in biological molecules and complex structures and without disturbing the systems, and requires no consumables or reagents [6,12,13]. However, little information was obtained about the applications of MALDI-TOF MS and FTIR spectra in plant pathogenic bacteria.

The objective of this study was to examine and compare the MALDI-TOF MS and FTIR spectra of bacteria from the two species of *Acidovorax*.

* Correspondence: libin0571@zju.edu.cn

¹State Key Laboratory of Rice Biology, Institute of Biotechnology, Zhejiang University, Hangzhou 310029, China

Full list of author information is available at the end of the article

Methods

Bacterial strains

The 10 virulent strains of *A. oryzae* used in this study were isolated from diseased rice seed and seedling, while the 10 virulent strains of *A. citrulli* were isolated from diseased watermelon and melon (Table 1). The identities of bacterial strains were determined and confirmed based on the biochemical and physiological characteristics as described by Krieg and Holt [14] and Schaad et al. [15], whole-cell fatty acid and Biolog metabolic assays as described by Li et al. [3,16,17], species-specific PCR [1,15,18] and 16 S ribosomal RNA gene sequence analysis [3,16,17]. The representative *A. oryzae* strain R1001 (Collection no: ACCC05733) and *A. citrulli* strain Ab1 (Collection no: ACCC05732) were deposited in Agricultural Culture Collection of China (ACCC).

MALDI-TOF MS

Sample preparation

One loop of bacterial cells grown on Luria-Bertani at 30°C for 48 h was suspended in 300 µl of Millipore water followed by adding 900 µl of absolute ethanol. Cell pellets were obtained by a centrifugation at 12000 rpm for 2 min and suspended in 50 µl of formic acid (70% v/v) followed by carefully adding 50 µl of acetonitrile. One microliter of supernatant after a centrifugation at 12000 rpm for 2 min was spotted on a steel target plate (Bruker Daltonic, Billerica, Massachusetts) and air dried at room temperature. Afterwards, 1 µl of matrix solution

(saturated solution of α -cyanohydroxycinnamic acid in 50% aqueous acetonitrile containing 2.5% trifluoroacetic acid) was quickly added onto the surface of each sample spot. Samples were prepared in duplicate.

MALDI-TOF MS analysis

Mass spectrometric measurements were performed with an AUTOFLEX Analyzer (Bruker Daltonics) as described in previous studies using the linear positive ion extraction [10,11,19]. The method of identification included the m/z from 2 to 12 kDa. *Escherichia coli* DH5 α was used as an external protein calibration mixture followed by the Bruker Test Standard [20]. Raw mass spectrum smooth, baseline correction and peak detection were performed using the corresponding programs installed in the MS system. Resulting mass fingerprints were exported to FLEX ANALYSIS (Bruker Daltonics) and analyzed. Spectral data were investigated for the presence of biomarkers characteristic for each of the two *Acidovorax* species. After visual inspection and comparison, the most intensive and predominantly present protein peaks were selected and screened in representatives of each species.

FTIR spectroscopy

Sample preparation

Bacterial cells were collected from overnight Luria-Bertani broth culture grown at 28°C by centrifugation at 10,000 rpm for 10 min. After removing the supernatants, the bacterial pellets were washed twice with double distilled water. After second wash in double distilled water, bacterial samples were stored at -70°C until lyophilisation. The samples for FTIR analysis were first grounded into fine particles using mortar and pestle. The 1 mg of each sample was then mixed with 100 mg potassium bromide (KBr) which extensively dried in microfuge tubes using a lyophiliser. These mixtures have been dried for an additional 2 h in the same microfuge tubes. The KBr based pellets were then compressed into a thin disk by establishing pressure of 100 kg/cm² (1200 psi) for about 8 min.

FTIR spectroscopy and data analysis

The FTIR spectroscopy data were analysed as previously described by Garip et al. [21] with a small modification. Pellets were scanned at 4 cm⁻¹ resolution with 100 scans in the spectral range of 4000–500 cm⁻¹ at room temperature. The sample compartment in the FTIR spectrometer was continuously purged with dry air to prevent water vapour. Analysis of the spectral data was performed by using Grams 32 (Galactic Industries, Salem, NH, USA) software. The spectral range of 4000–500 cm⁻¹ was analyzed. The band positions were measured according to the center of weight. The averages of

Table 1 Strains of *Acidovorax oryzae* (Ao) and *Acidovorax citrulli* (Ac) used in this study

Ao strains	Sources	Ac strains	Sources
R1001	Rice seedling, this lab	A1	Watermelon leaf, CAAS, China
R1002	Rice seedling, this lab	Aacf	Watermelon leaf, FAFFU, China
R1003	Rice seedling, this lab	Ab1	Watermelon leaf, this lab
R1004	Rice seedling, this lab	Njf4	Watermelon leaf, NAU, China
CB97012	Rice seeds, this lab	Ps96	Watermelon leaf, CAAS, China
CB97058	Rice seeds, this lab	Ab3	Melon leaf, this lab
CB97063	Rice seeds, this lab	Tw20	Melon leaf, CAAS, China
CB97181	Rice seeds, this lab	Ab5	Melon leaf, this lab
CB97095	Rice seeds, this lab	Ab8	Melon leaf, this lab
CB97128	Rice seeds, this lab	Ab9	Melon leaf, this lab

CAAS: Chinese Academy of Agricultural Sciences; FAFFU: Fujian Agricultural and Forestry University; NAU: Nanjing Agricultural University.

the spectra belonging to the same experimental groups, baseline correction, normalisation and the band areas were obtained by using the same software program. The average spectra and normalisation process were applied only for visual representation of the differences, however for the determination of the spectral parameters and calculation of mean values and statistical analysis each baseline corrected original spectrum was taken into consideration.

Statistics

The software STATGRAPHICS Plus, version 4.0 (Copyright Manugistics Inc., Rockville, Md., USA) was used to perform the statistical analysis. Levels of significance ($p < 0.05$) of main treatments and their interactions were calculated by analysis of variance after testing for normality and variance homogeneity.

Results and discussion

Bacterial identity

Results from this study indicated the rice strains should be identified as *A. oryzae* with Biolog similarity of 0.72 to 0.73, FAME similarity of 0.73 to 0.74, 16 S rRNA sequence similarity of 99% and confirmed by both pathogenicity tests and species-specific PCR, while the watermelon and melon strains should be identified as *A. citrulli* with Biolog similarity of 0.70 to 0.73, FAME similarity of 0.73 to 0.74, 16 S rRNA sequence similarity of 99%, and confirmed by both pathogenicity tests and

species-specific PCR in the newly proposed classification of subspecies of *A. avenae*. However, in general, the two species of *Acidovorax* were high similar, and difficult to be differentiated based on Biolog and FAME profile as well as 16 S rRNA sequence analysis.

MALDI-TOF MS

MALDI-TOF MS characterization of bacteria is based on differences in mass to charge ratio (m/z) fingerprints of whole cell proteins, mainly representing ribosomal proteins which are most abundantly expressed under all growth conditions [22]. In this type of mass spectrometry, samples were prepared by embedding analyte molecules in a crystal matrix of small acidic molecules. A brief laser pulse irradiates the sample and the matrix absorbs the laser energy resulting in ablation of a small volume of matrix and desorption of the embedded analyte molecules which are ionized. Subsequently, predominantly single charged analyte ions can be detected and analyzed [23].

Figure 1 presents a typical MALDI-TOF MS spectrum for the two species, which contain a contiguous sequence of about high-intensity ion peaks between 2000 and 12,000 Da. The obtained spectral profiles were further screened for the presence of recurring peaks or biomarker ions specific for both the species. Fifty selected m/z values were summarized in Table 2, while ten m/z values were detected in both species, making them characteristic for the genus *Acidovorax*. In addition, MALDI-




Figure 1 MALDI-TOF MS protein mass fingerprints of *Acidovorax oryzae* and *Acidovorax citrulli*. Similar and different marker masses for the identification of *A. oryzae* and *A. citrulli* are listed in Table 2. Intensity of ions is shown on the y axis and the mass (in Daltons) of the ions is shown on the x axis. The m/z values represent mass-to-charge ratios. *: Unique peaks positions for each of species.

Table 2 Characteristic MALDI-TOF masses (in Daltons) selected as possible biomarkers for identification of *Acidovorax oryzae* (Ao) and *Acidovorax citrulli* (Ac)

Ao	Ac	Ao	Ac
2178			6703
2568	2565		6845
2932	2930		7055
3169	3168	7067	
3281	3285	7349	
	3524		7461
3533		8387	8381
3675		8486	
	3729		8494
4184			8636
4353	4351	8642	
	4716	8709	
4777		9181	
4885		9545	
	4956		9503
4965		9746	
5135	5133		9919
5304	5305	9935	
	5674	10097	
5863	5861	10260	
6339	6337		10271
	6413	10503	
6420			10608
	6550	10609	
6568			11349

Masses observed in both species are marked in bold while species unique mass values marked in Figure 1. Assigned proteins calculated using RMIDb.

TOF MS revealed that 22 and 18 peaks were specific to *A. oryzae* and *A. citrulli*, respectively (Table 2, Figure 1). These unique peaks for each species offer a strong proof in differentiating the two species. This result is consistent with the review of Moore et al. [24], which found that MALDI-TOF MS is a valuable and reliable tool for microbial identification in a number of studies.

FTIR spectroscopy

In agreement with the result of bacteriological characterization, the 10 strains of *A. oryzae* had a very similar FTIR spectrum while the 10 strains of *A. citrulli* had a very similar FTIR spectrum regardless of bacterial origin (data not shown), indicating the stability and reliability of the FTIR spectroscopic system. In addition, characterization and differentiation of the two species were performed based on the average FTIR spectrum of

the 10 *A. oryzae* strains and the 10 *A. citrulli* strains, respectively (Figure 2). In general, there were significant differences in both the frequency and the intensity values of the 10 main functional groups between the two species except the frequency of PO₂ asymmetric stretching (Table 3), which indicated that the method of FTIR spectrum maybe have a higher level of differentiation between the two species compared to the biochemical and physiological characteristics tested in this study.

The average spectra in the 4000–500 cm⁻¹ region indicated that the *A. oryzae* strains have a higher frequency of the CH₃ asymmetric stretching vibration at 2959 cm⁻¹, the CH₂ asymmetric stretching vibration at 2927 cm⁻¹, the Amide I band at 1657 cm⁻¹, Amide II band at 1541 cm⁻¹, and the CH₂ bending band at 1452 cm⁻¹ compared to the *A. citrulli* strains, while the *A. citrulli* strains have a higher frequency of the CH₃ symmetric stretching vibration at 2876 cm⁻¹, the CH₂ symmetric stretching vibration at 2857 cm⁻¹, the COO⁻ symmetric stretch band at 1391 cm⁻¹ and the PO₂ symmetric stretching; phospholipids C-O stretch band at 1080 cm⁻¹ compared to the *A. oryzae* strains (Figure 2; Table 3; Additional file 1). In addition, the *A. oryzae* strains have a higher intensity of the absorption in the Amide I band at 1657 cm⁻¹, Amide II band at 1541 cm⁻¹, the COO⁻ symmetric stretch band at 1391 cm⁻¹, the PO₂ asymmetric stretching band at 1236 cm⁻¹, the PO₂ symmetric stretching; phospholipids C-O stretch band at 1080 cm⁻¹ compared to the *A. citrulli* strains, while the *A. citrulli* strains have a higher intensity of the absorption in the CH₃ asymmetric stretching vibration at 2959 cm⁻¹, the CH₂ asymmetric stretching vibration at 2927 cm⁻¹, the CH₃ symmetric stretching vibration at 2876 cm⁻¹, the CH₂ symmetric stretching vibration at 2857 cm⁻¹, the CH₂ bending band at 1452 cm⁻¹ compared to the *A. oryzae* strains. However, there was not significant difference in the frequency value of the PO₂ asymmetric stretching band at 1236 cm⁻¹ between the two species (Figure 2; Table 3; Additional file 1).

The average spectra in the 2800–1800 cm⁻¹ region were not detailed compared between the two species for no obvious peaks were found in the region (Figure 2; Table 3). Interestingly, this result indicated that five distinctive peaks around at 1738, 1311, 1128, 1078 and 989 cm⁻¹ was observed in the *A. oryzae* strains, but not in the *A. citrulli* strains, while five distinctive peaks centered at 1337, 968, 933, 916 and 786 cm⁻¹ was only observed in the *A. citrulli* strains, but not in the *A. oryzae* strains (Figure 2; Table 3; Additional file 1). These characteristic peaks are specific to either the *A. citrulli* strains or the *A. oryzae* strains. Therefore, it could be suggested that these characteristic peaks may be able to be used for the discrimination of the two species of *Acidovorax*.




Figure 2 The average FTIR spectra in the 4000–500 cm^{-1} region for both *Acidovorax oryzae* (n = 10) and *Acidovorax citrulli* (n = 10).

Previous related reports have revealed that the prominent peak centered at 2959 cm^{-1} is mainly due to lipids, the prominent peak centered at 2927 cm^{-1} is mainly due to lipids and with little contribution from proteins, carbohydrates and nucleic acids, the prominent peak centered at 2876 cm^{-1} is mainly due to proteins, the prominent peak centered at 2857 cm^{-1} is mainly due to

lipids, the band centered at 1739 cm^{-1} is mainly assigned to the C=O ester stretching vibration of triglycerides, the bands centered at 1657 cm^{-1} is mainly assigned to the stretching C=O (amide I) vibrational modes of the polypeptide and protein backbone, the band centered at 1541 cm^{-1} is mainly assigned to the bending N-H and stretching C-N (amide II), the band at 1452 cm^{-1} is

Table 3 The band frequencies and absorption intensity of various functional groups in the *Acidovorax oryzae* (Ao) and *Acidovorax citrulli* (Ac) strains

Functional groups	Frequency (cm^{-1})			Intensity		
	Ao (n = 10)	Ac (n = 10)	P-value	Ao (n = 10)	Ac (n = 10)	P-value
CH ₃ asymmetric stretching	2965.25 ± 0.35	2962.68 ± 0.14	***	1.14 ± 0.02	1.19 ± 0.02	*
CH ₂ asymmetric stretching	2930.14 ± 0.26	2927.85 ± 0.23	***	1.23 ± 0.02	1.25 ± 0.01	*
CH ₃ symmetric stretching	2873.22 ± 0.47	2875.97 ± 0.36	***	0.83 ± 0.01	0.89 ± 0.02	*
CH ₂ symmetric stretching	2853.15 ± 0.36	2855.22 ± 0.56	***	0.74 ± 0.05	0.86 ± 0.07	**
Amide I	1653.85 ± 0.21	1651.61 ± 0.14	***	2.97 ± 0.15	1.84 ± 0.25	***
Amide II	1542.53 ± 0.33	1539.82 ± 0.11	***	1.98 ± 0.25	1.57 ± 0.36	**
CH ₂ bending	1453.61 ± 0.43	1452.14 ± 0.14	**	0.90 ± 0.03	0.96 ± 0.02	*
COO ⁻ symmetric stretch	1394.20 ± 0.36	1397.09 ± 0.25	***	0.98 ± 0.02	0.92 ± 0.05	*
PO ₂ asymmetric stretching	1239.61 ± 0.12	1239.48 ± 0.19	0.12	1.01 ± 0.02	0.91 ± 0.02	*
PO ₂ symmetric stretching	1058.65 ± 1.78	1080.02 ± 0.56	***	1.14 ± 0.19	0.89 ± 0.08	***

Data are the mean of the 10 strains. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

Table 4 The band area values of various functional groups and protein/lipid ratio values in *Acidovorax oryzae* (Ao) and *Acidovorax citrulli* (Ac) strains

Functional groups	Ao (n = 10)	Ac (n = 10)	P-value
Band area value			
CH ₃ asymmetric stretching	0.152 ± 0.002	0.183 ± 0.010	*
CH ₃ symmetric stretching	0.053 ± 0.004	0.036 ± 0.002	*
Amide I	3.603 ± 0.021	1.668 ± 0.036	***
Amide II	1.931 ± 0.012	1.150 ± 0.011	**
PO ₂ asymmetric stretching	0.379 ± 0.062	0.801 ± 0.008	**
PO ₂ symmetric stretching	1.061 ± 0.051	1.182 ± 0.036	**
Protein/lipids ratio			
CH ₃ symmetric/CH ₃ asymmetric	0.349 ± 0.044	0.196 ± 0.015	***
DNA/Protein ratio			
PO ₂ asymmetric/Amide II	0.196 ± 0.006	0.697 ± 0.007	***

Data are the mean of the 10 strains. *: p < 0.05, **: p < 0.01, ***: p < 0.001.

mainly assigned to the CH₂ bending mode of lipids [6-9,12,13,25-29], the band at around 1337 cm⁻¹ was due to acetic acid which was produced by an acetate oxidation [30], the bands at 968, 933 and 916 cm⁻¹ were assigned to the vibration of C-O-C ring deoxyribose, the lipid C=O stretching vibration band at 1738 cm⁻¹ has been suggested as indicative of an increased concentration and difference in packing of the ester groups in bacteria [31]. Furthermore, the band at 1311 cm⁻¹ was due to the stretching mode of C-O of carboxylic acids which suggested an exopolymer formation in bacteria [32], while these bands at 1128, 1078 and 989 cm⁻¹ were due to DNA and RNA backbones, glycogen, and nucleic acids, respectively [6,21]. Therefore, the difference of FTIR spectra between the two species may be due mainly to the imparity of the macromolecular composition and concentration.

This study revealed that the protein-to-lipid ratio was significantly higher for the *A. oryzae* strains than for the *A. citrulli* strains in this study (Figure 2; Table 4), which was calculated by taking the ratio of the area of the CH₃ symmetric stretching band at 2876 cm⁻¹ to the area of the CH₃ asymmetric stretching band at 2959 cm⁻¹ [6,21]. In agreement with the result of the protein-to-lipid ratio, the ratio of DNA-to-protein was higher for the *A. citrulli* strains than for the *A. oryzae* strains (Figure 2; Table 4), which was calculated by taking the ratio of the area of PO₂ symmetric stretching band at 1080 cm⁻¹ to the area of the band at 1541 cm⁻¹ [6,21].

The ratio of protein-to-lipid in the membranes is an important factor affecting the membrane structure and dynamics [33]. Interestingly, the frequency of Amide I and Amide II has been regarded as indicative of conformation and structure of cellular proteins [31,34], while the absorption intensity of Amide I and Amide II has

been regarded as indicative of protein content in bacterial cells [6,21]. However, in this study, the *A. oryzae* strains not only have a higher value in the frequency and the absorption intensity of both Amide I and Amide II, but also in the triglyceride content that is indicative of the lipids compared to the *A. citrulli* strains. Therefore, the major contribution to the higher protein-to-lipid ratio in the *A. oryzae* strains comes from the significant increase of the area of both Amide I and Amide II.

Conclusions

In summary, our results indicated that there were significant differences in MALDI-TOF MS and FTIR spectra between the two species. In particular, several specific characteristic peaks were determined for each of the two species. Compared to the traditional time-consuming method, MALDI-TOF MS and FTIR spectroscopy is easy to implement and is an emergent physico-chemical technique in bacterial research. Therefore, result from this study may give a new strategy for the rapid bacterial identification and differentiation of the two species of *Acidovorax*.

Additional file

Additional file 1: The average FTIR spectra in the 4000–2800 cm⁻¹ (a); 1800–1400 cm⁻¹ (b); 1400–1000 cm⁻¹ (c); 1000–500 cm⁻¹ (d) region for both *Acidovorax oryzae* (n = 10) and *Acidovorax citrulli* (n = 10).

Competing interests

None declared.

Authors' contributions

Wang YL and Li B designed the experiments and wrote the paper. Liu BP, Zhou Q, Wu GX and Ibrahim M performed the experiments. Xie GL, Li HY and Sun GC coordinated the project. All authors have read and approved the manuscript.

Acknowledgements

This work was supported by Zhejiang Provincial Natural Science Foundation of China (Y3090150), Zhejiang Provincial Project (2010R10091), the Fundamental Research Funds for the Central Universities, the Agricultural Ministry of China (nyhyzx 201003029; 201003066), and Key Subject Construction Program of Zhejiang for Modern Agricultural Biotechnology and Crop Disease Control (2010DS700124-KF1101).

Author details

¹State Key Laboratory of Rice Biology, Institute of Biotechnology, Zhejiang University, Hangzhou 310029, China. ²State Key Laboratory Breeding Base for Zhejiang Sustainable Pest and Disease Control, Institute of Plant Protection and Microbiology, Zhejiang Academy of Agricultural Sciences, Hangzhou 310021, China. ³College of Plant Protection, Yunnan Agricultural University, Kunming 650201, China.

Received: 15 May 2012 Accepted: 23 July 2012

Published: 18 August 2012

References

1. Walcott RD, Gitaitis RD: Detection of *Acidovorax avenae* subsp. *citrulli* in watermelon seed using immunomagnetic separation and the polymerase chain reaction. *Plant Dis* 2000, **84**:470–474.

2. Zhao LH, Wang X, Xie GL, Xu FS, Xie GX: **Detection for pathogen of bacterial fruit blotch of watermelon by immuno-capture PCR.** *J Agr Biotechnol* 2006, **14**:946–951.
3. Li B, Liu BP, Yu RR, Tao ZY, Wang YL, Xie GL, Li HY, Sun GC: **Bacterial brown stripe of rice in soil-less culture system caused by *Acidovorax avenae* subsp. *avenae* in China.** *J Gen Plant Pathol* 2011, **77**:64–67.
4. Xie GL, Zhang GQ, Liu H, Lou MM, Tian WX, Li B, Zhou XP, Zhu B, Jin GL: **Genome sequence of the rice pathogenic bacterium *Acidovorax avenae* subsp. *avenae* RS-1.** *J Bacteriol* 2011, **193**:5013–5014.
5. Xu LH, Qiu W, Zhang WY, Li B, Xie GL: **Identification of the causal organism of bacterial brown stripe from rice seedling.** *Chinese J Rice Sci* 2008, **22**:302–306.
6. Garip S, Cetin GA, Severcan F: **Use of Fourier transform infrared spectroscopy for rapid comparative analysis of *Bacillus* and *Micrococcus* isolates.** *Food Chem* 2009, **113**:1301–1307.
7. Samelis J, Bleicher A, Delbes-Paus C, Kakouri A, Neuhaus K, Montel MC: **FTIR-based polyphasic identification of lactic acid bacteria isolated from traditional Greek Graviera cheese.** *Food Microbiol* 2011, **28**:76–83.
8. Wang J, Kim KH, Kim S, Kim YS, Li QX, Jun S: **Simple quantitative analysis of *Escherichia coli* K-12 internalized in baby spinach using Fourier Transform Infrared spectroscopy.** *Int J Food Microbiol* 2010, **144**:147–151.
9. Vodnar DC, Socaciu C, Rotar AM, Stanila A: **Morphology, FTIR fingerprint and survivability of encapsulated lactic bacteria (*Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *bulgaricus*) in simulated gastric juice and intestinal juice.** *Int J Food Sci Tech* 2010, **45**:2345–2351.
10. Lista F, Reubsaet FAG, De Santis R, Parchen RR, de Jong AL, Kieboom J, van der Laaken AL, Voskamp-Visser IAI, Fillo S, Jansen HJ, Van der Plas J, Paauw A: **Reliable identification at the species level of *Brucella* isolates with MALDI-TOF-MS.** *BMC Microbiol* 2011, **11**:267.
11. Ayyadurai S, Flaudrops C, Raoult D, Drancourt M: **Rapid identification and typing of *Yersinia pestis* and other *Yersinia* species by matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry.** *BMC Microbiol* 2010, **10**:285.
12. Toyran N, Lasch P, Naumann D, Turan B, Severcan F: **Early alterations in myocardia and vessels of the diabetic rat heart: An FTIR microspectroscopic study.** *Biochem J* 2006, **397**:427–436.
13. Lau NS, Tsuge T, Sudesh K: **Formation of new polyhydroxyalkanoate containing 3-hydroxy-4-methylvalerate monomer in *Burkholderia* sp.** *Appl Microbiol Biotechnol* 2011, **89**:1599–1609.
14. Krieg NR, Holt JG: *Bergey's Manual of Systematic Bacteriology.* Baltimore: Williams and Wilkins; 1984.
15. Schaad NW, Postnikova E, Sechler A, Clafin LE, Vidaver AK, Jones JB, Agarkova I, Ignatov A, Dickstein E, Ramundo BA: **Reclassification of subspecies of *Acidovorax avenae* as *A. avenae* (Manns 1905) emend., *A. cattleyae* (Pavarino, 1911) comb. nov., *A. citrulli* (Schaad et al., 1978) comb. nov., and proposal of *A. oryzae* sp. nov.** *Syst Appl Microbiol* 2008, **31**:434–446.
16. Li B, Xie GL, Zhang JZ, Janssens D, Swings J: **Identification of the bacterial leaf spot pathogen of poinsettia in China.** *J Phytopathol* 2006, **151**:711–715.
17. Li B, Yu RR, Yu SH, Qiu W, Fang Y, Xie GL: **First report on bacterial heart rot of garlic caused by *Pseudomonas fluorescens* in China.** *Plant Pathol J* 2009, **25**:91–94.
18. Song WY, Kim HM, Hwang CY, Schaad NW: **Detection of *Acidovorax avenae* ssp. *avenae* in rice seeds using BIO-PCR.** *J Phytopathol* 2004, **152**:667–676.
19. Decristophoris P, Fasola A, Benagli C, Tonolla M, Petrini O: **Identification of *Staphylococcus intermedius* group by MALDI-TOF MS.** *Syst Appl Microbiol* 2011, **34**:45–51.
20. Figueras MJ, Levican A, Collado L, Inza MI, Yustes C: ***Arcobacter ellisii* sp. nov., isolated from mussels.** *Syst Appl Microbiol* 2011, **34**:414–418.
21. Garip S, Bozoglu F, Severcan F: **Differentiation of mesophilic and thermophilic bacteria with Fourier transform infrared spectroscopy.** *Appl Spectrosc* 2007, **61**:186–192.
22. Ryzhov V, Fenselau C: **Characterization of the protein subset desorbed by MALDI from whole bacterial cells.** *Anal Chem* 2001, **73**:746–750.
23. Lay J: **MALDI-TOF mass spectrometry of bacteria.** *Mass Spectrom* 2001, **20**:172–194.
24. Moore ERB, Rosselló-Móra R: **MALDI-TOF MS: A return to phenotyping in microbial identification?** *Syst Appl Microbiol* 2011, **34**:1.
25. Savic D, Jokovic N, Topisirovic L: **Multivariate statistical methods for discrimination of lactobacilli based on their FTIR spectra.** *Dairy Sci Tech* 2008, **88**:273–290.
26. Dziuba B, Babuchowski A, Nalecz D, Niklewicz M: **Identification of lactic acid bacteria using FTIR spectroscopy and cluster analysis.** *Int Dairy J* 2007, **17**:183–189.
27. Rebuffo-Scheer CA, Schmitt J, Scherer S: **Differentiation of *Listeria monocytogenes* serovars by using artificial neural network analysis of Fourier-transformed infrared spectra.** *Appl Environ Microbiol* 2007, **73**:1036–1040.
28. Yu C, Irudayaraj J: **Identification of pathogenic bacteria in mixed cultures by FTIR spectroscopy.** *T ASABE* 2006, **49**:1623–1632.
29. Lamprell H, Mazerolles G, Kodjo A, Chamba JF, Noel Y, Beuvier E: **Discrimination of *Staphylococcus aureus* strains from different species of *Staphylococcus* using Fourier transform infrared (FTIR) spectroscopy.** *Int J Food Microbiol* 2006, **108**:125–129.
30. Liao LF, Lien CF, Lin JL: **FTIR study of adsorption and photoreactions of acetic acid on TiO₂.** *Phys Chem Chem Phys* 2001, **3**:831–837.
31. Jackson M, Ramjawan B, Hewko M, Mantsch HH: **Infrared microscopic functional group mapping and spectral clustering analysis of hypercholesterolemic rabbit liver.** *Cell Mol Biol* 1998, **44**:89–98.
32. Nichols PD, Henson JM, Guckert JB, Nivens DE, White DC: **FTIR methods microbial ecology: Analysis of bacteria, bacteria-polymer mixtures and biofilms.** *J Microbiol Meth* 1985, **4**:79–94.
33. Szalontai B, Nishiyama Y, Gombos Z, Murata N: **Membrane dynamics as seen by Fourier transform infrared spectroscopy in a cyanobacterium, *Synechocystis* PCC 6803. The effects of lipid unsaturation and the protein-tolipid ratio.** *Biochim Biophys Acta* 2000, **1509**:409–419.
34. Haris PI, Severcan F: **FTIR spectroscopic characterization of protein structure in aqueous and non-aqueous media.** *J Mol Catal B Enzym* 1999, **7**:207–221.

doi:10.1186/1471-2180-12-182

Cite this article as: Wang et al.: Differentiation in MALDI-TOF MS and FTIR spectra between two closely related species *Acidovorax oryzae* and *Acidovorax citrulli*. *BMC Microbiology* 2012 **12**:182.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

