



Assessment Methods for Problematic Eating Behaviors in Children and Adolescents With Autism Spectrum Disorder

Miji Lee^{1*}, Seolha Lee^{1*}, Jong-Woo Sohn², Ki Woo Kim³, and Hyung Jin Choi^{1,4,5,6}

¹Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul, Korea

²Department of Biological Sciences, Korea Advanced Institute of Science and Technology, Daejeon, Korea

³Division of Physiology, Departments of Oral Biology and Applied Life Science, BK21 Four, Yonsei University College of Dentistry, Seoul, Korea

⁴Department of Anatomy and Cell Biology, Seoul National University College of Medicine, Seoul, Korea

⁵Neuroscience Research Institute, Seoul National University College of Medicine, Seoul, Korea

⁶Wide River Institute of Immunology, Seoul National University, Hongcheon, Korea

J Korean Acad Child Adolesc Psychiatry 2024;35(1):57-65

<https://doi.org/10.5765/jkacap.230065>

In the article above, references 36 and 66 were listed incorrectly. The correct references are:

36) **Matson JL, LoVullo SV, Rivet TT, Boisjoli JA.** Validity of the Autism Spectrum Disorder-Comorbid for Children (ASD-CC). *Res Autism Spectr Disord* 2009;3:345-357.

66) **Lal R.** Effect of alternative and augmentative communication on language and social behavior of children with autism. *Educ Res Rev* 2010;5:119-125.

And in the subsection “Binge-eating (overeating)” on page 59, the seventh sentence should read as follows:

“Moreover, overeating symptoms among individuals with autism can be used in Autism Spectrum Disorders-Comorbid for Children (ASD-CC) [36].”