

SPECIAL POPULATIONS SECTION

Prevalence and Impact of Dysmenorrhea Among University Students in Ireland

Hannah Durand , PhD,^{*,†} Katie Monahan, BA,[†] and Brian E. McGuire, PhD^{*,†}

^{*}Centre for Pain Research, National University of Ireland, Galway, Ireland; and [†]School of Psychology, National University of Ireland, Galway, Ireland

Correspondence to: Hannah Durand, PhD, Centre for Pain Research, School of Psychology, National University of Ireland, Galway, H91EV56, Ireland. Tel: 0035391495831; E-mail: hannah.durand@nuigalway.ie.

Funding sources: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of interest: The authors have no conflicts of interest to declare.

Abstract

Objective. Primary dysmenorrhea (PD), or painful menstruation, is a common gynecological condition that can cause intense pain and functional disability in women of reproductive age. As a nonmalignant condition, PD is relatively understudied and poorly managed. The purpose of this study was to estimate the prevalence and impact of PD among third-level students in Ireland. **Design.** A cross-sectional observational design was used. **Methods.** Students ($n = 892$; age range = 18–45 years) completed an online survey on menstrual pain characteristics, pain management strategies, pain interference, and pain catastrophizing. **Results.** The prevalence of PD was 91.5% (95% confidence interval = 89.67–93.33). Nonpharmacological management strategies were most popular (95.1%); of these, heat application (79%), rest (60.4%), hot shower/bath (40.9%), and exercise (25.7%) were most common. Perceived effectiveness of these methods varied between participants. Analgesic use was also common (79.5%); of these, paracetamol was most used (60.5%) despite limited perceived effectiveness. Pain catastrophizing was a significant predictor of variance in both pain intensity and pain interference scores such that those with higher pain catastrophizing scores reported more intense pain and greater interference with daily activities and academic demands. **Conclusions.** This article presents the first investigation into PD among third-level students in Ireland. Poorly managed menstrual pain may impact functional ability across several domains. Future research should focus on improving menstrual pain management education and support and promoting menstrual health literacy for women affected by PD.

Key Words: Dysmenorrhea; Menstrual Pain; Menstruation; Pain Catastrophizing; Anxiety

Introduction

Primary dysmenorrhea (PD) is defined as painful menstrual cramps of uterine origin in the absence of pelvic pathology [1]. It is often characterized by congestive (deep, dull ache) and/or spasmodic (sharp spasms) pain and associated with a number of additional symptoms including fatigue, headache, backache, moodiness, irritability, constipation, and painful urination [2, 3]. PD is known to negatively affect quality of life and can result in absenteeism from school and work [4]. Despite this, given the nonmalignant nature of the condition, PD is generally understudied and poorly managed.

PD is the most common gynecological condition among women of reproductive age [5]. It affects an estimated 45–95% of menstruating women globally [1]; however, prevalence estimates vary widely, in part due to methodological differences but also cultural differences that may limit the generalizability of international estimates. Clinical risk factors for PD include younger age at menarche, irregular menstruation, and heavy menstrual flow [6, 7]. Women with PD report significantly lower quality of life during their menstruation phase in comparison with their pain-free follicular phases and compared with those without pain during menstruation [1]. PD can also lead to significant interference with daily activities

[8–11]. Importantly, for those in the critical developmental phases of adolescence and emerging adulthood, which typically coincide with secondary school and higher education, PD may negatively impact educational attainment through absenteeism and/or presenteeism, whereby people who menstruate may not be able to perform academically due to interference with concentration and performance during their period [2, 12–14].

Secondary dysmenorrhea, defined as menstrual pain resulting from anatomic or macroscopic pelvic pathology, is typically caused by a gynecologic disorder such as endometriosis, adenomyosis, or fibroids, or by congenital anomalies of the pelvic reproductive organs. Such conditions, particularly endometriosis, are not uncommon in young people [15, 16]; however, diagnoses are often delayed [17]. This may be in part due to low healthcare consultation rates for people with dysmenorrhea [18, 19]. Reasons for not seeking care for dysmenorrhea identified by Chen et al. [20] include assuming symptoms are normal, thinking providers would not offer help, being unaware of treatment options, and feeling embarrassed or afraid to seek care. In addition to the risk of underlying medical causes of menstrual pain going undetected, underutilization of healthcare for dysmenorrhea may contribute to suboptimal self-management through lack of awareness and underuse of effective strategies.

Pain catastrophizing, characterized by a tendency to magnify the threat value and to feel helpless in the context of pain, and by a relative inability to inhibit pain-related thoughts [21], may represent an important risk factor for adverse menstrual pain-related outcomes, such as pain intensity and disability. Females have been found to be more likely to engage in specific problematic coping strategies such as catastrophizing than males [22]. Indeed, pain catastrophizing has been shown to mediate the relationship between sex and pain for adolescents with chronic pain [22]. Relatively few studies [23, 24] have examined relationships between pain catastrophizing and dysmenorrhea to date, and these have typically focused on chronic pain samples. Research into the relationship between pain catastrophizing and dysmenorrhea-related outcomes in community samples is lacking.

The aim of the current study was to estimate the prevalence of dysmenorrhea among university students living in Ireland and to explore the predictive value of pain catastrophizing scores in explaining variance in pain intensity and pain interference.

Methods

Design

This study utilized a cross-sectional (observational) quantitative online survey design.

Participants

Third-level students in Ireland who menstruate were invited to self-select into a study about their menstruation. Inclusion criteria were being a student in a third-level educational institution in Ireland (e.g., university, college, institute of technology), aged 18 years or older, who has reached menarche and is premenopausal (i.e., has not reached perimenopause or menopause). Those on contraception who might use menstrual suppression were not specifically excluded. All who met these criteria were eligible to participate, regardless of how they identified in terms of gender.

Sample Size

A power calculation with finite population correction, assuming 5% precision, a conservative expected proportion of 50% based on the existing international literature, a large population to draw on (i.e., more than 120,000 females of reproductive age enrolled in higher education in Ireland) [25], and a 95% confidence interval, suggested a target sample of $N = 385$ would be sufficient to allow for estimation of the prevalence of PD in the general university population [26].

Measures

A copy of the survey tool is available via the Open Science Framework [27]. The survey was pilot tested with two students who met the eligibility criteria, and changes were made based on their feedback. The feedback related to the length of the questionnaire and the clarity of response options for certain items. Participants were asked to self-report basic demographic information as well as information about their menstrual cycle, menstrual pain experience, strategies they used to manage pain, and their perceived effectiveness of these. Prompts were used to elicit information about pain management strategies as well as any diagnosed gynecological conditions that may have been associated with menstrual pain. In responding to these items, participants selected from a list devised from previous research as many response options as were relevant to them. These lists also included an “other” option in which participants could enter via a free-text box any responses not listed on the survey. Effectiveness of pain management strategies was rated on a 5-point Likert scale ranging from 0 “not at all effective” to 4 “very effective.” To protect participant anonymity, only age ranges were recorded. Participants were also asked to complete the following measures.

Numerical Rating Scale (NRS). Pain intensity was measured using the NRS, a scale consisting of whole number scores between 0 and 10, with higher scores indicating a greater pain intensity. Across many different chronic pain conditions, the NRS has shown to be highly correlated with the visual analog scale [28]. This scale is a reliable and valid subjective measure of menstrual pain intensity [29, 30].

Patient-Reported Outcomes Measurement Information System (PROMIS). Pain interference was assessed using the PROMIS Short Form v1.0—Pain Interference 8a for adults [31]. This scale assessed self-reported impact of pain on aspects of the individual's life (social, cognitive, physical, recreational, and emotional). The prompt given at the start of the scale was modified from “In the past 7 days . . .” to “When you have menstrual pain . . .” to measure PD interference specifically. Participants responded on a 5-point Likert scale from 1 “not at all” to 5 “very much.” The total raw score was calculated by summing the response values to each question, with a possible range between 8 and 40. Additional specific questions relating to academic absenteeism and performance were also included. In the current study, the Cronbach's alpha coefficient for the PROMIS scale was 0.95, suggesting very good internal consistency.

Pain Catastrophizing Scale (PCS). Pain catastrophizing was measured using the PCS [32], a 13-item self-report measure that assesses three dimensions of pain catastrophizing (rumination, magnification, and helplessness). Participants indicated the extent to which they agree with statements on a 5-point Likert scale ranging from 0 “not at all” to 4 “all the time.” The rank score for the statements was summed to find the total catastrophizing score ranging from 0 to 52. The scale has shown good internal consistency ($\alpha=0.87$) [32]. In the current sample, the Cronbach's alpha coefficient was 0.94, suggesting very good internal consistency.

Procedure

Ethical approval was obtained from the School of Psychology Research Ethics Committee at NUI Galway. Potential participants were invited to complete the survey via institutional mailing lists, whereby each institution shared the study information with registered students via email on one occasion, and social media posts on Facebook, Twitter, and Instagram. Due to resource limitations, we could not pay for targeted social media advertising. Posts were made public and could be shared by other social media users. Participants were given information about the purpose and protocol of the study, and informed consent was obtained prior to data collection. Data collection ran from January to March 2020. Participants filled in the online questionnaire through the LimeSurvey platform on their smart phones, laptops, tablets, or personal computers in their own time. The survey took between 5 and 10 minutes to complete. Upon completion, participants were debriefed about the purpose of the study.

Statistical Analyses

Data were analyzed using SPSS v.25.0 [33]. A listwise deletion approach to handling missing data was taken, such that the current analyses include only participants with complete data on the variables of interest. Listwise

deletion was chosen due to the small and random nature of the missing data (<1% missing) and the large sample size [34]. Descriptive statistics were computed for demographics, pain, and menstruation characteristics listed above. Correlation coefficients (i.e., Pearson correlation coefficients [r] for two continuous variables and point-biserial correlation coefficients [r_{pb}] for one continuous and one binary variable) and hierarchical multiple linear regression were used to examine associations between predictor variables (i.e., clinical characteristics such as age at menarche, having an irregular menstrual cycle, having a heavy flow, and pain catastrophizing) and criterion variables (i.e., pain intensity and pain interference).

Results

Results are reported in accordance with the STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) statement [35]. A STROBE checklist is provided in [Supplementary Data, Appendix A](#).

Sample Characteristics

A sample of 899 university students in Ireland who menstruate initiated the online survey. Of these, 7 did not complete the survey, leaving a sample of 892 participants for analysis following listwise deletion. The majority (85.1%) were aged between 18 and 24 years, and most (81.5%) were undergraduate students. Demographic characteristics are presented in [Table 1](#).

The mean age of menarche was 12.54 years (range = 5–17¹). For most of the sample (87.6%), menstruation was regular (i.e., period occurs approximately every 28 days, between 21 and 35 days). The median menstrual frequency of those with irregular periods was 40 days (IQR = 24; total range = 6–180). Additional detail on menstrual patterns of the sample is provided in [Table 2](#). Beyond those listed on the survey, other related symptoms reported in a free-text box include nausea (1.6%), skin blemishes, pins and needles, mouth ulcers, heavy dreams, flatulence, ringing of the ears, and dermatographia (all <1%).

Prevalence and Characteristics of Menstrual Pain

Menstrual pain occurred in 96.9% of respondents. Of these, 5.4% were diagnosed with a gynecological

1 One participant reported the age at which they had their first period as 5 years old. While this may have been erroneous, it may be the case that this participant experienced precocious puberty (i.e., onset of puberty before the age of 8 years in girls). We therefore have not excluded this participant from these analyses. For information, the mean age of menarche in the sample with this participant excluded from the calculation is 12.55 years (1.40), the median is 12 (interquartile range = 1), and the range of responses is 8–17 years.

Table 1. Participant demographics

Demographics	No.	%
Age groups, y		
18–24	759	85.1
25–34	109	12.2
35–44	22	2.5
45–54	2	0.2
Year of study		
1	282	31.6
2	168	18.9
3	197	22.1
4	79	8.9
Postgraduate	165	18.5

Table 2. Cycle and menstruation characteristics

Characteristics	No.	%
Period duration, days		
1–2	12	1.5
3–4	269	33
5–6	443	54.3
7 or more	92	11.3
Menstrual flow		
Light	62	7.6
Moderate	492	60.3
Heavy	262	32.1
Experienced related symptom(s)		
Yes	811	99.4
No	5	0.6
Related symptoms		
Mood change	692	85.3
Fatigue	655	80.8
Bloating	632	77.9
Tender breasts	491	60.5
Appetite change	440	54.3
Headache	352	43.4
Diarrhea	351	43.3
Sweating	262	32.3
Dizziness	215	26.5
Constipation	211	26
Vomiting	95	11.7
Fainting	68	8.4
Other	21	2.6

Participants could choose more than one option; therefore, columns may add up to greater than 100%.

condition (i.e., may be more appropriately classed as having secondary dysmenorrhea): 2.7% had a diagnosis of endometriosis; 1.7% had polycystic ovary syndrome; and less than 1% reported another diagnosis (uterine fibroids, adenomyosis, or pelvic inflammatory disease). Therefore, the prevalence of PD in this sample was 91.5% (95% confidence interval = 89.67–93.33). Of these, most respondents experienced pain with every period (52.2%) or most periods (27.3%). The mean menstrual pain rating was 5.45 on average (1.79, median = 6, interquartile range [IQR] = 3, range = 1–10) and 7.41 at its most severe (1.73, median = 8, IQR = 3, range = 1–10). Only 37.2% had ever visited a healthcare professional regarding menstrual pain; those who did reported higher pain intensity on average ($t = -9.61$; $P < 0.001$) and at its

most severe ($t = -11.10$; $P < 0.001$). Beyond those listed on the survey (Table 3), other pain areas identified by participants included the lower legs (1.8%), the upper back, vagina, anus, shoulders, neck, chest, pelvis, and the cervix (all <1%). A full breakdown of dysmenorrhea-associated features is presented in Table 3.

Pain Management

Table 4 depicts the pain management strategies used as well as their perceived effectiveness. Nonpharmacological pain management strategies were most popular, with 95.1% of respondents reporting at least one such method; of these, direct heat application, having a hot shower or bath, rest, and exercise were most common. Though much less common, exercise was rated as more effective for managing pain than was rest. Other nonpharmacological methods identified by participants included chocolate, lying in the fetal position, masturbation, sex, rocking back and forth, sleep, teas, transcutaneous electrical nerve stimulator machine, prescription blood thinners, and topical essential oils (all <1%). Analgesic use was also common (79.5%); paracetamol was most used (60.5%). Nonsteroidal anti-inflammatory drugs (e.g., ibuprofen, aspirin), which are the first-line treatment for menstrual pain, were less commonly used. Other drugs identified by participants include butylscopolamine, diclofenac, and mefenamic acid (all <1%). The mean number of pain management strategies used was 4.37 (2.12, range = 1–14). The number of strategies used was significantly correlated with pain intensity, such that participants who reported more intense pain used more methods ($r = 0.24$; $P < 0.001$).

Pain Interference

Responses to the PROMIS measure of pain interference showed considerable impact of PD on social, cognitive, physical, recreational, and emotional aspects of life. Participants reported that pain interfered “quite a bit” or “very much” with day-to-day activities (36.6%); work around the home (34.1%) and household chores (30.5%); ability to participate in (42.4%) and enjoy social activities (47.2%); things usually done for fun (39.7%) and enjoyment of life (37.1%); and family life (19.6%).

Table 5 presents PD interference with academic demands. For those who reported they had missed lectures due to PD ($n = 513$), the median number of lecture hours missed during the previous menstruation was 3 (IQR = 2; total range = 1–48).

Inferential Analyses

Table 6 presents means, standard deviations, and correlation coefficients between predictor and criterion variables. Analyses of predictors of having PD were not undertaken given the degree of small sample bias (i.e., the small number of participants who did not report PD)

Table 3. Menstrual pain characteristics

Characteristics	No.	%
Menstrual pain		
Yes (PD)	816	91.5
Yes (SD)	48	5.4
No pain	28	3.1
Frequency of pain		
Every period	426	52.2
Most periods	223	27.3
Some periods	167	20.5
Stage at which pain occurs		
Premenstrual period	300	37.5
Beginning of period	699	87.5
Middle of period	191	23.9
End of period	36	4.5
Pain duration		
Less than 1 day	61	7.6
1 day	143	17.9
2 days	361	45.2
3 days	161	20.2
4 or more days	73	9.1
Pain site		
Abdominal region	733	91.7
Lumbar	428	53.6
Groin	228	28.5
Thigh(s)	85	10.6
Other	19	2.4
Consulted doctor for menstrual pain		
Yes	297	37.2
No	502	62.8

Participants could choose more than one option; therefore, columns may add up to greater than 100%. PD = primary dysmenorrhea; SD = secondary dysmenorrhea.

in this case [36]. No strong relationships were observed between predictor variables ($r/r_{pb} < 0.7$). Variance inflation factor (< 2) and tolerance values (> 0.1) for all predictor values were also adequate, thereby demonstrating that there was no issue with multicollinearity in the data. All factors were significantly correlated with pain intensity and pain interference bar having an irregular menstrual cycle, which was not included in any further analyses. All assumptions of hierarchical linear regression were met.

The first hierarchical multiple linear regression explored the impact of predictor variables on average pain intensity. Clinical variables (i.e., age at menarche and having a heavy menstrual flow) were entered into the first block, followed by pain catastrophizing in the second block. The overall model significantly predicted pain intensity ($F_{(3, 782)} = 81.50$; $P < 0.001$), accounting for 23.5% of the variance in pain intensity scores. In block one, having a heavy flow was associated with greater pain intensity. In block two, menstrual pain catastrophizing contributed significantly to the model, explaining an additional 15.3% in the variance explained.

Similar results were observed for the second regression analysis, which explored the impact of predictor variables on pain intensity at its most severe. The overall model significantly predicted pain intensity ($F_{(3, 782)} =$

71.72; $P < 0.001$), accounting for 21.3% of the variance in pain intensity scores. In block one, younger age at menarche and having a heavy flow were associated with greater pain intensity. In the block two, menstrual pain catastrophizing contributed significantly to the model and reduced the contribution of age at menarche to non-significance. Pain catastrophizing made a significant contribution to the model and accounted for an additional 15.4% in the variance explained.

The above procedure was followed for criterion variable pain interference, with the addition of average pain intensity as a predictor variable in block one. The overall model significantly predicted pain interference ($F_{(4, 781)} = 187.01$; $P < 0.001$), accounting for 48.7% of the variance in PROMIS scores. In block one, having a heavier flow and reporting greater pain intensity was associated with greater interference. In block two, menstrual pain catastrophizing contributed significantly to the model, explaining an additional 15.8% of variance beyond clinical characteristics in block one. Results of all hierarchical regression analyses are displayed in Table 7.

Discussion

The current study indicates that PD is highly prevalent among third-level students in Ireland. Our data show that 91% of students in Ireland who menstruate experience PD, with 52% having pain during every period. The majority had experienced pain since their first period. A wide variety of pharmacological and nonpharmacological pain management strategies were used, most commonly heat application, rest, paracetamol, and ibuprofen; the perceived effectiveness of these varied considerably, with many participants relying on strategies that have been shown to be less effective than others that are equally accessible in Ireland. Pain interference was common in this study; concentration during classes and ability to study and to complete assignments were most affected. Participants also reported that PD affected their enjoyment of life and impacted their ability to perform day-to-day activities. Given that the goal of pain management is not just to alleviate pain but also to maximize quality of life and functional ability, we contend that menstrual pain was poorly managed in this sample. Symptom burden was high, with a majority of those with PD experiencing changes in mood, fatigue, bloating, tender breasts, and changes in appetite. Given that more than 80% experienced changes in mood, it follows that PD may have a greater negative effect on mental health and well-being than has been previously understood.

This study is one of few to explore the role of pain catastrophizing in PD to date. We found that pain catastrophizing predicted both pain intensity and pain interference in this sample, after controlling for clinical variables known to predict menstrual pain (e.g., heavy flow). This is consistent with Payne et al. [23], in which pain catastrophizing scores correlated with participants'

Table 4. Pain management strategies used by participants with PD (n = 816)

Type	Prevalence of Use, No. (%)	Perceived Effectiveness, No. (%)				
		Very Effective	Somewhat Effective	A Little Effective	Not Very Effective	Not at All Effective
Analgesics	649 (79.5)					
Ibuprofen	410 (50.2)	93 (22)	230 (56.1)	63 (15.4)	18 (4.4)	6 (1.5)
Aspirin	39 (4.8)	4 (10.5)	18 (47.4)	6 (15.8)	9 (23.7)	1 (2.6)
Paracetamol	494 (60.5)	67 (13.6)	242 (49)	117 (23.7)	51 (10.3)	17 (3.4)
Codeine	167 (20.5)	75 (44.9)	81 (48.5)	10 (6)	1 (0.6)	
Prescription Analgesic	77 (9.4)	30 (39)	33 (42.9)	8 (10.4)	5 (6.5)	1 (1.3)
Other	8 (1)	5 (62.5)	3 (37.5)			
Contraceptives	255 (31.3)					
OCP, patch, ring	228 (27.9)	78 (34.2)	101 (44.3)	26 (11.4)	18 (7.9)	5 (2.2)
LARC	29 (3.6)	8 (27.6)	12 (41.4)	6 (20.7)	1 (3.4)	2 (6.9)
Nonpharmacological	776 (95.1)					
Heat application	645 (79.0)	182 (28.2)	293 (45.4)	145 (22.5)	25 (3.9)	
Hot shower/bath	334 (40.9)	47 (14.1)	164 (49.1)	94 (28.1)	29 (8.7)	
Cold shower/bath	12 (1.5)	1 (8.3)	4 (33.3)	4 (33.3)	3 (25)	
Exercise	210 (25.7)	48 (22.9)	106 (50.5)	50 (23.8)	6 (2.9)	
Yoga	39 (4.8)	9 (23.1)	15 (38.5)	11 (28.2)	4 (10.3)	
Rest	493 (60.4)	96 (19.5)	200 (40.6)	133 (27)	62 (12.6)	2 (0.4)
Relaxation techniques	53 (6.5)	6 (11.3)	24 (45.3)	19 (35.8)	3 (5.7)	1 (1.9)
Meditation	26 (3.2)	2 (7.7)	14 (53.8)	9 (34.6)	1 (3.8)	
Massage	94 (11.5)	9 (9.6)	41 (43.6)	35 (37.2)	9 (9.6)	
Acupuncture	9 (1.1)	3 (33.3)	4 (44.4)	1 (11.1)	1 (11.1)	
Diet change	49 (6)	8 (16.3)	20 (40.8)	13 (26.5)	8 (16.3)	
Dietary supplements	45 (5.5)	3 (6.7)	15 (33.3)	14 (31.1)	12 (26.7)	1 (2.2)
Homeopathy	7 (0.9)		3 (42.9)	3 (42.9)	1 (14.3)	
Herbs	36 (4.4)	4 (11.1)	18 (50)	9 (25)	4 (11.1)	1 (2.8)
Cannabis	23 (2.8)	9 (39.1)	14 (60.9)			
CBD products	12 (1.5)	3 (25)	6 (50)	2 (16.7)	1 (8.3)	
Alcohol	16 (2)	4 (25)	5 (31.3)	4 (25)	3 (18.8)	
Other	20 (2.5)	3 (15)	13 (65)	2 (10)	2 (10)	

CBD = cannabidiol; LARC = long-acting reversible; OCP = oral contraceptive pill; PD = primary dysmenorrhea contraception.

Table 5. Menstrual pain interference with academic demands

Academic Demand	Level of Interference				
	Very Much No. (%)	Quite a Bit No. (%)	Somewhat No. (%)	A Little Bit No. (%)	Not at All No. (%)
Lecture attendance	62 (8.2)	167 (22.1)	184 (24.3)	176 (23.3)	167 (22.1)
Exam study	90 (11.9)	220 (29.1)	195 (25.8)	179 (23.7)	72 (9.5)
Assignment completion	46 (6.1)	158 (20.9)	200 (26.5)	206 (27.2)	146 (19.3)
Concentration	146 (19.3)	237 (31.3)	175 (23.1)	157 (20.8)	41 (5.4)

menstrual pain ratings; however, this relationship was not significant for those without chronic pain. Walsh, LeBlanc, and McGrath [24] also found that women with higher pain catastrophizing scores reported greater disability. Similarly, Kapadi and Elander [37] found that lower physical quality of life was related to higher pain severity and catastrophizing scores. These findings indicate there may be a role for psychological intervention to improve outcomes for people with PD. Cognitive-behavioral therapy [38] or mindfulness-based [39] interventions targeting pain catastrophizing may serve to reduce pain intensity and inference for people with PD.

Psychological intervention may also help promote use of adaptive cognitive and behavioral coping strategies in place of maladaptive strategies such as catastrophic thinking. With research on pain catastrophizing and PD in its relative infancy, the current findings provide an important foundation of evidence on which to base future research and practice.

Clinical and Policy Implications

The current findings have implications for the assessment and clinical management of PD. Participants in this study reported menstrual pain at a variety of sites, including

Table 6. Descriptive statistics and correlations between predictor and criterion variables

Variable	M (SD)	No. (%)	1	2	3	4	5	6	7
1 Age at menarche	12.54 (1.42)		-						
2 Irregular menstruation		115 (12.9)	-0.06	-					
3 Heavy flow		295 (33.1)	-0.07*	-0.07*	-				
4 Pain catastrophizing	16.78 (11.72)		-0.14***	-0.08*	0.20***	-			
5 Pain intensity (on average)	5.51 (1.80)		-0.11**	0.03	0.28***	0.45***	-		
6 Pain intensity (most severe)	7.47 (1.73)		-0.14***	0.01	0.22***	0.44***	0.75***	-	
7 Pain interference	24.93 (8.01)		-0.11**	-0.05	0.27***	0.62***	0.57***	0.54***	-

Variables 2 and 3 were coded as binary variables, whereby 0 = regular menstruation, 1 = irregular menstruation and 0 = light/normal flow, 1 = heavy flow, - = to signify the negative correlation between two variables, respectively.

* $P < 0.05$.

** $P < 0.01$.

*** $P < 0.001$.

Table 7. Hierarchical regression analyses of the contribution of pain catastrophizing to pain intensity and interference

Variable	Adj. R^2	Adj. ΔR^2	ΔF	β
Pain intensity (on average)				
Block 1	0.08	0.08	35.88***	
Age at menarche				-0.04
Heavy flow				0.19***
Block 2	0.24	0.15	158.32***	
Pain catastrophizing				0.40***
Pain intensity (most severe)				
Pain interference				
Block 1	0.06	0.06	25.50***	
Age at menarche				-0.07*
Heavy flow				0.13***
Block 2	0.21	0.15	154.17***	
Pain catastrophizing				0.41***
Pain interference				
Pain intensity (most severe)				
Block 1	0.33	0.33	129.53***	
Age at menarche				0.00
Heavy flow				0.08**
Pain intensity				0.34***
Block 2	0.49	0.16	240.46***	
Pain catastrophizing				0.45***

* $P < 0.05$.

** $P < 0.01$.

*** $P < 0.001$.

the abdomen, lumbar, groin, and thighs. In addition, several gastrointestinal symptoms, such as nausea, vomiting, diarrhea, and bloating as well as headaches and tender breasts, were also reported. This is consistent with previous research, which has shown that increased inflammatory prostaglandins and pain sensitization are likely to contribute to pain at multiple sites as well as gastrointestinal symptoms among women with dysmenorrhea [4, 40–43]. However, this remains at odds with typical clinical practice, whereby abdominal pain intensity is often the only symptom assessed in dysmenorrhea cases [44]. Future dysmenorrhea assessment should involve comprehensive evaluation of pain at different locations and gastrointestinal symptoms as well as psychological factors such as pain catastrophizing and changes in mood to ensure adequate clinical care and facilitate effective self-management.

Consultation rates were relatively high in the current sample but still suboptimal given the symptom burden reported. This is consistent with previous research [2, 45–47]. Low consultation rates may also explain the low prevalence of secondary dysmenorrhea in this sample (5.4%); it is likely that some respondents classed in this study as having PD may have had undiagnosed gynecological conditions such as endometriosis or uterine fibroids. This may also partly explain why some participants reported unusually short durations between periods [48]. Low healthcare consultation rates may be in part due to enduring stigma surrounding menstruation [49] and a lack of education resulting in poor menstrual health literacy and reduced help seeking [46]. It may also be due to the enduring and harmful idea that menstrual pain is something that must be tolerated as “part of being a woman” [20]. Consequently, many people who

menstruate choose menstrual pain management strategies without consulting a medical professional, which may result in unnecessary pain and suffering. This is evident from the data, whereby many participants relied on less effective pain management strategies over those supported by evidence. For example, paracetamol was the most commonly used analgesic despite limited scientific evidence of clinical effectiveness for menstrual pain; non-steroidal anti-inflammatory drugs, the first-line treatment for menstrual pain, were less commonly used despite similar cost and ready availability in Ireland. This is in line with previous research [46], including a recent meta-analysis [50], which reported paracetamol to be the most widely used analgesic despite more than 50% of respondents reporting that it was not always effective for managing their pain. Varying perceived effectiveness ratings in this sample may indicate that what works for one individual may not work for another, particularly regarding nonpharmacological strategies. However, this may reflect that some young people who menstruate lack the necessary knowledge to allow them to self-manage menstrual pain effectively and may endure severe and debilitating symptoms without seeking medical attention and pain management support. Health education measures are needed to promote menstrual health literacy among young people and reduce stigma around menstruation. Comprehensive school- and community-based menstrual health education initiatives that are evidence-based, accessible, and inclusive can serve to increase health literacy, which in turn has positive impacts on understanding, agency and self-management, and appropriate help-seeking behavior [46, 51, 52]. To reduce stigma, educational initiatives should aim to normalize discussion of menstrual health issues by including boys and men in the conversation [53]. Providing this education to children at a younger age (i.e., primary school age) may also promote healthy physical and emotional development into adolescence and adulthood [53].

Limitations and Recommendations for Future Research

The current study used a large sample of third-level students who menstruate. Recruiting participants from third-level institutions across Ireland resulted in a sample that may be more representative than in previous research focusing on single institutions. There may, however, be certain biases in the recruited sample. Due to resource limitations in terms of both time and funding, this study used a nonprobability sampling approach, whereby students were invited to self-select into the study. Nonrepresentative sampling limits the validity of the conclusions we can draw regarding prevalence of PD based on this sample. People who experience menstrual pain may be more inclined to opt into such a study in comparison to those without. This could have contributed to the high prevalence of PD

observed. Furthermore, students may not be representative of young adults in the community [54] as well as adolescent schoolchildren who also experience PD. PD may affect these groups in different ways, for example, in terms of their development or the economic impact of missing work. Socioeconomic and occupational factors may also play a role in menstrual pain experiences. High levels of job strain, exhaustion, and stress related to working conditions have been associated with gynecological pain [55–57]. Additionally, access to gynecological healthcare, analgesic medications, and certain self-management supports may be more difficult for people with fewer economic resources. Finally, limitations of online surveys as regards sampling and access issues have been well documented [58–60] and must be considered when interpreting the current findings. Given advertisements were shared by social media users, it is not known how many eligible students saw the study invitation, and therefore it is not possible to calculate response rates. That said, given the reduced response time, relatively low cost, flexibility, acceptability, and constant advances in survey technology and household connectivity to the internet [61], online surveys such as this can provide useful information on research topics that have been historically deprioritized and underfunded. Future research should use probability sampling and a variety of data collection methods to obtain diverse clinical and community-based samples of people who menstruate, as well as school-based samples of menstruating adolescents, to validate the current prevalence estimate.

There are certain measurement limitations to consider. First, it is not possible to make a diagnosis of secondary dysmenorrhea without a gynecological exam. Reliance on self-report of gynecological health in a sample wherein consultation rates are low is therefore likely to underestimate the prevalence of secondary dysmenorrhea relative to estimates from clinical research. Second, we were limited in our ability to assess academic impact. For example, students may attend lectures but find their concentration in class negatively affected by pain. Previous research has shown that students are less likely to miss lectures because of pain when the lectures are mandatory [11]. Also, self-reported data on missed lectures may be subject to recall bias. Future research should use objective measures of lecture attendance in conjunction with other relevant indicators of academic performance that may be impacted by pain. Third, participants were not asked to report what phase of menstruation they were in at the time of participation. This may have affected pain intensity scores through both response and recall biases, given the cyclical nature of PD. Pain catastrophizing has also been demonstrated to fluctuate during the menstrual cycle [62], which is consistent with the growing literature on situational versus dispositional catastrophizing [63, 64]. Future cross-sectional

research should assess participants' menstrual phase at the time of responding and indeed aim to target phases during which pain is more likely (i.e., menstruation and ovulation phases). Longitudinal research may also provide important insight into the pain experience and its management. Fourth, the PROMIS pain interference measure has not been validated in dysmenorrhea, and so results should be interpreted with caution. Fifth, in an effort to keep survey completion time to a minimum to achieve a sufficiently large sample, several factors thought to affect menstrual pain experiences were not assessed in this study; notably, modifiable and behavioral factors associated with increased risk of PD such as body mass index and smoking were not included. Importantly, there are likely other psychological factors beyond pain catastrophizing that play a role in menstrual pain experiences (e.g., pain acceptance) [37]; in particular, only 23.5% and 21.3% of variance in pain intensity was predicted by our model versus 48.7% variance in pain interference. Future research should explore the predictive value of pain catastrophizing relative to other theoretically and empirically informed predictors. Finally, this study did not address other factors that may be associated with the effectiveness of pain management strategies, particularly dosage of analgesic drugs and timing of administration. Future research should attend to these variables to advance the science and practice of menstrual pain management.

Conclusions

Limitations notwithstanding, the current study provides important data on the prevalence and impact of PD among third-level students in Ireland. Based on these findings, we conclude that menstrual pain is highly prevalent and poorly managed. This has the potential to impact women's functional ability across personal and academic domains. The relationship between pain catastrophizing and pain intensity and interference warrants further investigation, given its implications for pain management and risk of pain-related disability. Further clinical research and health education measures are needed to promote menstrual health literacy and reduce stigma around menstruation to ensure adequate menstrual pain assessment and management.

Acknowledgements

The authors would like to acknowledge and thank the participants who took part in the study.

Supplementary Data

Supplementary data are available at *Pain Medicine* online.

References

- Iacovides S, Avidon I, Baker FC. What we know about primary dysmenorrhea today: A critical review. *Hum Reprod Update* 2015;21(6):762–78.
- Söderman L, Edlund M, Marions L. Prevalence and impact of dysmenorrhea in Swedish adolescents. *Acta Obstet Gynecol Scand* 2019;98(2):215–21.
- Moronkola OA, Uzuegbu VU. Menstruation: Symptoms, management and attitude of female nursing students in Ibadan, Nigeria. *African J Reprod Health* 2006;10(3):84–9.
- Dawood MY. Primary dysmenorrhea: Advances in pathogenesis and management. *Obstet Gynecol* 2006;108(2):428–41.
- Coco AS. Primary dysmenorrhea. *Am Family Phys* 1999;60(2):489–96.
- Hu Z, Tang L, Chen L, Kaminga AC, Xu H. Prevalence and risk factors associated with primary dysmenorrhea among Chinese female university students: A cross-sectional study. *J Pediatr Adolesc Gynecol* 2020;33(1):15–22.
- Azagew AW, Kassie DG, Walle TA. Prevalence of primary dysmenorrhea, its intensity, impact and associated factors among female students' at Gondar town preparatory school, Northwest Ethiopia. *BMC Women's Health* 2020;20(1):1–7.
- Fernández-Martínez E, Onieva-Zafra MD, Parra-Fernández ML. The impact of dysmenorrhea on quality of life among Spanish female university students. *Int J Environ Res Public Health* 2019;16(5):713.
- Ghaderi F, Jafarabadi MA, Bandpei MAM. Dysmenorrhea and self-care strategies in Iranian female students: A regression modeling of pain severity and underlying factors. *Int J Adolesc Med Health* 2017;29(6):20160017.
- Parker MA, Sneddon AE, Arbon P. The menstrual disorder of teenagers (MDOT) study: Determining typical menstrual patterns and menstrual disturbance in a large population-based study of Australian teenagers. *BJOG* 2010;117(2):185–92.
- Vlachou E, Owens DA, Lavdaniti M, et al. Prevalence, well-being, and symptoms of dysmenorrhea among university nursing students in Greece. *Diseases* 2019;7(1):5.
- Armour M, Parry K, Manohar N, et al. The prevalence and academic impact of dysmenorrhea in 21,573 young women: A systematic review and meta-analysis. *J Women's Health* 2019;28(8):1161–71.
- Armour M, Ferfolja T, Curry C, et al. The prevalence and educational impact of pelvic and menstrual pain in Australia: A national online survey of 4202 young women aged 13–25 years. *J Pediatr Adolesc Gynecol* 2020;33(5):511–8.
- Hailemeskel S, Demissie A, Assefa N. Primary dysmenorrhea magnitude, associated risk factors, and its effect on academic performance: Evidence from female university students in Ethiopia. *Int J Women's Health* 2016;8(1):489–96.
- Smorgick N, As-Sanie S, Marsh CA, Smith YR, Quint EH. Advanced stage endometriosis in adolescents and young women. *J Pediatr Adolesc Gynecol* 2014;27(6):320–3.
- Janssen EB, Rijkers ACM, Hoppenbrouwers K, Meuleman C, d'Hooghe TM. Prevalence of endometriosis diagnosed by laparoscopy in adolescents with dysmenorrhea or chronic pelvic pain: A systematic review. *Hum Reprod Update* 2013;19(5):570–82.

17. Greene R, Stratton P, Cleary SD, Ballweg ML, Sinaii N. Diagnostic experience among 4,334 women reporting surgically diagnosed endometriosis. *Fertil Steril* 2009;91(1):32–9.
18. Chen CX, Kwekkeboom KL, Ward SE. Beliefs about dysmenorrhea and their relationship to self-management. *Res Nurs Health* 2016;39(4):263–76.
19. Tanaka E, Momoeda M, Osuga Y, et al. Burden of menstrual symptoms in Japanese women—an analysis of medical care-seeking behavior from a survey-based study. *Int J Women's Health* 2014;6(11).
20. Chen CX, Shieh C, Draucker CB, Carpenter JS. Reasons women do not seek health care for dysmenorrhea. *J Clin Nurs* 2018;27(1-2):e301–8.
21. Quartana PJ, Campbell CM, Edwards RR. Pain catastrophizing: A critical review. *Expert Rev Neurother* 2009;9(5):745–58.
22. Keogh E, Eccleston C. Sex differences in adolescent chronic pain and pain-related coping. *Pain* 2006;123(3):275–84.
23. Payne LA, Rapkin AJ, Lung KC, Seidman LC, Zeltzer LK, Tsao JC. Pain catastrophizing predicts menstrual pain ratings in adolescent girls with chronic pain. *Pain Med* 2016;17(1):16–24.
24. Walsh TM, LeBlanc L, McGrath PJ. Menstrual pain intensity, coping, and disability: The role of pain catastrophizing. *Pain Med* 2003;4(4):352–61.
25. Higher Education Authority. 2018/19 Student Demographics, All HEA-Funded HEIs. Dublin, Ireland: HEA Statistics Archive; 2019.
26. Naing L, Winn T, Rusli B. Practical issues in calculating the sample size for prevalence studies. *Arch Orofacial Sci* 2006;1(1):9–14.
27. Durand H. Prevalence and impact of dysmenorrhea among University students in Ireland. *Open Science Framework*; 2021.
28. Hawker GA, Mian S, Kendzerska T, et al. Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF). *Arthritis Care Res* 2011;63(suppl 11):S240–52.
29. Ameade EPK, Mohammed BS. Menstrual pain assessment: Comparing verbal rating scale (VRS) with numerical rating scales (NRS) as pain measurement tools. *Int J Women's Health Wellness* 2016;2(1):17.
30. Bourdel N, Alves J, Pickering G, Ramilo I, Roman H, Canis M. Systematic review of endometriosis pain assessment: How to choose a scale? *Hum Reprod Update* 2015;21(1):136–52.
31. Amtmann D, Cook KF, Jensen MP, et al. Development of a PROMIS item bank to measure pain interference. *Pain* 2010;150(1):173–82.
32. Sullivan MJL, Bishop SR, Pivik J. The pain catastrophizing scale: Development and validation. *Psychol Assess* 1995;7(4):524–32.
33. IBM Corp. IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY: IBM Corp; 2017.
34. Schafer JL. *Analysis of Incomplete Multivariate Data*. London: Chapman and Hall; 1997.
35. Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Ann Intern Med* 2007;147(8):573–7.
36. King G, Zeng L. Logistic regression in rare events data. *Polit Anal* 2001;9(2):137–63.
37. Kapadi R, Elander J. Pain coping, pain acceptance and analgesic use as predictors of health-related quality of life among women with primary dysmenorrhea. *Eur J Obstet Gynecol Reprod Biol* 2020;246(1):40–4.
38. Lazaridou A, Kim J, Cahalan CM, Loggia ML, et al. Effects of cognitive-behavioral therapy (CBT) on brain connectivity supporting catastrophizing in fibromyalgia. *Clin J Pain* 2017;33(3):215–21.
39. Lazaridou A, Franceschelli O, Protsenko K, Napadow V, Edwards R. (318) The association between mindfulness, catastrophizing and pain interference among patients with fibromyalgia: The moderating role of mindfulness. *J Pain* 2017;18(4):S54–5.
40. Chen CX, Kwekkeboom KL, Ward SE. Self-report pain and symptom measures for primary dysmenorrhoea: A critical review. *Eur J Pain* 2015;19(3):377–91.
41. Jarrett M, Cain KC, Heitkemper M, Levy RL. Relationship between gastrointestinal and dysmenorrheic symptoms at menses. *Res Nurs Health* 1996;19(1):45–51.
42. Mannix LK. Menstrual-related pain conditions: Dysmenorrhea and migraine. *J Women's Health* 2008;17(5):879–91.
43. Nguyen AM, Humphrey L, Kitchen H, Rehman T, Norquist JM. A qualitative study to develop a patient-reported outcome for dysmenorrhea. *Qual Life Res* 2015;24(1):181–91.
44. Chen CX, Draucker CB, Carpenter JS. What women say about their dysmenorrhea: A qualitative thematic analysis. *BMC Women's Health* 2018;18(1):47.
45. Acheampong K, Baffour-Awuah D, Ganu D, et al. Prevalence and predictors of dysmenorrhea, its effect, and coping mechanisms among adolescents in Shai Osudoku District, Ghana. *Obstet Gynecol Int* 2019;2019:1–7.
46. Armour M, Hyman MS, Al-Dabbas M, et al. Menstrual health literacy and management strategies in young women in Australia: A national online survey of young women aged 13–25. *J Pediatr Adolesc Gynecol* 2021;34(2):135–43.
47. Ortiz MI, Rangel-Flores E, Carrillo-Alarcón LC, Veras-Godoy HA. Prevalence and impact of primary dysmenorrhea among Mexican high school students. *Int J Gynecol Obstet* 2009;107(3):240–3.
48. Shafir AL, Farland LV, Shah DK, et al. Risk for and consequences of endometriosis: A critical epidemiologic review. *Best Pract Res Clin Obstet Gynaecol* 2018;51(1):1–15.
49. Johnston-Robledo I, Chrisler JC. The menstrual mark: Menstruation as social stigma. *Sex Roles* 2013;68(1–2):9–18.
50. Armour M, Parry K, Al-Dabbas MA, et al. Self-care strategies and sources of knowledge on menstruation in 12,526 young women with dysmenorrhea: A systematic review and meta-analysis. *PLoS One* 2019;14(7):e0220103.
51. Bush D, Brick E, East MC, Johnson N. Endometriosis education in schools: A New Zealand model examining the impact of an education program in schools on early recognition of symptoms suggesting endometriosis. *Aust N Z J Obstet Gynaecol* 2017;57(4):452–7.
52. Wong CL, Ip WY, Choi KC, Lam LW. Examining self-care behaviors and their associated factors among adolescent girls with dysmenorrhea: An application of Orem's self-care deficit nursing theory. *J Nurs Scholarsh* 2015;47(3):219–27.
53. Fahs B, Perianes MB. Transnational engagement: Designing an ideal menstrual health (MH) curriculum—stories from the field. In: *The Palgrave Handbook of Critical Menstruation Studies*, London, UK: Palgrave Macmillan 2020:449–65.
54. Hanel PH, Vione KC. Do student samples provide an accurate estimate of the general public? *PLoS One* 2016;11(12):e0168354.
55. Attarchi M, Darkhi H, Kashanian M, et al. Characteristics of menstrual cycle in shift workers. *Global J Health Sci* 2013;5(3):163–72.

56. Figà-Talamanca I. Occupational risk factors and reproductive health of women. *Occup Med* 2006;56(8):521–31.
57. Nohara M, Momoeda M, Kubota T, Nakabayashi M. Menstrual cycle and menstrual pain problems and related risk factors among Japanese female workers. *Ind Health* 2011;49(2):228–34.
58. Thompson LF, Surface EA, Martin DL, Sanders MG. From paper to pixels: Moving personnel surveys to the Web. *Personnel Psychol* 2003;56(1):197–227.
59. Howard PE, Rainie L, Jones S. Days and nights on the Internet: The impact of a diffusing technology. *Am Behav Scientist* 2001;45(3):383–404.
60. Andrews D, Nonnecke B, Preece J. Electronic survey methodology: A case study in reaching hard-to-involve Internet users. *Int J Hum-Comput Interact* 2003;16(2):185–210.
61. Granello DH, Wheaton JE. Online data collection: Strategies for research. *J Counsel Dev* 2004;82(4):387–93.
62. Cosic A, Ferhatovic L, Banozic A, et al. Pain catastrophizing changes during the menstrual cycle. *Psychol Health Med* 2013;18(6):735–41.
63. Campbell CM, Kronfli T, Buenaver LF, et al. Situational versus dispositional measurement of catastrophizing: Associations with pain responses in multiple samples. *J Pain* 2010;11(5):443–53. e2.
64. Durand H, Birnie KA, Noel M, et al. State versus trait: Validating state assessment of child and parental catastrophic thinking about children's acute pain. *J Pain* 2017;18(4):385–95.