

Increased endothelin-1 in colorectal cancer and reduction of tumour growth by ET_A receptor antagonism

E Asham, A Shankar, M Loizidou, S Fredericks³, K Miller¹, PB Boulos, G Burnstock² and I Taylor

Departments of Surgery and ¹Histopathology, The Royal Free and University College Medical School, 67–73 Riding House Street, London W1W 7EJ; ²The Autonomic Neuroscience Institute, The Royal Free and University College Medical School, Rowland Hill Street, London NW3 2PF; ³The Analytical Unit, St George's Hospital Medical School, Blackshaw Road, Tooting, London SW17 0QT, UK

Summary Endothelin-1 (ET-1) is a vasoconstrictor peptide which stimulates proliferation in vitro in different cell types, including colorectal cancer cells. Raised ET-1 levels have been detected both on tissue specimens and in the plasma of patients with cancers. To investigate the role of ET-1 in colorectal cancer: (i) ET-1 plasma levels in patients with colorectal cancer were measured by radioimmunoassay: group 1 = controls ($n = 22$), group 2 = primary colorectal cancer only ($n = 39$), group 3 = liver metastases only ($n = 26$); (ii) ET-1 expression in primary colorectal cancer specimens ($n = 10$) was determined immunohistochemically and (iii) the effect of intraportally infused antagonists to the two ET-1 receptors, ET_A and ET_B, on the growth of liver metastases in a rat model was assessed. ET-1 plasma levels were significantly increased in both patients with primary tumour and patients with metastases, compared to controls ($P < 0.01$, 3.9 ± 1.4 , 4.5 ± 1.5 , vs. 2.75 ± 1.37 pg/ml, respectively). Immunohistochemically, strong expression of ET-1 was found in the cytoplasm, stroma and blood vessels of cancers, unlike the normal colon where only the apical layer of the epithelium, vascular endothelial cells and surrounding stroma were positively stained. In the rat model, there was significant reduction in liver tumour weights compared to controls, following treatment with the ET_A antagonist (BQ123) 30 min after the intraportal inoculation of tumour cells ($P < 0.05$). These results suggest ET-1 is produced by colorectal cancers and may play a role in the growth of colorectal cancer acting through ET_A receptors. ET_A antagonists are indicated as potential anti-cancer agents. © 2001 Cancer Research Campaign <http://www.bjcancer.com>

Keywords: ET-1; colorectal cancer; colorectal liver metastases; ET_A antagonist; ET_B antagonist; BQ123

Endothelin-1 (ET-1), a recognised vasoconstrictor peptide, has been implicated in the growth regulation of several tumours. Its actions are mediated via two receptor subtypes ET_A and ET_B (Yanagisawa et al, 1988; Clozel and Clozel 1989; Clozel et al, 1992). Both receptors have a role in vasomotor control, although only ET_A appears to be involved in the mitogenic action of ET-1 (Ali et al, 2000a; Bagnato et al, 1997, 1999; Nelson et al, 1996; Kitigawa et al, 1994).

ET-1 stimulates proliferation in vitro of fibroblasts, renal mesangial cells, smooth muscle and several tumour cell lines, including colorectal cancer (Hirata et al, 1989; Kusuvara et al, 1989; Simonson et al, 1989; Shichiri et al, 1991; Ali et al, 2000a). Immunohistochemical studies on breast and prostate cancer demonstrate increased expression of ET-1, with patients exhibiting elevated plasma levels of ET-1 (Kojima and Nihei, 1995; Nelson et al, 1995). Other malignant tumours associated with increased ET-1 plasma levels include pancreatic and hepatocellular carcinomas (Oikawa et al, 1994; Nakamuta et al, 1993).

We have previously shown in a pilot study that patients with colorectal cancer, with and without liver metastases, have elevated plasma levels of ET-1 and that the peptide is overexpressed by different cell types within colorectal liver metastases (Shankar et al, 1998). ET-1 binding sites in primary colorectal cancer are known to be unregulated (Inigagi et al, 1992). Recently, we investigated binding to receptor subtypes and showed specifically that ET_A

receptors are overexpressed while ET_B receptors are underexpressed in colorectal cancer tissue compared to normal colon (Ali et al, 2000b). These results suggest ET-1 may play a paracrine or autocrine role in the development of colorectal cancer both locally and systemically.

The purpose of this study is to explore the role of ET-1 in colorectal cancer from three perspectives: firstly, to confirm in a further series of observations the elevated plasma levels of ET-1 in patients with colorectal cancer; secondly, to assess the expression of ET-1 in primary colorectal cancer specimens using immunohistochemistry; and finally to assess the effect of intraportally infused ET_A and ET_B receptor antagonists on the growth of liver metastases in an animal model.

METHODS

Assay for circulating ET-1

Blood collection

Three groups of patients were studied: Group 1—controls ($n = 22$), group 2—patients with primary colorectal cancer without liver metastases on CT scanning or at laparotomy ($n = 39$), group 3—patients with liver metastases whose primary colorectal cancer had already been resected ($n = 26$). Information was collected on serum carcinoembryonic antigen (CEA) levels and Dukes stage, for all patients.

Patients with the following conditions—which are independently associated with elevated ET-1 levels—were excluded;

Received 11 September 2001

Accepted 20 September 2001

Correspondence to: M Loizidou

hypertension, renal failure, heart disease, pre-existing liver disease and Raynauds phenomenon. We have previously demonstrated that elevated ET-1 plasma levels are not associated with endothelial cell trauma (as determined by the plasma levels of thrombomodulin, a marker of endothelial cell integrity) and as such represent an increased production of ET-1 rather than a consequence of tumour invasion and damage of adjacent blood vessels.

All patients had a fasting venous specimen taken prior to surgery. This was collected into precooled blood bottles containing ethylene diamine tetraacetic acid. The specimens were then centrifuged in a chillspin (4° for 15 min at 1000 g) and the plasma frozen at -70° within 1 h of collection.

ET-1 radioimmunoassay

Plasma ET-1 was measured by radioimmunoassay (RIA, Nichols Institute Diagnostics Ltd, Saffron Walden, Essex, UK) according to the manufacturer's instructions. The anti ET-1 antibody had a cross-reactivity of 67% with ET-2, 84% with ET-3, 2.6% with Big ET-1, 5.3% with Big ET-2 and 0.2% with Big ET-3. Endothelin concentrations were calculated by plotting the % bound against log concentrations of known standards.

Immunohistochemistry for ET-1

Specimens of primary colorectal cancer and normal colonic tissue (taken 10 cm away from the tumour) were collected at the time of resection from 10 patients, fixed in neutral buffered formalin for 24 h and paraffin wax embedded.

Sections were cut (5 µm thick), dewaxed in xylene, rehydrated through decreasing concentrations of ethanol and washed in Tris Buffered Saline (TBS, 3 × 5 min). Sections were taken to water and antigen unmasking was achieved by microwaving slides in 600 ml of citrate buffer (0.01 M, pH = 6), for 20 min (full power, 800 W). Hot buffer was flushed out with cold running water and sections were transferred to the incubation chamber and washed with TBS. The protocol for ET-1 staining was as follows: Tissue endogenous peroxidase was blocked by incubating the sections for 10 min in 0.5% methanolic H₂O₂. Sections were washed in running tap water and incubated in mouse monoclonal anti-ET-1 antibody for 1 h (1:100 in TBS, Department of Immunology, University College London, UK, Ong et al, 1993; the antibody cross-reacts with ET-2 and ET-3 (15%), but not big ET). The specimens were washed again in TBS and incubated with rabbit anti-mouse immunoglobulin for 35 min. Routine streptavidin-biotin peroxidase complexing followed (Dako Ltd, Ely, UK) according to the manufacturer's instructions. For the final colour reaction, freshly prepared 3,3 diaminobenzidine (DAB) solution incorporating 0.03% H₂O₂ was applied on the slides (10 min). DAB was rinsed off with TBS and sections were washed in running tap water. Nuclei were counterstained with haematoxylin (4 min), dehydrated and mounted.

Assessment of ET-1 antagonists in an animal model of colorectal liver metastases

In vivo protocol

The animal model used for colorectal liver metastases has been described in detail previously (Loizidou et al, 1991). The intention of this component of the study was to assess the effect of intraportally inoculated antagonists to ET_A and ET_B on the develop-

ment of liver metastases when given at varying times after tumour inoculation. Given the vasoconstricting effect of ET-1 primarily mediated via ET_A, application of ET_A and mixed ET_A/ET_B antagonists lead to vasodilation and subsequent hypotension, while application of the ET_B antagonist lead to hypertension; as such the doses of antagonists used in all arms of the study were those which produced an approximate 20% change in systemic blood pressure (confirmed in a small series of experiments in our lab). This level of blood pressure change was arbitrarily chosen as the safety limit. The study was conducted under a Home Office approved project licence.

Male Hooded Lister rats weighing between 250–300 g were anaesthetised using halothane inhalation. A lower midline incision was then made and a tributary of the ileocolic vein was cannulated. All animals received a bolus injection of 1 × 10⁶ MC28 syngeneic tumour cells in 0.2 ml saline (the dose known to produce reproducible tumour growth by 14 days, Loizidou et al, 1991).

In the first arm of the study the effect of infusions of ET-1 antagonists 30 min after tumour inoculation were assessed. The animals were divided into 5 groups (6 animals per group) and received; BQ123 (Alexis Corporation, Nottingham, UK) an ET_A antagonist, PD142893 (Alexis Corporation) a nonselective ET_A and ET_B antagonist, A-192621 an ET_B receptor antagonist (kindly donated by Abbott, Chicago, IL, USA), saline only and saline plus an intramuscular injection of 0.1 mg/kg acepromazine, (National Veterinary Supplies, Stroke-on-Trent, UK) a hypotensive agent which acts as a control for the vascular effects of ET-1 antagonism. Each antagonist was infused at a rate of 100 µg/kg/min for 30 min.

In the second arm of the study, the ileocolic vein was cannulated after tumour cell inoculation was performed. A fine 2F gauge portacath (Becton-Dickinson UK Ltd, Oxford, UK) was tunnelled under the skin to the interscapular area. This allowed access to the portal circulation without the need for a further laparotomy. The animals were then divided into 4 groups (6 animals per group) with infusions of agents given via the portal circulation on days 2, 4 and 6 after tumour inoculation. The infusions were administered under halothane anaesthesia for 30 min, at the same rate as in the previous study. The groups were; saline, BQ123, PD142893, A-192621 and saline plus intramuscular acepromazine.

Given that different passages of the same tumour produce different degrees of percentage hepatic replacement with tumour, in order to compare the different agents one animal from each group received tumour from the same passage. This allowed comparisons to be made between the animals in each of the groups and was applied to both arms of the study.

All animals were sacrificed 14 days after intraportal tumour inoculation, the livers were harvested and the tumours dissected out. Wet weights were obtained for uninvolved liver and for total tumour, per animal. The percent hepatic replacement (PHR) with tumour was calculated for every animal, by dividing tumour weight by total liver (i.e., liver plus tumour) weight and then multiplying by 100. Results were presented as ratios between PHRs of experimental animals to PHRs of control animals.

RESULTS

Plasma levels of ET-1

The plasma levels of ET-1 in the control group (age range 42–82 years, median age 59.5 years, 8 females and 14 males, *n* = 22) had a mean of 2.75 pg/ml, (SD = 1.37). In the colorectal cancer group




Figure 1 Levels of ET-1 in plasma from patients with or without colorectal cancer. Individual values are shown for controls (Δ, *n* = 22), patients with primary colorectal cancer (○, *n* = 39), and patients with previously removed primary colorectal cancer who have subsequently developed liver metastases (□, *n* = 26). Bars are means (± SD). Significance was reached with both patient groups compared to controls (*P* < 0.01, Mann-Whitney test)

without metastases (median age 64.6 years, range 55–77 years, 16 females and 23 males, *n* = 39) the mean ET-1 level was 3.9 pg/ml, (SD = 1.4) and in the group of patients with liver metastases (median age 55, range 46–75 years, 3 females and 23 males, *n* = 26) the mean ET-1 level was 4.5 pg/ml (SD = 1.5), as illustrated in Figure 1. There was a significant difference between the cancer patients and the control group (*P* < 0.001 and *P* < 0.01 for the metastases and primary cancer groups, respectively) whilst there was no significant difference between the two groups of cancer patients.

In the colorectal cancer group without metastases, there were 3 Dukes A, 26 Dukes B, and 10 Dukes C patients with CEA levels ranging from <1–43 (median = 2). All 26 patients with liver metastases were classed as Dukes D, by definition, with CEA levels recorded between <1–1650 (median = 11). No correlation was found between ET-1 levels and serum CEA or Dukes stage.

ET-1 Immunohistochemistry

Five patients had Dukes B and 5 had Dukes C carcinomas (3 females and 7 males). ET-1 immunohistochemistry of control tissue showed ET-1 present in the apical layer of colonic epithelium in 5 of 10 patients studied, vascular endothelial cells and surrounding stroma (Figure 2A). In the colorectal cancer specimens there was strong expression of ET-1 in the cytoplasm of epithelial cells, stroma and blood vessels, in all the tumour specimens studied (Figure 2B). There was no correlation between Dukes staging and intensity of staining.

ET-1 antagonists

Day 0 infusion group

The only group which demonstrated a significant reduction in tumour load compared to the saline control group (paired *t*-test) was those animals which received BQ123. The mean ratio (compared to the saline control group) was 0.64 ± 0.20, (*P* < 0.05, paired *t*-test).




Figure 2 ET-1 immunoreactivity in colon. (A) Control: note ET-1 staining (brown) limited to the apical segment of the colonic epithelium (arrows) together with some staining of the stroma (S). (B) Colorectal cancer showing strong staining for ET-1, in the cytoplasm of epithelial cancer cells (arrows) and in the stroma (S). Counterstained with haematoxylin. Magnification × 700

PD142893 gave a mean of 1.10 ± 0.9 (NS), A-192621 was 1.13 ± 0.88 (NS) and the saline/acepromazine group was 1.4 ± 0.70 (NS). These results are illustrated in Figure 3, which demonstrates reductions in tumour load compared to the saline only group.




Figure 3 Liver tumours in rats which received Endothelin receptor antagonists. After tumour cell inoculation, animals received one of the following: saline (controls □, dotted line), BQ123 (○), PD142893 (Δ), A-192621 (▽), or saline plus the hypotensive agent acepromazine (+) (*n* = 6 per group). Fourteen days later, liver tissue (tumour and normal) was weighed. Results (Y-axis) are shown as a ratio of the % hepatic replacement by tumour (PHR) in the experimental animals compared to control PHR. The horizontal dotted line represents the average (mean) of control values (SD = 0.23). The BQ123 group had significantly smaller tumours than the controls (*P* < 0.05, paired Student *t*-test)

Day 2, 4 and 6 group

In this group of experiments, none of the agents produced significant reductions in tumour load compared to the control group. The mean values (SD) were 1.11 ± 0.81 for BQ123, 1.02 ± 0.58 for PD142893, and 1.80 ± 1.90 for the saline/acepromazine group.

DISCUSSION

ET-1 is a vasoconstrictor peptide the physiological functions of which are well described. ET-1 acts via two G-linked protein receptors ET_A and ET_B , with ET_A mediating vasoconstriction and ET_B vasodilation (Yanagisawa et al, 1988; Clozel and Clozel, 1989; Clozel et al, 1992).

Recently ET-1 has been implicated as a possible regulator of tumour growth in a number of human malignancies including colorectal cancer (Bagnato et al, 1997, 1999; Kojima and Nihei, 1995; Nelson et al, 1995, 1996, 1999; Oikawa et al, 1994; Nakamura et al, 1993; Shankar et al, 1998; Ali et al, 2000a). A variety of tumour cell lines including colorectal cancer produce ET-1, which when added to the growth medium results in an increase in tumour cell turnover (Shichiri et al, 1991; Bagnato et al, 1997; Moraitis et al, 1977). Studies involving colorectal, ovarian and prostate cancer suggest that the receptor responsible for the ET-1 mitogenic action is ET_A , which is probably upregulated (Bagnato et al, 1997, 1999; Nelson et al, 1995; Ali et al, 2000a). Binding of ligand to receptor usually activates phospholipase C followed by Ca^{++} /protein kinase C signalling. It has also been shown that ET-1 acts as a co-mitogen with other factors such as epidermal growth factor, EGF (Bagnato et al, 1997). Synergism might be in part explained by the fact that Ca^{++} /protein kinase C cross-talk with (activate) molecules belonging to EGF-initiated tyrosine kinase cascades within the cell (Bagnato et al, 1997; Luttrell et al, 1999). In a recent study (Eberl et al, 2000), a mixed ET_A/ET_B receptor antagonist potentiated FasL-induced apoptosis in cultured colorectal cancer cells. This suggests that ET-1 may not only enhance growth, but also protect against apoptosis in certain cell types.

In colorectal cancer Inagaki et al (1992) demonstrated a high density of ET-1 binding sites within tumour vessels and stromal tissues surrounding primary colorectal cancers. Shankar et al (1998) using immunoelectron microscopy have demonstrated increased expression of ET-1 by a variety of cell types within colorectal liver metastases and that patients with colorectal cancer with and without liver metastases had elevated plasma levels of ET-1. Such findings suggest ET-1 may act by a paracrine/autocrine mechanism in these tumours.

This study has extended the earlier work with regard to elevated plasma levels of ET-1 in patients with colorectal cancer (Shankar et al, 1998) and also shows that primary tumours as well as liver metastases produce ET-1. These findings complement recently published data on Big ET-1, the precursor of ET-1, in colorectal cancer (Simpson et al, 2000). The authors found significantly higher plasma levels of Big ET-1 in patients with cancer compared to controls and that tissues from colorectal cancer overexpress Big ET-1, compared to controls. Immunohistochemically, we demonstrated that in colorectal cancer tissue, cancer epithelial cells, endothelial cells and stromal cells overexpress ET-1, compared to normal colon. The fact that, in some cases, normal colonic mucosa also expresses ET-1 suggests it may play a role in the control of normal cell shedding and perhaps apoptosis as suggested by other groups (Shichiri et al, 1998). It has been proposed that primary tumours control the growth of distant metastases by the secretion

of a variety of messenger compounds (Hanahan and Folkman, 1998) and ET-1 may have such a role.

During the metastatic cascade, tumour emboli are shed from primary colorectal cancers and reach the liver via the portal circulation. These micrometastases initially obtain their blood supply from the portal vein. This is the basis for the rationale behind peri-operative portal vein chemotherapy (Shankar et al, 1996).

Since ET-1 appears to play a role in colorectal cancer development, administration of intraportally infused ET-1 antagonists at this micrometastatic phase might produce a reduction in hepatic tumour load. We have previously shown that MC28 tumours possess similar anatomical and physiological profiles to colorectal liver metastases and also express elevated levels of ET-1 (Loesch et al, 1997; Ashraf et al, 1997). As such, the behaviour of these tumours should mimic those of colorectal cancer.

The findings in this study confirm the potential involvement of ET_A receptors in the development of colorectal liver metastases. Only the ET_A antagonist BQ123 produced a significant reduction in tumour weight. Also of interest is the timing of delivery; since only BQ123 given at the time of inoculation, but not if administered at 2, 4, or 6 days after tumour inoculation, produced any effect. These findings are in keeping with previous studies involving cytotoxic portal vein infusions designed to prevent hepatic tumour development, and suggest the importance of perinoculation infusions compared to "delayed" treatment (Sutanto-Ward et al, 1992). One explanation for this finding is that after 24–48 h these tumours may switch their blood supply from the portal vein to the hepatic artery and hence insufficient antagonist reaches the tumour. Another suggestion is that the role of ET-1 in metastatic development is to promote tumour implantation and the initial establishment of micrometastases. Accordingly, the administration of these antagonists after this stage would have little effect. Further experimentation with different antagonists and different in vivo models would be necessary to elucidate possible mechanisms of action. Use of recently reported ET_A antagonists that could be delivered systemically (or orally) may resolve this issue (Liu et al, 1998; Wilson et al, 1999).

The absence of any effect in the hypotensive control group suggests that changes in blood flow caused by ET-1 receptor antagonism are not the cause of the effect seen.

These results suggest that ET-1 is produced by colorectal cancer, and may play a role both locally and systemically in cancer development by acting through ET_A receptors. In addition application of ET_A antagonists via the portal vein at the time of tumour implantation reduces subsequent hepatic involvement in this model of colorectal liver metastases.

ACKNOWLEDGEMENTS

We are grateful to the Middlesex Hospital Special Trustees, London, W1, for financial support for this project. We thank Abbott Laboratories, Chicago, Illinois for donating endothelin receptor antagonists. We thank Dr R. Jordan and Mr T. Robson for their invaluable help with imaging.

REFERENCES

- Ali H, Loizidou M, Dashwood M, Savage F, Sheard C and Taylor I (2000a) Stimulation of colorectal cancer cell line growth by ET-1 and its inhibition by ET_A antagonists. *Gut* **47**: 685–688

- Ali H, Dashwood M, Dawas K, Loizidou M, Savage F and Taylor I (2000b) Endothelin receptor expression in colorectal cancer. *J Cardiovasc Pharm* **36**: S69–S71
- Ashraf S, Loizidou M, Crowe R, Turmaine M, Burnstock G and Taylor I (1997) Blood vessels in liver metastases from both sarcoma and carcinoma lack perivascular innervation and smooth muscle cells. *Clin Exp Metastasis* **15**: 484–498
- Bagnato A, Tecce R, Dicastro V and Catt KJ (1997) Activation of mitogenic signalling by endothelin-1 in ovarian carcinoma cells. *Cancer Res* **57**: 1306–1311
- Bagnato A, Salani D, Di Castro V, Wu-Wong JR, Tecce R, Nicotra MR, Kenuti A and Natali PG (1999) Expression of ET-1 and ET_A receptor in ovarian carcinoma: evidence for an autocrine role in tumour growth. *Cancer Res* **59**: 720–727
- Clozel JP and Clozel M (1989) Effects of endothelin on the coronary vascular bed in open-chest dogs. *Circ Res* **65**: 1193–1200
- Clozel M, Gray GA, Breu V, Löffler BM and Osterwalder R (1992) The endothelin ET_B receptor mediates both vasodilation and vasoconstriction *in vivo*. *Biochem Biophys Res Commun* **186**: 867–873
- Eberl LP, Valdenaire O, Saintgiorgio V, Jeannin J-F and Juillerat-Jeanneret L (2000) Endothelin receptor blockade potentiates FasL-induced apoptosis in rat colon carcinoma cells. *Int J Cancer* **86**: 182–187
- Hanahan D and Folkman J (1996) Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *Cell* **86**: 353–364
- Hirata Y, Takagi Y, Fukuda Y and Marumi F (1989) Endothelin is a potent mitogen for rat vascular smooth muscle cells. *Atherosclerosis* **78s**: 225–228
- Inigagi H, Bishop AE, Eimoto T and Polak JM (1992) Autoradiographic localization of ET-1 binding sites in human colonic cancer tissue. *J Pathol* **168**: 263–267
- Kitigawa KN, Tsutsumi K, Niwa M, Yamaga S, Anda T and Khalid H (1994) A selective ET_A antagonist, BQ123, inhibits ¹²⁵I-ET-1 binding of human meningiomas and antagonizes ET-1 induced proliferation. *Cell Mol Neurobiol* **14**: 105–118
- Kojima K and Nihei Z (1995) Expression of ET-1 immunoreactivity in breast cancer. *Surg Oncol* **4**: 309–315
- Kusuhara M, Yamaguchi K, Ohnishi A, Abe K, Kimura S, Oono H, Hori S and Nakamura Y (1989) Endothelin potentiates growth factor stimulated DNA synthesis in Swiss 3T3 cells. *Jpn J Cancer Res* **80**: 302–305
- Liu G, Henry KH Jr, Szczepankiewicz BG, Winn M, Kozmina NS, Boyd SA, Wasicak J, von Geldern TW, Wu-Wong JR, Chiou WJ, Dixon DG, Nguyen B, Marsh KC and Opgenorth TJ (1998) Pyrrolidine-3-carboxylic acids as endothelin antagonists. 3. Discovery of a potent 2-nonyl, highly selective ET_A antagonist (A-216546). *J Med Chem* **41**: 3261–3275
- Loesch A, Turmaine M, Loizidou M, Crowe R, Ashraf S, Taylor I and Burnstock G (1997) Increase in immunoreactivity for ET-1 in blood vessels of rat liver metastases: experimental sarcoma and carcinoma. *J Anat* **191**: 291–299
- Loizidou MC, Lawrance RJ, Holt S, Carty NJ, Cooper AJ, Alexander P and Taylor I (1991) Facilitation by partial hepatectomy of tumour growth within the rat liver following intraportal injection of syngeneic tumour cells. *Clin Exp Metastasis* **9**: 335–349
- Luttrell LM, Daaka Y and Lefkowitz J (1999) Regulation of tyrosine kinase cascades by G protein coupled receptors. *Curr Opin Cell Biol* **11**: 177–183
- Moraitis S, Langdon SP and Miller WR (1997) Endothelin expression and responsiveness in human ovarian cancer cell lines. *Eur J Cancer* **33**: 661–668
- Nakamuta M, Ohashi M, Tabata S, Tanabe Y, Goto K, Naruse M, Naruse K, Hiroshige K and Nawata H (1993) High plasma concentrations of ET-like immunoreactivities in patients with hepatocellular carcinoma. *Am J Gastroenterol* **88**: 248–252
- Nelson JB, Hedican SP, George DJ, Reddi AH, Piantadosi S, Eisenberger MA and Simon JW (1995) Identification of ET-1 in the pathophysiology of metastatic adenocarcinoma of the prostate. *Nat Med* **1**: 944–949
- Nelson JB, Chan-Tack K, Hedican SP, Magnuson SR, Opgenorth TJ, Bova GS and Simons JW (1996) Endothelin-1 production and decreased ETB receptor expression in advanced prostatic cancer. *Cancer Res* **56**: 663–668
- Oikawa T, Kushuhara M, Ishikawa S, Hitomi J, Kono A, Iwanaga T and Yamaguchi K (1994) Production of ET-1 and thrombomodulin by human pancreatic cancer cells. *Br J Cancer* **69**: 1059–1064
- Ong A, Jowett T, Scoble J, O'Shea P, Varghese Z and Moorhead J (1993) Effect of cyclosporin A on endothelin synthesis by cultured human renal cortical epithelial cells. *Nephrol Dial Transplant* **8**: 748–753
- Shankar A, Loizidou M and Taylor I (1996) The vascularity of colorectal liver metastases. *Eur J Surg Oncol* **22**: 389–396
- Shankar A, Loizidou M, Aliev G, Fredericks S, Holt D, Boulos PB, Burnstock B and Taylor I (1998) Raised ET-1 levels in patients with colorectal liver metastases. *Br J Surg* **85**: 502–506
- Shichiri M, Hirata Y, Nakajima T, Ando K, Imai T, Yanagisawa M, Masaki T and Marumo F (1991) Endothelin-1 is an autocrine/paracrine growth factor for human cancer cell lines. *J Clin Invest* **87**: 1867–1871
- Shichiri M, Marumo F and Yukio H (1998) Endothelin-B receptor-mediated suppression of endothelial apoptosis. *J Cardiovasc Pharmacol* **31**: S138–S141
- Simonson MS, Wamm S, Mene P, Dubyak GR, Kester M, Nakazato Y, Sedor JR and Dunn MJ (1989) Endothelin stimulated phospholipase C, Na⁺/H⁺ exchange c-fos expression and mitogenesis in rat mesangial cells. *J Clin Invest* **83**: 708–712
- Simpson RA, Dickinson T, Porter KE, London NJM and Hemingway DM (2000) Raised levels of plasma Big ET-1 in patients with colorectal cancer. *Br J Surg* **87**: 1409–1413
- Sutanto-Ward E, Sigurdson ER, Tremiteira S, Lincer R, Chapman D and Niedzwiecki D (1992) Adjuvant chemotherapy for colorectal hepatic metastases: role of route of administration and timing. *Surg Oncol* **1**: 87–95
- Wilson J, Hunt SJ, Tang E, Wright N, Kelly E, Palmer S, Heys C, Mellor S, James R and Bialecki R (1999) Pharmacological profile of 2D1611, an orally active ET_A antagonist. *J Pharmacol Exp Ther* **290**: 1085–1091
- Yanagisawa M, Kurihara H, Kimura S, Tomobe Y, Kobayashi M, Mitsui Y, Yazaki Y, Goto K and Masaki T (1988) A novel potent vasoconstrictor peptide produced by vascular endothelial cells. *Nature* **332**: 411–415