

**Guest editorial:**

**HIGHLIGHT REPORT:  
PROTECTION OF CHOLESTATIC LIVERS BY SHUNTING OF BILE  
FROM CANALICULI TO SINUSOIDS**

Tahany Abbass<sup>1</sup>, Walaa Murad<sup>1</sup>, Abdel-latif Seddek<sup>2\*</sup>

<sup>1</sup> Histology Department, Faculty of Medicine, South Valley University, Qena, Egypt

<sup>2</sup> Forensic Medicine and Toxicology Department, Faculty of Veterinary Medicine, South Valley University, Qena, Egypt

\* Corresponding author: Abdel-latif Seddek, Department of Forensic Medicine and Toxicology, Faculty of Veterinary Medicine, South Valley University, Qena, Egypt;  
E-mail: [tahany\\_abbass@yahoo.com](mailto:tahany_abbass@yahoo.com); [walaa.ahmed@med.svu.edu.eg](mailto:walaa.ahmed@med.svu.edu.eg);  
[abdellatief-shakir@vet.svu.edu.eg](mailto:abdellatief-shakir@vet.svu.edu.eg)

<http://dx.doi.org/10.17179/excli2018-1738>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

Cholestatic liver diseases have been reported to show an anatomically ascending course of disease (Jansen et al., 2017). The first lesions occur in downstream bile ducts followed by damage of the upstream liver parenchyma. In the acute phase after obstruction this may lead to a strong increase of bile acids in bile canaliculi up to concentrations that are toxic to hepatocytes. In the recent issue of *Hepatology* an interesting mechanism has been reported that prevents deterioration of liver tissue in this acute phase of cholestasis (Ghallab et al., 2018). After bile duct ligation in mice bile acid concentrations increase into cytotoxic range of 0.2–2 mM. Between days 1–3 after bile duct ligation the apical hepatocyte membrane ruptures in individual, dispersed hepatocytes. Within minutes the membrane of the same cell also becomes leaky at the sinusoidal/blood side. This creates a shunt between the bile canaliculus and the sinusoidal blood (Ghallab et al., 2018). As a consequence, bile acids leak from overloaded canaliculi into the blood. Therefore, bile acid concentrations in the canaliculi decrease after

shunt formation. In principle, the acute cholestatic liver sacrifices some dispersed individual hepatocytes that later can easily be regenerated in order to save the entire organ.

Analysis of the mechanisms of hepatotoxicity represents an intensively studied field of research (Fickert et al., 2014; Tag et al., 2015; Stöber, 2016; Ghallab, 2017; Ghallab et al., 2016; Vartak et al., 2016). In recent years procedures of mathematical modeling (Schenk et al., 2017; Sezgin et al., 2018; Bartl et al., 2015; Schliess et al., 2014) and advanced imaging techniques such as combined MALDI and two-photon microscopy as applied in the highlighted study (Ghallab et al., 2018) have contributed to a deeper understanding of the pathomechanisms of liver diseases. Although, the here described mechanism of physical shunting of bile salts from high to low concentration areas protects the liver from more severe damage, it should be considered that this leads to higher concentrations in blood with the long term consequence of kidney damage usually in the form of cholemic nephropathy.

**Conflict of interest**

There is no conflict of interest.

**REFERENCES**

- Bartl M, Pfaff M, Ghallab A, Driesch D, Henkel SG, Hengstler JG, et al. Optimality in the zonation of ammonia detoxification in rodent liver. *Arch Toxicol.* 2015;89:2069-78.
- Fickert P, Pollheimer MJ, Beuers U, Lackner C, Hirschfield G, Housset C, et al. Characterization of animal models for primary sclerosing cholangitis (PSC). *J Hepatol.* 2014;60:1290-303.
- Ghallab A. Highlight report: Metabolomics in hepatotoxicity testing. *EXCLI J.* 2017;16:1323-5.
- Ghallab A, Cellière G, Henkel SG, Driesch D, Hoehme S, Hofmann U, et al. Model-guided identification of a therapeutic strategy to reduce hyperammonemia in liver diseases. *J Hepatol.* 2016;64:860-71.
- Ghallab A, Hofmann U, Sezgin S, Vartak N, Hassan R, Zaza A, et al. Bile micro-infarcts in cholestasis are initiated by rupture of the apical hepatocyte membrane and cause shunting of bile to sinusoidal blood. *Hepatology.* 2018 Aug 13, epub ahead of print.
- Jansen PL, Ghallab A, Vartak N, Reif R, Schaap FG, Hampe J, et al. The ascending pathophysiology of cholestatic liver disease. *Hepatology.* 2017;65:722-38.
- Schenk A, Ghallab A, Hofmann U, Hassan R, Schwarz M, Schuppert A, et al. Physiologically-based modeling in mice suggests an aggravated loss of clearance capacity after toxic liver damage. *Sci Rep.* 2017;7(1):6224.
- Schliess F, Hoehme S, Henkel SG, Ghallab A, Driesch D, Böttger J, et al. Integrated metabolic spatial-temporal model for the prediction of ammonia detoxification during liver damage and regeneration. *Hepatology.* 2014;60:2040-5.
- Sezgin S, Hassan R, Zühlke S, Kuepfer L, Hengstler JG, Spittler M, et al. Spatio-temporal visualization of the distribution of acetaminophen as well as its metabolites and adducts in mouse livers by MALDI MSI. *Arch Toxicol.* 2018;92:2963-77.
- Stöber R. Pathophysiology of cholestatic liver disease and its relevance for in vitro tests of hepatotoxicity. *EXCLI J.* 2016;15:870-1.
- Tag CG, Sauer-Lehnen S, Weiskirchen S, Borkham-Kamphorst E, Tolba RH, Tacke F, et al. Bile duct ligation in mice: induction of inflammatory liver injury and fibrosis by obstructive cholestasis. *J Vis Exp.* 2015;96.
- Vartak N, Damle-Vartak A, Richter B, Dirsch O, Dahmen U, Hammad S, et al. Cholestasis-induced adaptive remodeling of interlobular bile ducts. *Hepatology.* 2016;63:951-64.