

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-Amino-6-methylpyridinium 2,2,2-trichloroacetate

K. Syed Suresh Babu,<sup>a,b</sup> G. Peramaiyan,<sup>a</sup> M. NizamMohideen<sup>b\*</sup> and R. Mohan<sup>a\*</sup><sup>a</sup>Department of Physics, Presidency College (Autonomous), Chennai 600 005, Tamil Nadu, India, and <sup>b</sup>Department of Physics, The New College (Autonomous), Chennai 600 014, Tamil Nadu, India

Correspondence e-mail: mnizam\_new@yahoo.in, professorsomohan@yahoo.co.in


Received 26 February 2014; accepted 27 February 2014

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.046;  $wR$  factor = 0.129; data-to-parameter ratio = 19.5.

In the asymmetric unit of the title molecular salt,  $\text{C}_6\text{H}_9\text{N}_2^+ \cdot \text{C}_2\text{Cl}_3\text{O}_2^-$ , there are two independent 2-amino-6-methylpyridinium cations and two independent trichloroacetate anions. The pyridine N atom of the 2-amino-6-methylpyridine molecule is protonated and the geometries of these cations reveal amine–imine tautomerism. Both protonated 2-amino-6-methylpyridinium cations are essentially planar [maximum deviations = 0.026 (2) and 0.012 (2) Å]. In the crystal, the protonated N atom and the 2-amino group of the cation are hydrogen bonded to the carboxylate O atoms of the anion via a pair of  $\text{N}-\text{H} \cdots \text{O}$  hydrogen bonds, forming an  $R_2^2(8)$  ring motif. These motifs are connected via  $\text{N}-\text{H} \cdots \text{O}$  and  $\text{C}-\text{H} \cdots \text{O}$  hydrogen bonds to form slabs parallel to (101).

## Related literature

For applications of pyridinium derivatives, see: Akkurt *et al.* (2005). For pyridine derivatives as templating agents, see: Desiraju (2001); Jeffrey (1997). For details of 2-aminopyridine and its derivatives, see: Katritzky *et al.* (1996); Tomaru *et al.* (1991). For bond lengths and angles in similar structures, see: Jin, Shun *et al.* (2005); Feng *et al.* (2007); Nahringsbauer & Kvik (1977). For other 2-aminopyridinium structures, see: Jin *et al.* (2000, 2001); Jin, Tu *et al.* (2005). For studies on the tautomeric forms of 2-aminopyridine systems, see: Ishikawa *et al.* (2002). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



## Experimental

## Crystal data

$\text{C}_6\text{H}_9\text{N}_2^+ \cdot \text{C}_2\text{Cl}_3\text{O}_2^-$   
 $M_r = 271.52$   
 Monoclinic,  $P2_1/n$   
 $a = 11.6376$  (5) Å  
 $b = 14.6648$  (6) Å  
 $c = 13.9100$  (6) Å  
 $\beta = 96.024$  (1)°

$V = 2360.81$  (17) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.76$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.35 \times 0.30 \times 0.30$  mm

## Data collection

Bruker Kappa APEXII CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 2004)  
 $T_{\min} = 0.777$ ,  $T_{\max} = 0.805$

39609 measured reflections  
 5784 independent reflections  
 3922 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.032$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$   
 $wR(F^2) = 0.129$   
 $S = 1.04$   
 5784 reflections  
 297 parameters  
 6 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.66$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.64$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N1}-\text{H1A} \cdots \text{O3}^{\text{i}}$	0.90 (1)	1.96 (1)	2.856 (3)	172 (3)
$\text{N2}-\text{H2B} \cdots \text{O4}^{\text{i}}$	0.89 (1)	1.95 (1)	2.825 (3)	167 (3)
$\text{N2}-\text{H2A} \cdots \text{O1}$	0.89 (1)	2.11 (1)	2.982 (3)	167 (3)
$\text{N3}-\text{H3A} \cdots \text{O2}$	0.90 (1)	1.84 (1)	2.729 (2)	175 (3)
$\text{N4}-\text{H4B} \cdots \text{O1}$	0.88 (1)	2.06 (1)	2.929 (3)	167 (3)
$\text{N4}-\text{H4A} \cdots \text{O3}^{\text{ii}}$	0.88 (1)	2.29 (2)	3.096 (3)	152 (3)
$\text{C6}-\text{H6C} \cdots \text{O1}^{\text{iii}}$	0.96	2.50	3.413 (4)	158
$\text{C11}-\text{H11} \cdots \text{O4}^{\text{iv}}$	0.93	2.50	3.371 (4)	157

Symmetry codes: (i)  $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (ii)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (iii)  $-x + 1, -y, -z$ ; (iv)  $-x, -y, -z + 1$ .

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT and XPREP (Bruker, 2004); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: WinGX (Farrugia, 2012) and PLATON (Spek, 2009).

The authors are thankful to the SAIF, IIT Madras, for the data collection.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU2705).

## References

- Akkurt, M., Karaca, S., Jarrahpour, A. A., Zarei, M. & Büyükgüngör, O. (2005). *Acta Cryst.* **E61**, o776–o778.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2004). *APEX2*, *SAINTE* and *XPREF*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Desiraju, G. R. (2001). *Curr. Sci.* **81**, 1038–1055.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Feng, W.-J., Wang, H.-B., Ma, X.-J., Li, H.-Y. & Jin, Z.-M. (2007). *Acta Cryst.* **E63**, m1786–m1787.
- Ishikawa, H., Iwata, K. & Hamaguchi, H. (2002). *J. Phys. Chem. A*, **106**, 2305–2312.
- Jeffrey, G. A. (1997). In *An Introduction to Hydrogen Bonding*. Oxford University Press.
- Jin, Z.-M., Pan, Y.-J., Hu, M.-L. & Shen, L. (2001). *J. Chem. Crystallogr.* **31**, 191–195.
- Jin, Z.-M., Pan, Y.-J., Liu, J.-G. & Xu, D.-J. (2000). *J. Chem. Crystallogr.* **30**, 195–198.
- Jin, Z.-M., Shun, N., Lü, Y.-P., Hu, M.-L. & Shen, L. (2005). *Acta Cryst.* **C61**, m43–m45.
- Jin, Z.-M., Tu, B., He, L., Hu, M.-L. & Zou, J.-W. (2005). *Acta Cryst.* **C61**, m197–m199.
- Katritzky, A. R., Rees, C. W. & Scriven, E. F. V. (1996). In *Comprehensive Heterocyclic Chemistry II*. Oxford: Pergamon Press.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Nahringbauer, I. & Kvik, Å. (1977). *Acta Cryst.* **B33**, 2902–2905.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Tomaru, S., Matsumoto, S., Kurihara, T., Suzuki, H., Oobara, N. & Kaino, T. (1991). *Appl. Phys. Lett.* **58**, 2583–2585.

## supplementary materials

*Acta Cryst.* (2014). E70, o391–o392 [doi:10.1107/S1600536814004553]

## 2-Amino-6-methylpyridinium 2,2,2-trichloroacetate

K. Syed Suresh Babu, G. Peramaiyan, M. NizamMohideen and R. Mohan

### 1. Comment

2-Aminopyridine and its derivatives play an important role in heterocyclic chemistry (Katritzky *et al.*, 1996). Pyridine heterocycles and their derivatives are present in many large molecules having photo-chemical, electro-chemical and catalytic applications. Some pyridine derivatives possess nonlinear optical (NLO) properties (Tomaru *et al.*, 1991) and often possess antibacterial and antifungal activities (Akkurt *et al.*, 2005). The use of pyridine derivatives as templating agents for the self-assembly of organic-inorganic supramolecular materials has been widely studied (Desiraju, 2001). They are often involved in hydrogen bonding (Jeffrey, 1997). In order to further study the hydrogen bonding in these systems, the synthesis and structure of the title salt is presented herein.

The asymmetric unit of title compound, Fig. 1, consists of two crystallographically independent protonated 2-amino-6-methylpyridinium cation and trichloroacetate anion. The pyridinium rings are almost planar with mean deviations of only 0.011 (2) and 0.007 (2) Å for atoms N1 and N3, respectively. The dihedral angle between a 2-amino-6-methylpyridinium cation and trichloroacetate anion group is 18.3 (2) and 47.7 (2)° for the both molecules respectively. In both the molecules, the protonated 2-amino-6-methylpyridinium cations are essentially planar, with a maximum deviation of 0.026 (2) Å for atom N1 and 0.012 (2) Å for atom N3.

In the cations, the N2—C1 [1.324 (4) Å] and N4—C9 bonds [1.335 (3) Å] are shorter than the N1—C1 [1.357 (3) Å], N1—C5 [1.361 (3) Å], N3—C9 [1.341 (3) Å] and N3—C13 [1.360 (3) Å] bonds, and the C1—C2 [1.394 (4) Å], C3—C4 [1.401 (5) Å], C9—C10 [1.404 (3) Å] and C11—C12 [1.385 (5) Å] bonds are significantly longer than bonds C2—C3 [1.346 (5) Å], C4—C5 [1.360 (4) Å], C10—C11 [1.350 (5) Å] and C12—C13 [1.362 (4) Å]. This are similar to the bond distances observed previously for the amino-methylpyridinium cation (Jin, Shun *et al.*, 2005; Feng *et al.*, 2007). In contrast, in the solid state structure of amino-methylpyridinium, the exocyclic N—C bond is clearly longer than that in the ring (Nahringbauer & Kvik, 1977). The geometric features of amino-methylpyridinium cation (N1/N2/C1/C6 and N3/N4/C9—C13) resemble those observed in other 2-aminopyridinium structures (Jin *et al.*, 2000; Jin *et al.*, 2001; Jin, Tu *et al.*, 2005) that are believed to be involved in amine-imine tautomerism (Ishikawa *et al.*, 2002). Similar features are also provided by the cation amino-methylpyridinium (N3/N4/C7/C12). However, a previous study shows that a pyridinium cation always possesses an expanded angle of C—N—C [C1N1—C5 [123.7 (2)°] and C9—N3—C13 [124.6 (2)°] in comparison with the parent pyridine (Jin, Shun *et al.*, 2005).


In the crystal, the protonated atoms (N1 and N3) and a nitrogen atom of the 2-amino group (N2 and N4) are hydrogen bonded to the carboxylate oxygen atoms (O1, O2, O3 and O4) *via* pairs of intermolecular N—H—O [(N1—H1A···O3 and N3—H3A···O2) and (N2—H2B···O4 and N4—H4B···O1)] hydrogen bonds (Fig. 2 and Table 1), forming an  $R_2^2(8)$  ring motif (Bernstein *et al.*, 1995). Furthermore, these motifs are connected *via* N2—H2A···O1, N4—H4A···O3 and C—H···O hydrogen bonds to form slabs parallel with (101) [Fig. 2 and Table 1].

## 2. Experimental

Crystals of the title compound were obtained by slow evaporation of a 1:1 mol. mixture of 2-Amino-6-methylpyridine and trichloroacetic acid in methanol at room temperature.


## 3. Refinement

N-bound H atoms were located in a difference Fourier map and refined with distance restraints: N—H = 0.89 (1) and 0.90 (1) Å for NH<sub>2</sub> and NH H atoms, respectively; N—H distances of the NH<sub>2</sub> groups were restrained to be equal. The C-bound H atoms were positioned geometrically and refined using a riding model: C—H = 0.93–0.96 Å with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C-methyl})$  and  $= 1.2U_{\text{eq}}(\text{C})$  for other H atoms. A rotating group model was used for the methyl group.



**Figure 1**

A view of the molecular structure of the two independent trichloroacetate anions and the two independent 2-amino-6-methylpyridinium cations of the title salt. Displacement ellipsoids are drawn at the 30% probability level.



**Figure 2**

The crystal packing of the title compound, viewed normal to (101). The N—H···O and C—H···O hydrogen bonds are shown as dashed lines (see Table 1 for details; H atoms not involved in hydrogen bonding have been omitted for clarity).

### 2-Amino-6-methylpyridinium 2,2,2-trichloroacetate

#### Crystal data

$C_6H_9N_2^+ \cdot C_2Cl_3O_2^-$

$M_r = 271.52$

Monoclinic,  $P2_1/n$

Hall symbol:  $-P\ 2_1/n$

$a = 11.6376\ (5)\ \text{\AA}$

$b = 14.6648\ (6)\ \text{\AA}$

$c = 13.9100\ (6)\ \text{\AA}$

$\beta = 96.024\ (1)^\circ$

$V = 2360.81\ (17)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1104$

$D_x = 1.528\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 5784 reflections

$\theta = 2.0\text{--}28.1^\circ$

$\mu = 0.76\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Block, colourless

$0.35 \times 0.30 \times 0.30\ \text{mm}$

*Data collection*

Bruker Kappa APEXII CCD diffractometer	39609 measured reflections
Radiation source: fine-focus sealed tube	5784 independent reflections
Graphite monochromator	3922 reflections with $I > 2\sigma(I)$
$\omega$ and $\varphi$ scans	$R_{\text{int}} = 0.032$
Absorption correction: multi-scan (SADABS; Sheldrick, 2004)	$\theta_{\text{max}} = 28.3^\circ$ , $\theta_{\text{min}} = 2.0^\circ$
$T_{\text{min}} = 0.777$ , $T_{\text{max}} = 0.805$	$h = -15 \rightarrow 12$
	$k = -19 \rightarrow 19$
	$l = -16 \rightarrow 18$

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.046$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.129$	$w = 1/[\sigma^2(F_o^2) + (0.0512P)^2 + 1.4785P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
5784 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
297 parameters	$\Delta\rho_{\text{max}} = 0.66 \text{ e } \text{\AA}^{-3}$
6 restraints	$\Delta\rho_{\text{min}} = -0.64 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3813 (2)	0.07767 (18)	0.03971 (18)	0.0507 (6)
C2	0.3271 (3)	0.0212 (2)	0.1018 (2)	0.0713 (8)
H2	0.3343	0.0326	0.1679	0.086*
C3	0.2640 (3)	-0.0500 (3)	0.0649 (3)	0.0852 (10)
H3	0.2284	-0.0880	0.1063	0.102*
C4	0.2509 (3)	-0.0681 (2)	-0.0346 (3)	0.0751 (9)
H4	0.2071	-0.1176	-0.0588	0.090*
C5	0.3029 (2)	-0.01251 (17)	-0.0952 (2)	0.0549 (6)
C6	0.2932 (3)	-0.0214 (2)	-0.2021 (2)	0.0737 (9)
H6A	0.2729	0.0366	-0.2311	0.111*
H6B	0.2345	-0.0653	-0.2226	0.111*
H6C	0.3658	-0.0412	-0.2216	0.111*
C7	0.49848 (19)	0.20079 (16)	0.35230 (16)	0.0429 (5)
C8	0.5578 (2)	0.29325 (16)	0.33022 (17)	0.0464 (5)
C9	0.3521 (2)	-0.03512 (16)	0.41212 (17)	0.0476 (5)
C10	0.2838 (3)	-0.1102 (2)	0.4323 (2)	0.0663 (8)

H10	0.2952	-0.1667	0.4044	0.080*
C11	0.2013 (3)	-0.0996 (2)	0.4928 (2)	0.0753 (9)
H11	0.1557	-0.1492	0.5060	0.090*
C12	0.1831 (3)	-0.0162 (2)	0.5356 (2)	0.0710 (9)
H12	0.1254	-0.0100	0.5766	0.085*
C13	0.2502 (2)	0.0565 (2)	0.51737 (16)	0.0544 (6)
C14	0.2420 (3)	0.1494 (2)	0.5583 (2)	0.0767 (9)
H14A	0.3029	0.1580	0.6096	0.115*
H14B	0.1686	0.1565	0.5832	0.115*
H14C	0.2490	0.1939	0.5086	0.115*
C15	0.0091 (2)	0.26412 (17)	0.33994 (19)	0.0491 (6)
C16	0.0641 (2)	0.19779 (16)	0.26897 (18)	0.0460 (5)
N1	0.36821 (17)	0.05787 (13)	-0.05609 (15)	0.0459 (4)
N3	0.33292 (17)	0.04379 (13)	0.45650 (13)	0.0437 (4)
N2	0.4457 (2)	0.14867 (19)	0.06970 (17)	0.0668 (6)
N4	0.4353 (2)	-0.03672 (15)	0.35294 (18)	0.0605 (6)
O1	0.48960 (16)	0.14569 (12)	0.28478 (12)	0.0549 (4)
O2	0.46713 (18)	0.19317 (14)	0.43335 (12)	0.0660 (5)
O3	-0.01711 (18)	0.34018 (12)	0.30739 (14)	0.0637 (5)
O4	0.0005 (2)	0.23281 (15)	0.42110 (15)	0.0791 (6)
C11	0.55673 (9)	0.37373 (6)	0.42317 (6)	0.0896 (3)
C12	0.70242 (6)	0.27091 (5)	0.31008 (6)	0.0710 (2)
C13	0.48451 (7)	0.34145 (5)	0.22382 (6)	0.0752 (2)
C14	0.21459 (6)	0.22009 (6)	0.28120 (6)	0.0697 (2)
C15	0.01042 (7)	0.21709 (6)	0.14738 (5)	0.0719 (2)
C16	0.04405 (9)	0.08231 (5)	0.29620 (7)	0.0861 (3)
H4B	0.463 (2)	0.0160 (11)	0.3358 (19)	0.059 (8)*
H2B	0.474 (2)	0.1848 (16)	0.0269 (17)	0.064 (9)*
H4A	0.447 (3)	-0.0864 (14)	0.320 (2)	0.082 (10)*
H2A	0.455 (3)	0.157 (2)	0.1332 (8)	0.082 (10)*
H1A	0.403 (2)	0.0949 (15)	-0.0955 (16)	0.057 (8)*
H3A	0.379 (2)	0.0908 (13)	0.446 (2)	0.061 (8)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0481 (14)	0.0549 (14)	0.0483 (13)	0.0081 (11)	0.0017 (10)	0.0075 (11)
C2	0.0690 (19)	0.084 (2)	0.0605 (17)	-0.0020 (16)	0.0039 (14)	0.0247 (16)
C3	0.078 (2)	0.083 (2)	0.095 (3)	-0.0068 (19)	0.0106 (19)	0.039 (2)
C4	0.0634 (19)	0.0526 (16)	0.108 (3)	-0.0103 (14)	0.0013 (17)	0.0084 (17)
C5	0.0451 (14)	0.0417 (12)	0.0764 (18)	0.0072 (10)	-0.0007 (12)	-0.0040 (12)
C6	0.076 (2)	0.0650 (18)	0.079 (2)	0.0008 (15)	0.0022 (16)	-0.0237 (16)
C7	0.0384 (11)	0.0478 (12)	0.0423 (12)	-0.0102 (10)	0.0031 (9)	0.0052 (10)
C8	0.0464 (13)	0.0488 (13)	0.0453 (12)	-0.0116 (10)	0.0105 (10)	0.0016 (10)
C9	0.0539 (14)	0.0453 (13)	0.0428 (12)	-0.0052 (11)	0.0019 (10)	0.0071 (10)
C10	0.079 (2)	0.0514 (15)	0.0682 (18)	-0.0218 (14)	0.0046 (15)	0.0071 (13)
C11	0.077 (2)	0.084 (2)	0.0644 (18)	-0.0401 (18)	0.0034 (15)	0.0191 (16)
C12	0.0600 (17)	0.104 (2)	0.0517 (15)	-0.0272 (17)	0.0167 (13)	0.0075 (16)
C13	0.0502 (14)	0.0788 (18)	0.0352 (12)	-0.0098 (13)	0.0087 (10)	0.0016 (12)
C14	0.082 (2)	0.092 (2)	0.0611 (17)	-0.0084 (18)	0.0311 (16)	-0.0186 (17)

C15	0.0401 (12)	0.0493 (14)	0.0595 (15)	-0.0035 (10)	0.0123 (10)	-0.0009 (11)
C16	0.0422 (12)	0.0440 (12)	0.0525 (13)	-0.0031 (10)	0.0074 (10)	0.0012 (10)
N1	0.0439 (11)	0.0427 (10)	0.0513 (11)	0.0070 (9)	0.0057 (9)	0.0027 (9)
N3	0.0475 (11)	0.0494 (11)	0.0349 (9)	-0.0116 (9)	0.0073 (8)	0.0031 (8)
N2	0.0811 (17)	0.0736 (16)	0.0452 (13)	-0.0144 (13)	0.0042 (12)	-0.0005 (12)
N4	0.0733 (16)	0.0448 (12)	0.0675 (14)	-0.0029 (11)	0.0263 (12)	-0.0012 (11)
O1	0.0684 (11)	0.0488 (9)	0.0492 (10)	-0.0155 (8)	0.0144 (8)	-0.0021 (8)
O2	0.0833 (14)	0.0737 (12)	0.0437 (10)	-0.0383 (11)	0.0188 (9)	-0.0022 (9)
O3	0.0745 (13)	0.0496 (10)	0.0696 (12)	0.0085 (9)	0.0205 (10)	0.0019 (9)
O4	0.1060 (18)	0.0731 (13)	0.0642 (13)	0.0115 (12)	0.0377 (12)	0.0076 (10)
C11	0.1227 (8)	0.0727 (5)	0.0798 (5)	-0.0399 (5)	0.0406 (5)	-0.0284 (4)
C12	0.0438 (4)	0.0737 (5)	0.0975 (6)	-0.0119 (3)	0.0168 (3)	0.0099 (4)
C13	0.0823 (5)	0.0593 (4)	0.0805 (5)	-0.0066 (4)	-0.0083 (4)	0.0246 (4)
C14	0.0392 (3)	0.0947 (6)	0.0764 (5)	0.0005 (3)	0.0112 (3)	-0.0015 (4)
C15	0.0810 (5)	0.0774 (5)	0.0538 (4)	0.0001 (4)	-0.0092 (3)	-0.0080 (3)
C16	0.1113 (7)	0.0446 (4)	0.1068 (7)	-0.0056 (4)	0.0323 (5)	0.0062 (4)

*Geometric parameters (Å, °)*

C1—N2	1.324 (4)	C10—C11	1.350 (5)
C1—N1	1.357 (3)	C10—H10	0.9300
C1—C2	1.394 (4)	C11—C12	1.385 (5)
C2—C3	1.346 (5)	C11—H11	0.9300
C2—H2	0.9300	C12—C13	1.362 (4)
C3—C4	1.401 (5)	C12—H12	0.9300
C3—H3	0.9300	C13—N3	1.360 (3)
C4—C5	1.360 (4)	C13—C14	1.483 (4)
C4—H4	0.9300	C14—H14A	0.9600
C5—N1	1.361 (3)	C14—H14B	0.9600
C5—C6	1.485 (4)	C14—H14C	0.9600
C6—H6A	0.9600	C15—O3	1.230 (3)
C6—H6B	0.9600	C15—O4	1.232 (3)
C6—H6C	0.9600	C15—C16	1.569 (3)
C7—O2	1.226 (3)	C16—C16	1.756 (2)
C7—O1	1.235 (3)	C16—C15	1.763 (3)
C7—C8	1.567 (3)	C16—C14	1.772 (2)
C8—C11	1.752 (2)	N1—H1A	0.898 (10)
C8—C12	1.765 (3)	N3—H3A	0.896 (10)
C8—C13	1.776 (3)	N2—H2B	0.885 (10)
C9—N4	1.335 (3)	N2—H2A	0.886 (10)
C9—N3	1.341 (3)	N4—H4B	0.881 (10)
C9—C10	1.404 (3)	N4—H4A	0.879 (10)
N2—C1—N1	118.6 (2)	C10—C11—C12	121.4 (3)
N2—C1—C2	123.3 (3)	C10—C11—H11	119.3
N1—C1—C2	118.0 (3)	C12—C11—H11	119.3
C3—C2—C1	119.2 (3)	C13—C12—C11	119.7 (3)
C3—C2—H2	120.4	C13—C12—H12	120.2
C1—C2—H2	120.4	C11—C12—H12	120.2
C2—C3—C4	121.5 (3)	N3—C13—C12	117.7 (3)



C2—C3—H3	119.3	N3—C13—C14	116.3 (2)
C4—C3—H3	119.3	C12—C13—C14	126.0 (3)
C5—C4—C3	119.3 (3)	C13—C14—H14A	109.5
C5—C4—H4	120.3	C13—C14—H14B	109.5
C3—C4—H4	120.3	H14A—C14—H14B	109.5
C4—C5—N1	118.2 (3)	C13—C14—H14C	109.5
C4—C5—C6	125.2 (3)	H14A—C14—H14C	109.5
N1—C5—C6	116.6 (2)	H14B—C14—H14C	109.5
C5—C6—H6A	109.5	O3—C15—O4	129.4 (2)
C5—C6—H6B	109.5	O3—C15—C16	115.6 (2)
H6A—C6—H6B	109.5	O4—C15—C16	115.0 (2)
C5—C6—H6C	109.5	C15—C16—C16	112.94 (17)
H6A—C6—H6C	109.5	C15—C16—C15	112.14 (17)
H6B—C6—H6C	109.5	C16—C16—C15	108.66 (13)
O2—C7—O1	129.1 (2)	C15—C16—C14	106.82 (16)
O2—C7—C8	116.1 (2)	C16—C16—C14	108.04 (13)
O1—C7—C8	114.87 (19)	C15—C16—C14	108.03 (13)
C7—C8—C11	113.64 (16)	C1—N1—C5	123.7 (2)
C7—C8—C12	108.53 (17)	C1—N1—H1A	117.3 (18)
C11—C8—C12	108.85 (13)	C5—N1—H1A	118.9 (18)
C7—C8—C13	108.95 (16)	C9—N3—C13	124.6 (2)
C11—C8—C13	107.87 (14)	C9—N3—H3A	117.3 (18)
C12—C8—C13	108.92 (12)	C13—N3—H3A	118.1 (18)
N4—C9—N3	117.7 (2)	C1—N2—H2B	120 (2)
N4—C9—C10	124.9 (3)	C1—N2—H2A	115 (2)
N3—C9—C10	117.4 (2)	H2B—N2—H2A	125 (3)
C11—C10—C9	119.2 (3)	C9—N4—H4B	117.6 (19)
C11—C10—H10	120.4	C9—N4—H4A	120 (2)
C9—C10—H10	120.4	H4B—N4—H4A	120 (3)
N2—C1—C2—C3	-179.4 (3)	C11—C12—C13—N3	0.3 (4)
N1—C1—C2—C3	-0.2 (4)	C11—C12—C13—C14	-179.4 (3)
C1—C2—C3—C4	-0.6 (5)	O3—C15—C16—C16	-154.9 (2)
C2—C3—C4—C5	0.0 (5)	O4—C15—C16—C16	26.5 (3)
C3—C4—C5—N1	1.4 (4)	O3—C15—C16—C15	-31.7 (3)
C3—C4—C5—C6	-177.7 (3)	O4—C15—C16—C15	149.7 (2)
O2—C7—C8—C11	-5.5 (3)	O3—C15—C16—C14	86.5 (2)
O1—C7—C8—C11	174.28 (18)	O4—C15—C16—C14	-92.2 (2)
O2—C7—C8—C12	115.8 (2)	N2—C1—N1—C5	-179.0 (2)
O1—C7—C8—C12	-64.5 (2)	C2—C1—N1—C5	1.7 (4)
O2—C7—C8—C13	-125.8 (2)	C4—C5—N1—C1	-2.3 (4)
O1—C7—C8—C13	54.0 (3)	C6—C5—N1—C1	176.9 (2)
N4—C9—C10—C11	-179.3 (3)	N4—C9—N3—C13	178.8 (2)
N3—C9—C10—C11	1.5 (4)	C10—C9—N3—C13	-1.9 (4)
C9—C10—C11—C12	-0.3 (5)	C12—C13—N3—C9	1.0 (4)
C10—C11—C12—C13	-0.6 (5)	C14—C13—N3—C9	-179.3 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N1—H1A $\cdots$ O3 <sup>i</sup>	0.90 (1)	1.96 (1)	2.856 (3)	172 (3)
N2—H2B $\cdots$ O4 <sup>i</sup>	0.89 (1)	1.95 (1)	2.825 (3)	167 (3)
N2—H2A $\cdots$ O1	0.89 (1)	2.11 (1)	2.982 (3)	167 (3)
N3—H3A $\cdots$ O2	0.90 (1)	1.84 (1)	2.729 (2)	175 (3)
N4—H4B $\cdots$ O1	0.88 (1)	2.06 (1)	2.929 (3)	167 (3)
N4—H4A $\cdots$ O3 <sup>ii</sup>	0.88 (1)	2.29 (2)	3.096 (3)	152 (3)
C6—H6C $\cdots$ O1 <sup>iii</sup>	0.96	2.50	3.413 (4)	158
C11—H11 $\cdots$ O4 <sup>iv</sup>	0.93	2.50	3.371 (4)	157

Symmetry codes: (i)  $x+1/2, -y+1/2, z-1/2$ ; (ii)  $-x+1/2, y-1/2, -z+1/2$ ; (iii)  $-x+1, -y, -z$ ; (iv)  $-x, -y, -z+1$ .