



POSTER PRESENTATION

Open Access

Comparison of the depth of vaccine-elicited HIV-1 Env epitope-specific CD8+ T lymphocyte responses

SL Hulot^{1*}, BT Korber², G Pantaleo³, J Tartaglia⁴, B Jacobs⁵, B Perdiguero⁶, CE Gomez⁶, M Esteban⁶, N Letvin¹, MS Seaman¹, B Haynes⁷, S Santra¹

From AIDS Vaccine 2012
Boston, MA, USA. 9-12 September 2012

Background

One of the major challenges in the development of an effective HIV-1 vaccine is the extraordinary genetic diversity of the virus. Immunizations of nonhuman primates using consensus and mosaic immunogens have been shown to elicit cross-reactive CD8+T lymphocyte responses that increase the depth of epitope recognition. However, one of the limitations of vaccine-induced epitope-specific CD8+T lymphocytes includes lack of protection against diverse strains and emergent forms of HIV-1 due to altered T cell receptor (TCR) affinity for variant peptide:MHC class I complexes.

Methods

In this study, we immunized a cohort of fifteen Mamu-A*01+ rhesus monkeys with either a 3-valent mosaic Env, group M consensus Env, or single clade B Env vaccine and compared the ability of the CD8+T lymphocyte populations elicited by each immunogen to recognize variants of an HIV-1 envelope epitope sequence p41A (YI9). We identified vaccine-induced CD8+T lymphocytes populations using tetramers constructed with 9 variants of p41A epitope. We assessed the ability of those variant peptides to activate CTL by measuring cytokine production and CD107a expression. We evaluated proliferation using carboxyfluorescein succinimidyl ester. Finally, we investigated the functional avidity of these CD8+T lymphocytes for the variant peptide:Mamu-A*01 complexes using surface plasmon resonance technology.

Results

Our data show that Env immunizations can generate cross-reactive CD8+T lymphocytes that recognize 2 of 9 (22%) of the variants of p41A epitope, with higher responses induced by the consensus and the 3-valent mosaic immunogens (variant from clade C and variant from clade A/E) compared to the single clade Env immunogen. Tetramer-binding data also show that CD8+T lymphocytes from monkeys immunized with mosaic immunogen have a trend of higher binding to majority of the variant peptides tested.

Conclusion

This underscores the potential of mosaic immunogens for generating cellular immune responses with greater depth.

Author details

¹Beth Israel Deaconess Medical Center, Boston, MA, USA. ²Los Alamos National Laboratory, Los Alamos, NM, USA. ³Centre Hospitalier Universitaire Vaudois (CHUV), Lausanne, Switzerland. ⁴Sanofi Pasteur, Toronto, Canada. ⁵Center for Infectious Diseases and Vaccinology, Tempe, AZ, USA. ⁶Centro Nacional de Biotecnología, Madrid, Spain. ⁷Duke University Medical Center / Duke Human Vaccine Institute, Durham, NC, USA.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-P288

Cite this article as: Hulot *et al.*: Comparison of the depth of vaccine-elicited HIV-1 Env epitope-specific CD8+ T lymphocyte responses. *Retrovirology* 2012 **9**(Suppl 2):P288.

¹Beth Israel Deaconess Medical Center, Boston, MA, USA
Full list of author information is available at the end of the article