




Taibah University  
Journal of Taibah University Medical Sciences

www.sciencedirect.com



Review Article

## Protein-based salivary biomarkers for the diagnosis of periodontal diseases: Systematic review and meta-analysis



Esteban Arroyo, PhD<sup>a</sup>, Mónica G. Oliveira-Alves, PhD<sup>b,c</sup>,  
Cintia M. Chamorro-Petronacci, PhD<sup>d,e</sup>, Xabier Marichalar-Mendia, PhD<sup>f,g,\*</sup>,  
Susana B. Bravo-López, PhD<sup>h</sup>, Juan Blanco-Carrión, PhD<sup>i,j</sup> and  
Mario Pérez-Sayáns, PhD<sup>d,e</sup>

<sup>a</sup> Department of Diagnosis and Surgery, Araraquara, School of Dentistry, Sao Paulo State University (Unesp), Araraquara, SP, Brazil

<sup>b</sup> Technology Research Center (NPT), Universidade Mogi das Cruzes, Mogi das Cruzes, Brazil

<sup>c</sup> School of Medicine, Anhembi Morumbi University, Sao José dos Campos, Brazil

<sup>d</sup> Oral Medicine, Oral Surgery and Implantology Unit (MedOralRes), Faculty of Dentistry, University of Santiago de Compostela, Spain

<sup>e</sup> ORALRES Group Instituto de Investigación de Santiago (IDIS), Spain

<sup>f</sup> Grupo GIU21/042, Department of Nursing I, University of the Basque Country (UPV/EHU), Barrio Sarriena s/n, Leioa, Bizkaia, Spain

<sup>g</sup> BioCruces-Bizkaia Health Research Institute, Barakaldo, Spain

<sup>h</sup> Proteomic Platform, Health Research Institute, Santiago de Compostela, A Coruña, Spain

<sup>i</sup> Periodontology Unit, School of Medicine and Dentistry, University of Santiago de Compostela, Santiago de Compostela, Spain

<sup>j</sup> Odontología Médico-Quirúrgica (OMEQUI) Research Group, Health Research Institute of Santiago de Compostela (IDIS), Santiago de Compostela, Spain

Received 6 July 2022; revised 20 October 2022; accepted 9 December 2022; Available online 20 December 2022

### المخلص

**أهداف البحث:** هدفت هذه المراجعة المنهجية والتحليل التلوي إلى تحديد نقاط التعبير التفاضلي للمؤشرات الحيوية القائمة على البروتين والتي يمكن اكتشافها في اللعاب لأمراض اللثة الرئيسية من خلال مراجعة منهجية وتحليل تلوي.

**طريقة البحث:** أجريت مراجعة الأدبيات حتى 31 يناير 2022. تم تقييم الجودة المنهجية وخطر التحيز باستخدام مقياس نيوكاسل أوناناو لدراسات الحالات والشواهد. تم إجراء التحليل الإحصائي للنتائج باستخدام أداة "ميتا المفتوحة" (محلل). تم إجراء تحليل المسارات البيولوجية داخل علامات البروتين التفاضلية باستخدام أداة تحليل تكامل "ستيتش"، اقتصر على التفاعلات مع مستويات ثقة عالية (0.7).

**النتائج:** من بين جميع المؤشرات الحيوية القائمة على البروتين التي تم اكتشافها، يمكن تحليل 12 مؤشرا تلويًا: انترليوكين-1بيتا، والبروتينات الالتهابية الضامة-1 ألفا، والألبومين، وعامل نخر الورم-ألفا، والتيلوبيبتيد المتقاطع من الكولاجين ألفا-1، والغلوبيولين المناعي أ، واللاكتوفيرين، وبروتيناز فليزي شبكي-8، وانترليوكين-6، وانترليوكين-8، وانترليوكين-17 وبروستاغلاندين-2. كانت العلامات اللعابية ذات التطبيق الرئيسي هي انترليوكين-1بيتا، للتمييز بين مجموعة أمراض اللثة المزمنة والتهاب اللثة مع التعبير الكلي 73.5 بيكوجرام/ملي، والتيلوبيبتيد المتقاطع من الكولاجين ألفا-1 لأمراض اللثة المزمنة مقابل مرضى التحكم الصحي مع التعبير الكلي 0,091 نانوجرام/ملي، وبروستاغلاندين-2 لأمراض اللثة المزمنة مقابل مرضى التحكم الأصحاء الذين لديهم تعبير الكلي 36,3 بيكوجرام/ملي.

**الاستنتاجات:** المؤشرات الحيوية ذات أعلى تعبير تفاضلي وأكبر إمكانات للتطبيق السريري هي انترليوكين-1بيتا للتمييز بين التهاب اللثة والمرحلة المتقدمة من التهاب اللثة، والتيلوبيبتيد المتقاطع من الكولاجين ألفا-1 والبروستاغلاندين-2 لالتهاب اللثة مقابل المرضى الأصحاء.

**الكلمات المفتاحية:** المؤشرات الحيوية؛ التهاب اللثة؛ أمراض اللثة؛ المرحلة المتقدمة من التهاب اللثة

\* Corresponding address: Department of Nursing I, University of the Basque Country (UPV/EHU), Barrio Sarriena s/n, 48940, Leioa, Bizkaia, Spain.

E-mail: [xabier.marichalar@ehu.eus](mailto:xabier.marichalar@ehu.eus) (X. Marichalar-Mendia)

Peer review under responsibility of Taibah University.



Production and hosting by Elsevier

## Abstract

**Objective:** This systematic review and meta-analysis was aimed at determining differentially expressed protein-based biomarkers detectable in the saliva for the diagnosis of major periodontal diseases.

**Methods:** A literature review was conducted through January 31, 2022. The methodological quality and risk of bias were assessed with the Newcastle–Ottawa scale for case-control studies. Heterogeneity among studies was analysed with the Q statistical test and the  $I^2$  test.  $p$ -values lower than 0.10 and  $I^2$  values higher than 50% indicated high heterogeneity among studies; therefore, the random-effects model was used. The analysis of biological pathways associated with the differentially expressed protein markers was performed with the STITCH integration analysis tool and was limited to interactions with high confidence levels (0.7).

**Results:** Of all protein-based biomarkers detected, 12 were suitable for meta-analysis: IL-1 $\beta$ , MIP-1 $\alpha$ , albumin, TNF- $\alpha$ , ICTP, Ig-A, lactoferrin, MMP-8, IL-6, IL-8, IL-17 and PGE2. The salivary markers with high applicability were IL-1 $\beta$  for differentiating patients with chronic periodontal disease from patients with gingivitis with an OE = 73.5 pg/mL; ICTP for differentiating patients with chronic periodontal disease from healthy control patients with an OE = 0.091 ng/mL; and PGE2 for differentiating patients with chronic periodontal disease from healthy control patients with an OE = 36.3 pg/mL.

**Conclusions:** The biomarkers with the highest differential expression and the greatest potential for clinical applicability are IL-1 $\beta$  for differentiating periodontitis from gingivitis, and ICTP and PGE2 for differentiating periodontitis from healthy status.

**Keywords:** Biomarkers; Gingivitis; Periodontal diseases; Periodontitis

© 2022 The Authors.

Production and hosting by Elsevier Ltd on behalf of Taibah University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Gingivitis can be diagnosed according to specific signs of inflammation on the gums in response to the biofilm accumulated on the tooth surface over several days. Various factors, such as the systemic conditions of the patient or the intake of specific drugs, may favour development of this disease.<sup>1</sup> The pathology of gingivitis is characterised on the basis of clinical reversibility and the presence of a moderate to high bacterial load. In addition, gingivitis is understood as a precursor of periodontal disease and therefore must be controlled to prevent irreversible outcomes.<sup>2</sup>

Periodontal disease is a chronic inflammatory disease of the periodontium, which results in loss of the periodontal ligament and the destruction of the underlying alveolar bone.<sup>3</sup> The host's immune system defends against bacteraemia through an inflammatory response, which results in chronic inflammatory infection. Additionally, the immune response modulates the expression and virulence of biofilms, which interfere with multiple protective homeostatic mechanisms.<sup>4</sup>

Periodontal diseases affect as much as 50% of the world's population and rank sixth among the most prevalent pathologies worldwide.<sup>5</sup> Currently, the diagnosis of periodontitis is a precise, consistent and simple process enabling identification of disease stage, and the standardisation of specific treatments.<sup>6</sup> However, these clinical and radiological methods are useful only for confirming the presence and severity of this disease, whereas information on the biological behaviour at the stage of diagnosis is lacking. Moreover, clinical monitoring is a time-consuming activity in daily practice.<sup>7</sup>

Saliva is a known reservoir of large amounts of molecules and may potentially be influenced by bacterial dysbiosis and simultaneous effects of the host's immune system response. Many of these molecules are proteins that are involved in various molecular processes and can be quantified with diverse molecular techniques to identify diagnostic cut-offs for periodontal pathologies.<sup>8</sup>

This systematic review and meta-analysis was aimed at exploring the available literature to identify differentially expressed protein-based biomarkers that can be detected in saliva and used for diagnosis of major periodontal diseases.

## Materials and Methods

### *Protocols and inclusion criteria*

A literature review was conducted through January 31, 2022. The review was organised according to the PRISMA guidelines<sup>9</sup> for systematic reviews. The PICO elements were as follows: P: patients with gingivitis (GV), chronic periodontal disease (CPD) or aggressive periodontal disease (APD); I: salivary protein study; C: healthy control patients (HCG); O: differences in salivary protein expression between CPD and GV, compared with HCG.

The inclusion criteria were as follows: 1) human case-control studies in English; 2) studies including quantified salivary protein-based markers; and 3) studies using ELISA and/or equivalent analytic methods with quantitative expression determination (bromocresol green method, Immunoturbidimetry and LUMINEX).

The exclusion criteria were as follows: 1) studies performing differential expression proteomic analysis through mass spectrometry or LC-MS/MS; 2) *in vitro* or animal studies; 3) studies quantifying protein expression in units of measurement not transformable to ng/dl; 4) studies in which the marker was expressed below a minimum cut-off point; 5) studies with unclear or poorly defined control groups; and 6) studies with a risk of bias rating of less than 7 on the Newcastle–Ottawa Scale.<sup>10</sup>

## Sources

The electronic research was conducted in MEDLINE through PubMed; EMBASE through OVID; Web of Science; Scopus; Cochrane Library; Clinical Trials; the five regional bibliographic databases of the WHO; and databases from the Conference Citation Index.<sup>11</sup> The terms used in this process were as follows: gingivitis OR chronic periodontitis OR periodontal disease OR aggressive periodontal disease AND salivary protein OR biomarker OR salivary biomarker OR protein expression.

## Study selection and data extraction process

The research was performed by five observers (EAA, MGOA, SBBL, JBC and MPS), and CMCP acted as an independent observer in the event of any discrepancies. First, the title and abstract of the retrieved articles were read. Subsequently, the full text of all eligible studies was examined, and a decision was made regarding study inclusion. The data were extracted with a standardised, pilot-tested form.

The form included the following components: title; authors; publication year; study group; sample size; biomarker; quantitative expression; analysis method; and diagnostic criteria for the pathology under study.

## Diagnostic criteria

The GV definition was established according to Armitage's 1999 classification defining GV as gingival disease induced by dental plaque without any other contributing factor.<sup>12</sup> The diagnosis of CPD varied from radiographic to arbitrary criteria based on probing depth thresholds (PPD) or clinical attachment loss (CAL) for a determined number of affected sites, or PPD or CAL averages for the full mouth. Because of modifications that have occurred over the years, broad criteria were established as follows: CPD Group: Two or more sites with PPD  $\geq 4$  mm and CAL  $\geq 2$  mm. APD group: any PPD  $\geq 6$  mm with associated CAL  $> 2$  mm. GV group:  $\geq 20\%$  bleeding on probing, with  $< 3\%$  of sites with PPD  $\geq 4$  mm and no sites with CAL  $\geq 2$  mm. HCG group: no systemic diseases, no immunosuppressive medication,  $< 20\%$  of sites bleeding on probing,  $< 3\%$  of sites with a probing depth  $\geq 4$  mm and no sites with clinical insertion loss  $\geq 2$  mm.<sup>12</sup>

## Risk of bias assessment, data synthesis and analysis

The methodological quality and risk of bias were assessed with the Newcastle–Ottawa scale for case-control studies.<sup>10</sup> Two researchers (MPS and EAA) performed this analysis independently, and in the event of any discrepancies, a third party (XMM) served as a mediator. Studies with a score greater than 7 were included.

## Qualitative analysis

The expression of each protein biomarker in saliva was analysed, in conjunction with the author and year of

publication. The expression estimates are shown as integer values with standard deviation for each biomarker and subgroup. The weighted mean difference between groups was obtained for each model, thus yielding the overall expression (OE) and 95% confidence interval (CI 95%).

## Quantitative analysis

A meta-analysis was performed when at least two articles provided data on the same biomarker and units of measurement. Our analysis strategy was to include all expression studies regardless of their threshold values. The statistical analysis of the results was performed with the OpenMeta tool (Analyst).

## Functional networks and pathway mapping

The analysis of biological pathways associated with the differentially expressed protein markers was performed with the STITCH integration analysis tool and was limited to interactions with high confidence levels (0.7)<sup>13 (P4)</sup>.

## Assessment of heterogeneity


First, we evaluated the threshold effect of expression graphically by examining the forest plot; second, we statistically evaluated the threshold effect with the Q test to determine the  $I^2$  index. A p-value  $> 0.1$  and an index  $I^2 < 50\%$  indicated low heterogeneity among studies. In contrast, a p-value of  $< 0.1$  and an index  $I^2 > 50\%$  indicated the presence of considerable heterogeneity. Study specific estimates were combined with random effects models.

## Results

A total of 81 studies were included in the qualitative assessment, of which 26 were suitable for quantitative analysis<sup>14–39</sup> (Figure 1). Of all the detected protein-based biomarkers (Supplementary 1), 12 were suitable for meta-analysis (Table 1).

According to the Newcastle–Ottawa Scale, 2 (6.7%) were determined to be of low quality, 20 (66.7%) were determined to be of moderate quality, and 8 (26.6%) were determined to be of high quality (Supplementary 2).

The biomarkers included were IL-1 $\beta$ , MIP-1 $\alpha$ , albumin, TNF- $\alpha$ , ICTP, Ig-A, lactoferrin, MMP-8, IL-6, IL-8, IL-17 and PGE2. Forest plots were drawn for the different subgroups, CPD vs. GV (Figure 2), CPD vs. HCG (Figure 3) APD vs. HCG (Figure 4) and GV vs. HCG (Figure 5). The most studied biomarker was IL-1 $\beta$ . All markers were positive and overexpressed in the CPD group, in contrast to the GV and HCG groups, except for albumin (CPD vs. HCG; OE =  $-16.8$  mg/mL; CI 95%:  $-50.3$  to  $16.6$  mg/mL,  $p = 0.324$ ;  $I^2 = 98.02\%$ ,  $p < 0.001$ ) and Ig-A (CPD vs. HCG; OE =  $-47.9$  mg/mL; CI 95%:  $-223.9$  to  $127.9$  mg/mL,  $p = 0.593$ ;  $I^2 = 95.59\%$ ,  $p < 0.001$ ). MMP-8 was the most overexpressed marker in the CPD vs. HCG group, although it had a high degree of heterogeneity (OE =  $319.4$  ng/mL; CI 95%:  $206.1$ – $432.8$  ng/mL,  $p < 0.001$ ;  $I^2 = 92.20\%$ ,  $p < 0.001$ ).





**Figure 1:** Flowchart of the systematic review.

**Table 1: Complete description of the overall expression values (OE) of the different salivary biomarkers according to the subgroup under study. CPD (chronic periodontal disease), GV (gingivitis), APD (aggressive periodontitis), HCG (healthy control), CI (confidence interval).**

Biomarker	Subgroup	OE	IC 95% (p)	Heterogeneity ( $I^2$ , p-value)
IL-1B pg/mL	CPD/GV	73.598	70.660–76.536 (p < 0.001)	( $I^2$ = 0%, p = 0.961)
	CPD/HCG	93.801	52.393–135.208 (p < 0.001)	( $I^2$ = 99.30%, p < 0.001)
	CPD-AGR/HCG	16.628	–1.265 to 34.521 (p = 0.069)	( $I^2$ = 0%, p = 0.775)
	GV/HCG	62.421	–41.856 to 166.698 (p = 0.241)	( $I^2$ = 90.38%, p = 0.001)
MIP-1 $\alpha$ pg/mL	CPD/GV	10.131	–1.491 to 21.754 (p = 0.088)	( $I^2$ = 99.79%, p < 0.001)
	CPD/HCG	20.077	6.200–33.953 (p = 0.005)	( $I^2$ = 99.89%, p < 0.001)
	GV/HCG	9.954	7.700–12.208 (p < 0.001)	( $I^2$ = 95.48%, p < 0.001)
Albumin mg/mL	CPD/HCG	–16.867	–50.366 to 16.632 (p = 0.324)	( $I^2$ = 98.02%, p < 0.001)
TNF- $\alpha$ pg/mL	CPD/HCG	1.236	0.373–2.099 (p = 0.005)	( $I^2$ = 93.57%, p < 0.001)
ICTP ng/mL	CPD/HCG	0.091	0.019–0.164 (p = 0.013)	( $I^2$ = 16.22%, p = 0.275)
Ig-A mg/mL	CPD/HCG	25.755	–84.664 to 136.174 (p = 0.593)	( $I^2$ = 88.31%, p < 0.001)
Lactoferrin mg/mL	CPD/HCG	10.449	–7.580 to 28.477 (p = 0.256)	( $I^2$ = 95.91%, p < 0.001)
MMP-8 ng/mL	CPD/HCG	319.489	206.139–432.838 (p < 0.001)	( $I^2$ = 92.20%, p < 0.001)
IL-6 pg/mL	CPD/HCG	49.166	18.380–79.952 (p = 0.002)	( $I^2$ = 99.93%, p < 0.001)
IL-8 pg/mL	CPD/HCG	43.563	–330.540 to 417.667 (p = 0.819)	( $I^2$ = 71.59%, p = 0.061)
IL-17 pg/mL	CPD/HCG	1.843	–3.086 to 6.772 (p = 0.464)	( $I^2$ = 98.82%, p < 0.001)
PGE2 pg/mL	CPD/HCG	36.305	18.350–54.260 (p < 0.001)	( $I^2$ = 0%, p = 0.850)

## CPD vs GV

## IL-1B

MIP-1 $\alpha$ 

**Figure 2:** Forest plots of the chronic periodontitis (CPD) subgroups vs. the gingivitis (GV) group, and the IL-1 $\beta$  and MIP-1 $\alpha$  biomarkers.



CPD vs HCG




Figure 3: Forest plots of the chronic periodontitis (CPD) subgroups vs. the healthy control (HCG) group and the IL-1β, MIP-1α, albumin, TNF-α, ICTP, Ig-A, lactoferrin, MMP-8, IL-6, IL-8, IL-17 and PGE2 biomarkers.

APD vs HCG

IL-1B





Figure 4: Forest plot of the aggressive periodontal disease (APD) subgroup vs. the healthy control (HCG) group, and the IL-1β biomarker.



**Figure 5:** Forest plots of the gingivitis (GV) subgroup vs. the healthy control (HCG) group, and the IL-1 $\beta$  and MIP-1 $\alpha$  biomarkers.

On the basis of the significance between groups and the low degree of heterogeneity of the results, the salivary markers found to have strong applicability were IL-1 $\beta$  for differentiating the I and GV groups with an OE = 73.5 pg/mL (CI 95%: 70.6–76.5 pg/mL,  $p < 0.001$ ;  $I^2 = 0\%$ ,  $p = 0.961$ ); ICTP for differentiating CPD from HCG with an OE = 0.091 ng/mL (CI 95%: 0.019–0.164 ng/mL,  $p = 0.013$ ;  $I^2 = 16.22\%$ ,  $p = 0.275$ ); and PGE2 for differentiating CPD from HCG with an OE = 36.3 pg/mL (CI 95%: 18.3–54.2 pg/mL,  $p < 0.001$ ;  $I^2 = 0\%$ ,  $p = 0.850$ ). However, the number of studies included in the meta-analysis was highly low for these markers (Supplementary 3).

We conducted a functional interaction study for these three markers and observed a strong network between IL-1 $\beta$  and PGE2, whereas ICTP remained outside this node (Supplementary 4).

## Discussion

This study was aimed at determining differentially expressed salivary proteins for diagnosis of the main periodontal diseases, CPD and GV. Despite analysing 81 studies, we were able to perform statistical comparisons of protein expression through meta-analysis on 26 studies on 12 previously described biomarkers.

IL-1 $\beta$ , a cytokine produced by macrophages, is a key mediator of the inflammatory response and is stimulated in the presence of bacterial components associated with the osseous reabsorption process in periodontal diseases. Furthermore, clinical studies have indicated that high levels of IL-1 $\beta$  are associated with gingival inflammation, the severity of periodontitis and periodontal disease progression.<sup>40</sup> In the present study, the IL-1 $\beta$  levels were significantly higher in the CPD group with respect to the GV and HCG groups. The APD and GV groups presented lower values of OE than the CPD group, and no significant

differences were detected with respect to the HCG group (Table 1). IL-1 $\beta$  was the marker assessed in the most studies in the meta-analysis (10 for the CPD–HCG comparison,<sup>17,18,20,23,28–30,34,36,37,39</sup> 2 for the CPD–GV comparison,<sup>29,36</sup> 2 for the CPD/AGR–HCG comparison<sup>18,19</sup> and 2 for the GV–HCG comparison<sup>29,36</sup>).

Research conducted by Miller in a case study from the USA reported the highest average expression of IL-1 $\beta$  among all studies included in the meta-analysis, with a level of 540,900 pg/mL (CI 95%: 157,335–924,565) in the CPD and HCG groups.<sup>39</sup> Rangbulla<sup>23</sup> reported an IL-1 $\beta$  expression of 440.9 pg/mL (CI 95%: 317.3–564.4). In contrast, the studies including populations from Medellín, Colombia<sup>18</sup>; Belo Horizonte, Brazil<sup>28</sup>; and Boston, USA<sup>37</sup> showed an IL-1 $\beta$  expression lower than the average in the meta-analysis (Figure 3). The differences in detection for the studies conducted by Miller<sup>39</sup> and Rangbulla<sup>23</sup> compared with the study conducted by Teles,<sup>37</sup> which had the highest and lowest IL-1 $\beta$  expression in the meta-analysis, respectively, might have been due to differences in disease levels/severity, the collection and storage methods, processing and centrifugation, or salivary analysis, as well as potential cytokine degradation. The patients included in the study conducted by Miller<sup>39</sup> presented relatively more severe periodontal disease. This explanation might also apply to the findings of Rangbulla,<sup>23</sup> who selected patients with a profile similar to those included in the study conducted by Miller.<sup>39</sup>

MIP-1 $\alpha$ , a cytokine secreted by a wide variety of cells, contributes to the pathogenesis of a wide range of inflammatory diseases through processes including cell recruitment, stem cell inhibition and immune response maintenance.<sup>35</sup> MIP-1 $\alpha$  is associated with the formation of multinuclear cells, thus indicating its potential role in bone destruction associated with periodontal diseases.<sup>41</sup> In the present work, we observed differences in MIP-1 $\alpha$  expression between CPD–HCG and GV–HCG. However, no significant

differences were observed among pathological conditions. Nonetheless, all comparisons showed high heterogeneity, with  $I^2$  values of approximately 95%. The variability of MIP-1 $\alpha$  expression observed among studies may be associated with the sample sizes, given that in the study by Nisha, the same number of individuals was tested in each group,<sup>35</sup> whereas in the study by Ebersole, the CPD group had more than twice the number of participants as the GV group.<sup>17</sup> The variability might also be associated with the detection methods, given that Nisha used ELISA,<sup>35</sup> and Ebersole used the Luminex 100IS system.<sup>17</sup>

Albumin is the most abundant protein in human blood, and its main role is regulating the osmotic pressure in the blood plasma; it exhibits an esterase-like activity with broad substrate specificity.<sup>42</sup> Novakovic<sup>16</sup> has reported significantly higher levels of albumin in healthy patients, and suggested that non-enzymatic antioxidants may be a promising indicator of periodontal health. In the present study, we observed no significant differences in albumin expression in the CPD–HCG comparison. Furthermore, high heterogeneity was observed ( $I^2 > 95\%$ ). The wide heterogeneity in the results may be explained by use of different analytic methods.

TNF- $\alpha$ , a cytokine secreted predominantly by macrophages, can induce cell death in certain tumour cell lines, stimulate cell proliferation and induce cell differentiation; it also plays a key role in the pathogenesis of periodontal disease.<sup>42</sup> In the present work, we observed significant differences in the CPD–HCG comparison, with high heterogeneity values ( $I^2 = 93.57\%$ ). Ding<sup>43</sup> has reported contradictory results in the association between periodontitis and TNF- $\alpha$ . However, each of the studies included in the meta-analysis observed that the TNF- $\alpha$  expression was higher in CPD than in HCG.

Pyridinoline cross-linked carboxyterminal telopeptide of type-I collagen (ICTP) is a type 1 bone collagen molecule that is released into the circulation after bone resorption and the degradation of the collagen matrix by protease and bacterial collagen. This marker indicates the progression of periodontal collapse, including alveolar bone loss.<sup>44</sup> Al-Shammari<sup>45</sup> has observed high levels of ICTP in gingival crevicular fluid in periodontitis. In the present work, we observed significant differences in ICTP expression in the CPD–HCG comparison, with reasonably low heterogeneity values ( $I^2 = 16.22\%$ ). Among the selected studies, Frodge<sup>21</sup> did not observe any significant differences in ICTP expression or any correlation between ICTP expression and smoking/non-smoking status. In contrast, in the study conducted by Gursoy,<sup>22</sup> a significant difference was observed only between the CPD and HCG when the participants were non-smokers. These differences might have been due to differences in harmful habits or the geographical origin of the study populations, given that Frodge studied an American population,<sup>21</sup> whereas Gursoy studied a Finnish population.<sup>22</sup>

Ig-A is predominant in saliva and it is derived from plasma cells of the salivary glands.<sup>46</sup> Rangulla has highlighted that the decreased Ig-A levels following treatment suggest an association between this biomarker and periodontal disease.<sup>23</sup> In the present work, we observed no significant differences in Ig-A expression in the CPD–HCG comparison. However, significant differences were

observed between CPD and HCG in the studies by Rangulla<sup>23</sup> and Olayanju,<sup>24</sup> whereas Glimvall<sup>25</sup> did not report any differences. These differences might have been because smokers accounted for almost 50% of Glimvall's CPD group, whereas the samples assessed in the other two studies contained only non-smokers. Therefore, given that Ig-A is an inflammatory marker, the vasoconstrictive effects of tobacco have been suggested to affect Ig-A expression.

Lactoferrin, a metalloprotein with antimicrobial, anti-inflammatory and anticancer characteristics, can be found in tears, saliva, sweat, colostrum and milk. It is also found in polymorphonuclear leukocyte granules,<sup>25</sup> Glimvall has correlated the salivary lactoferrin concentration with the progression of periodontal disease.<sup>25</sup> In the present study, we observed no significant differences in lactoferrin expression in the CPD–HCG comparison. However, the study conducted by Lourenço<sup>34</sup> has indicated significantly higher lactoferrin expression in the CPD group than the HCG group, and similar results have been observed by Glimvall.<sup>25</sup> Both studies used the ELISA method to detect lactoferrin, although the study performed by Lourenço<sup>34</sup> in a Brazilian population observed expression seven times greater than those found by Glimvall<sup>25</sup> in a study in a Swiss cohort.

Extracellular matrix metalloproteinase-8 (MMP-8) is involved in extracellular matrix breakdown, and in various physiological and pathological processes. Multiple active forms of the enzyme exist, which act in the degradation of type I–III collagens.<sup>42</sup> Sorsa<sup>47</sup> has observed high MMP-8 levels in gingival crevicular fluid in patients with periodontal disease, which were correlated with periodontitis levels. Gursoy<sup>22</sup> has concluded that gingival inflammation can affect secretion of both the MMP-8 and the matrix metalloproteinase inhibitor (TIMP-1), thus modulating saliva concentrations. In the present work, we observed significant differences in the CPD–HCG comparison, with high heterogeneity ( $I^2 = 92.20\%$ ). Within each study, significantly higher expression of MMP-8 was observed in the CPD group. The study by Rangulla has indicated that the MMP-8 expression in CPD patients significantly decreased after periodontal treatment.<sup>23</sup> Johnson<sup>48</sup> has noted several limitations in the detection of high levels of MMP-8, including the sample size and the cross-sectional nature of the study.

Interleukin-6 (IL-6) is a cytokine that acts on inflammation and B-cell maturation, and it is predominantly produced in areas with high and chronic inflammation, where serum is secreted.<sup>42</sup> We observed significant differences in the CPD–HCG comparison, with high heterogeneity ( $I^2 = 99.93\%$ ). All but two included studies<sup>15,17,29,32,36</sup> showed significantly higher IL-6 expression in CPD.<sup>20,30</sup>

Interleukin-8 (IL-8), a chemoattractant cytokine produced by a variety of tissue and blood cells, attracts and activates neutrophils, which migrate to regions of inflammation and release granulate enzymes,<sup>49</sup> Finoti has observed higher levels of protein in the gingival tissues of patients with chronic periodontal disease in the presence of high expression of the IL-8 gene.<sup>49</sup> In contrast, we found no significant differences in the CPD–HCG comparison, with high heterogeneity ( $I^2 = 71.59\%$ ). Cheng<sup>32</sup> has not observed significant differences in the presence of IL-8 in saliva between the CP and control groups. Likewise, Arellano-Garcia



has not observed any differences in IL-8 expression in the saliva between the CPD and HCG groups.<sup>33</sup> Teles and Cheng<sup>32,37</sup> underscored existing doubts regarding the use of salivary cytokines as biomarkers for periodontal disease. Teles<sup>37</sup> has also highlighted the need for standardisation of disease level in these types of studies. Furthermore, Cheng<sup>32</sup> has noted the importance of using the total number of salivary proteins in unstimulated saliva as a point of reference for standardisation in salivary protein biomarker research.

Interleukin 17 (IL-17), a pro-inflammatory cytokine produced by T lymphocytes in tissues with periodontal disease, is involved in the modulation of T helper 1 (Th1) lymphocytes and the enhancement of inflammatory reactions via gingival fibroblast-derived mediators. It is also involved in osteoclastic bone resorption.<sup>50</sup> In the present systematic review and meta-analysis, we found no significant differences ( $p = 0.464$ ) in IL-17 expression between patients with periodontitis and healthy patients, and extreme heterogeneity in values was observed ( $I^2 = 98.82\%$ ). Özçaka<sup>27</sup> has observed a significant difference in salivary IL-17 concentration between the COMPOUND and HCG groups ( $p = 0.009$ ). Likewise, Yang<sup>14</sup> has reported a higher IL-17 concentration in CPD than HCG, and observed a significant decrease in concentration after treatment of periodontal disease.

Prostaglandin E2 (PGE2) is a product of cyclooxygenase in arachidonic acid metabolism whose functional effects include fever, pain, vasodilation and bone resorption. Cyclooxygenase-2 expression is elevated in inflamed gingival tissues, and this enzyme is responsible for PGE2 production, thus stimulating proinflammatory reactions.<sup>51</sup> In the present systematic review and meta-analysis, we observed significant differences ( $p < 0.001$ ) in PGE2 expression in the CPD–HCG comparison, and there was but no heterogeneity in the data included in the meta-analysis ( $I^2 = 0\%$ ). Kosaka<sup>29</sup> has observed a significant and proportional correlation between PGE2 concentration and the Community Periodontal Index.

## Conclusion

The most studied protein based salivary biomarker was IL-1 $\beta$ . The biomarkers with the highest differential expression and the greatest potential for clinical applicability are IL-1 $\beta$  for differentiating periodontitis from gingivitis; ICTP for differentiating patients with periodontitis from healthy patients; and PGE2 for differentiating patients with periodontitis from healthy patients. The number of studies must be increased in the future to support clinical translation of the findings.

## Source of funding

Esteban Arroyo received funding for this research from Estudantes-Convênio de Pós-Graduação–PEC-PG, CAPES/CNPq, Brazil (88881.154490/2017-01). This study was financed partly by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior–Brasil (CAPES)–Finance Code 001.

## Conflict of interest

The authors have no conflict of interest to declare.

## Ethical approval

Not applicable.

## Authors contributions

EA: Conceptualization, Investigation, Writing - Original Draft, Visualization. MGOA: Methodology, Investigation, Writing - Original Draft, Visualization. CMC: Methodology, Validation, Investigation, Visualization, Supervision. Xabier Marichalar-Mendia: Formal analysis, Investigation, Visualization. SBB: Investigation, Writing - Review & Editing, Visualization. JB: Investigation, Writing - Review & Editing, Visualization. MP: Conceptualization, Methodology, Formal analysis, Investigation, Writing - Review & Editing, Visualization, Supervision, Project administration. All authors have critically reviewed and approved the final draft and are responsible for the content and similarity index of the manuscript.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jtumed.2022.12.004>.

## References

- Mariotti A. Dental plaque-induced gingival diseases. *Ann Periodontol* 1999; 4(1): 7–19. <https://doi.org/10.1902/annals.1999.4.1.7>.
- Lang NP, Bartold PM. Periodontal health. *J Clin Periodontol* 2018; 45(S20): S9–S16. <https://doi.org/10.1111/jcpe.12936>.
- Pihlstrom BL, Michalowicz BS, Johnson NW. Periodontal diseases. *Lancet* 2005; 366(9499): 1809–1820. [https://doi.org/10.1016/S0140-6736\(05\)67728-8](https://doi.org/10.1016/S0140-6736(05)67728-8).
- Meyle J, Chapple I. Molecular aspects of the pathogenesis of periodontitis. *Periodontology* 2015; 69(1): 7–17. <https://doi.org/10.1111/prd.12104>.
- Bernabe E, Marcenes W, Hernandez CR, Bailey J, Abreu LG, Alipour V, et al. Global, regional, and national levels and trends in burden of oral conditions from 1990 to 2017: a systematic analysis for the global burden of disease 2017 study. *J Dent Res* 2020; 99(4): 362–373. <https://doi.org/10.1177/0022034520908533>.
- Tonetti MS, Greenwell H, Kornman KS. Staging and grading of periodontitis: framework and proposal of a new classification and case definition. *J Periodontol* 2018; 89(S1): S159–S172. <https://doi.org/10.1002/JPER.18-0006>.
- Savage A, Eaton KA, Moles DR, Needleman I. A systematic review of definitions of periodontitis and methods that have been used to identify this disease. *J Clin Periodontol* 2009; 36(6): 458–467. <https://doi.org/10.1111/j.1600-051X.2009.01408.x>.
- Lorenzo-Pouso AI, Pérez-Sayáns M, Bravo SB, López-Jornet P, García-Vence M, Alonso-Sampedro M, et al. Protein-based salivary profiles as novel biomarkers for oral diseases. *Dis Markers* 2018; 2018: 1–22. <https://doi.org/10.1155/2018/6141845>.
- Liberati A, Altman D, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP A, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol* 2009; 62: e1–e34. <https://doi.org/10.1016/j.jclinepi.2009.06.006>.

10. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010; 25(9): 603–605. <https://doi.org/10.1007/s10654-010-9491-z>.
11. Khurshid Z, Tariq R, Asiri FY, Abid K, Zafar MS. Literature search strategies in dental education and research. *J Taibah Univ Med Sci* 2021; 16(6): 799–806. <https://doi.org/10.1016/j.jtumed.2021.05.012>.
12. Armitage GC. Development of a classification system for periodontal diseases and conditions. *Ann Periodontol* 1999; 4(1): 1–6. <https://doi.org/10.1902/annals.1999.4.1.1>.
13. Kuhn M, Szklarczyk D, Pletscher-Frankild S, Blicher TH, von Mering C, Jensen LJ, et al. STITCH 4: integration of protein–chemical interactions with user data. *Nucleic Acids Res* 2014; 42(D1): D401–D407. <https://doi.org/10.1093/nar/gkt1207>.
14. Yang X, Li C, Pan Y. The influences of periodontal status and periodontal pathogen quantity on salivary 8-hydroxydeoxyguanosine and interleukin-17 levels. *J Periodontol* 2016; 87(5): 591–600. <https://doi.org/10.1902/jop.2015.150390>.
15. Sharma M, Bairy I, Pai K, Satyamoorthy K, Prasad S, Berkovitz B, et al. Salivary IL-6 levels in oral leukoplakia with dysplasia and its clinical relevance to tobacco habits and periodontitis. *Clin Oral Investig* 2011; 15(5): 705–714. <https://doi.org/10.1007/s00784-010-0435-5>.
16. Novakovic N, Todorovic T, Rakic M, Milinkovic I, Dozic I, Jankovic S, et al. Salivary antioxidants as periodontal biomarkers in evaluation of tissue status and treatment outcome. *J Periodontol Res* 2014; 49(1): 129–136. <https://doi.org/10.1111/jre.12088>.
17. Ebersole JL, Schuster JL, Stevens J, Dawson 3rd D, Kryscio RJ, Lin Y, et al. Patterns of salivary analytes provide diagnostic capacity for distinguishing chronic adult periodontitis from health. *J Clin Immunol* 2013; 33(1): 271–279. <https://doi.org/10.1007/s10875-012-9771-3>.
18. Isaza Guzmán DM, Medina-Piedrahíta VM, Gutiérrez-Henao C, Tobón-Arroyave SI. Salivary levels of NLRP3 inflammasome-related proteins as potential biomarkers of periodontal clinical status. *J Periodontol* 2017; 88(12): 1329–1338. <https://doi.org/10.1902/jop.2017.170244>.
19. Fine DH, Markowitz K, Furgang D, Fairlie K, Ferrandiz J, Nasri C, et al. Macrophage inflammatory protein-1 $\alpha$ : a salivary biomarker of bone loss in a longitudinal cohort study of children at risk for aggressive periodontal disease? *J Periodontol* 2009; 80(1): 106–113. <https://doi.org/10.1902/jop.2009.080296>.
20. Gursoy UK, Könönen E, Uitto V, Pussinen PJ, Hyvärinen K, Suominen-Taipale L, et al. Salivary interleukin-1b concentration and the presence of multiple pathogens in periodontitis. *J Clin Periodontol* 2009; 36(11): 922–927. <https://doi.org/10.1111/j.1600-051X.2009.01480.x>.
21. Frodge BD, Ebersole JL, Kryscio RJ, Thomas MV, Miller CS. Bone remodeling biomarkers of periodontal disease in saliva. *J Periodontol* 2008; 79(10): 1913–1919. <https://doi.org/10.1902/jop.2008.080070>.
22. Gursoy UK, Könönen E, Pradhan-Palikhe P, Tervahartiala T, Pussinen PJ, Suominen-Taipale L, et al. Salivary MMP-8, TIMP-1, and ICTP as markers of advanced periodontitis: MMP-8 in periodontitis. *J Clin Periodontol* 2010; 37(6): 487–493. <https://doi.org/10.1111/j.1600-051X.2010.01563.x>.
23. Rangbulla V, Nirola A, Gupta M, Batra P, Gupta M. Salivary IgA, interleukin-1B and MMP-8 as salivary biomarkers in chronic periodontitis patients 2017; 20(1): 9.
24. Olayanju O, Rahamon S, Joseph I, Arinola O. Salivary immunoglobulin classes in Nigerians with periodontitis. *J Contemp Dent Pract* 2012; 13(2): 163–166. <https://doi.org/10.5005/jp-journals-10024-1114>.
25. Glimvall P, Wickström C, Jansson H. Elevated levels of salivary lactoferrin, a marker for chronic periodontitis?: salivary lactoferrin and chronic periodontitis. *J Periodontol Res* 2012; 47(5): 655–660. <https://doi.org/10.1111/j.1600-0765.2012.01479.x>.
26. Chauhan A, Yadav SS, Dwivedi P, Lal N, Usman K, Khattri S. Correlation of serum and salivary cytokines level with clinical parameters in metabolic syndrome with periodontitis: role of salivary cytokines in periodontitis. *J Clin Lab Anal* 2016; 30(5): 649–655. <https://doi.org/10.1002/jcla.21917>.
27. Özçaka Ö, Nalbantsoy A, Buduneli N. Interleukin-17 and interleukin-18 levels in saliva and plasma of patients with chronic periodontitis: interleukin-17, interleukin-18 and chronic periodontitis. *J Periodontol Res June* 2011. <https://doi.org/10.1111/j.1600-0765.2011.01377.x>, no–no.
28. Moura MF, Navarro TP, Silva TA, Costa LOM, Soares Dutra Oliveira AM, Costa FO. Periodontitis and endothelial dysfunction: periodontal clinical parameters and levels of salivary markers interleukin-1 $\beta$ , tumor necrosis factor- $\alpha$ , matrix metalloproteinase-2, tissue inhibitor of metalloproteinases-2 complex, and nitric oxide. *J Periodontol* 2017; 88(8): 778–787. <https://doi.org/10.1902/jop.2017.170023>.
29. Kosaka T, Kokubo Y, Ono T, Sekine S, Kida M, Kikui M, et al. Salivary inflammatory cytokines may be novel markers of carotid atherosclerosis in a Japanese general population: the Suita study. *Atherosclerosis* 2014; 237(1): 123–128. <https://doi.org/10.1016/j.atherosclerosis.2014.08.046>.
30. Rathnayake N, Akerman S, Klinge B, Lundegren N, Jansson H, Tryselius Y, et al. Salivary biomarkers of oral health – a cross-sectional study. *J Clin Periodontol* 2013; 40(2): 140–147. <https://doi.org/10.1111/jcpe.12038>.
31. Gursoy UK, Könönen E, Pussinen PJ, Tervahartiala T, Hyvärinen K, Suominen AL, et al. Use of host- and bacteria-derived salivary markers in detection of periodontitis: a cumulative approach. *Dis Markers* 2011; 30(6): 299–305. <https://doi.org/10.1155/2011/621484>.
32. Cheng LYS, Jordan L, Gorugantula LM, Schneiderman E, Chen HS, Rees T. Salivary interleukin-6 and -8 in patients with oral cancer and patients with chronic oral inflammatory diseases. *J Periodontol* 2014; 85(7): 956–965. <https://doi.org/10.1902/jop.2013.130320>.
33. Arellano-Garcia ME, Hu S, Wang J, Henson B, Zhou H, Chia D, et al. Multiplexed immunobead-based assay for detection of oral cancer protein biomarkers in saliva. *Oral Dis* 2008; 14(8): 705–712. <https://doi.org/10.1111/j.1601-0825.2008.01488.x>.
34. Lourenço AG, Nakao C, Machado AA, Motta ACF, Tonani L, Candido RC, et al. Lactoferrin, a marker for periodontal disease. *Curr HIV Res* 2013; 11(3): 220–225. <https://doi.org/10.2174/1570162X11311030006>.
35. Nisha KJ, Suresh A, Anilkumar A, Padmanabhan S. MIP-1 $\alpha$  and MCP-1 as salivary biomarkers in periodontal disease. *Saudi Dent J* 2018; 30(4): 292–298. <https://doi.org/10.1016/j.sdentj.2018.07.002>.
36. Ebersole JL, Nagarajan R, Akers D, Miller CS. Targeted salivary biomarkers for discrimination of periodontal health and disease(s). *Front Cell Infect Microbiol* 2015; 5. <https://doi.org/10.3389/fcimb.2015.00062>.
37. Teles RP, Likhari V, Socransky SS, Haffajee AD. Salivary cytokine levels in subjects with chronic periodontitis and in periodontally healthy individuals: a cross-sectional study. *J Periodontol Res* 2009; 44(3): 411–417. <https://doi.org/10.1111/j.1600-0765.2008.01119.x>.
38. Gupta N, Gupta ND, Gupta A, Khan S, Bansal N. Role of salivary matrix metalloproteinase-8 (MMP-8) in chronic periodontitis diagnosis. *Front Med* 2015; 9(1): 72–76. <https://doi.org/10.1007/s11684-014-0347-x>.
39. Miller CS, King CP, Langub MC, Kryscio RJ, Thomas MV. Salivary biomarkers of existing periodontal disease. *J Am Dent Assoc* 2006; 137(3): 322–329. <https://doi.org/10.14219/jada.archive.2006.0181>.

40. Engebretson SP, Grbic JT, Singer R, Lamster IB. GCF IL-1beta profiles in periodontal disease. **J Clin Periodontol** 2002; 29(1): 48–53. <https://doi.org/10.1034/j.1600-051x.2002.290108.x>.
41. Ryu OH, Choi SJ, Linares AMG, Song IS, Kim YJ, Jang KT, et al. Gingival epithelial cell expression of macrophage inflammatory protein-1alpha induced by interleukin-1beta and lipopolysaccharide. **J Periodontol** 2007; 78(8): 1627–1634. <https://doi.org/10.1902/jop.2007.070066>.
42. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, et al. Proteomics. Tissue-based map of the human proteome. **Science** 2015; 347(6220): 1260419. <https://doi.org/10.1126/science.1260419>.
43. Ding C, Ji X, Chen X, Xu Y, Zhong L. TNF- $\alpha$  gene promoter polymorphisms contribute to periodontitis susceptibility: evidence from 46 studies. **J Clin Periodontol** 2014; 41(8): 748–759. <https://doi.org/10.1111/jcpe.12279>.
44. Golub LM, Lee HM, Greenwald RA, Ryan ME, Sorsa T, Salo T, et al. A matrix metalloproteinase inhibitor reduces bone-type collagen degradation fragments and specific collagenases in gingival crevicular fluid during adult periodontitis. **Inflamm Res** 1997; 46(8): 310–319. <https://doi.org/10.1007/s000110050193>.
45. Al-Shammari KF, Giannobile WV, Aldredge WA, Iacono VJ, Eber RM, Wang HL, et al. Effect of non-surgical periodontal therapy on C-telopeptide pyridinoline cross-links (ICTP) and interleukin-1 levels. **J Periodontol** 2001; 72(8): 1045–1051. <https://doi.org/10.1902/jop.2001.72.8.1045>.
46. Napimoga MH, Nunes LHAC, Maciel AAB, Demasi APD, Benatti BB, Santos VR, et al. Possible involvement of IL-21 and IL-10 on salivary IgA levels in chronic periodontitis subjects. **Scand J Immunol** 2011; 74(6): 596–602. <https://doi.org/10.1111/j.1365-3083.2011.02605.x>.
47. Sorsa T, Hernández M, Leppilähti J, Munjal S, Netuschil L, Mäntylä P. Detection of gingival crevicular fluid MMP-8 levels with different laboratory and chair-side methods. **Oral Dis** 2010; 16(1): 39–45. <https://doi.org/10.1111/j.1601-0825.2009.01603.x>.
48. Johnson N, Ebersole JL, Kryscio RJ, Danaher RJ, Dawson 3rd D, Al-Sabbagh M, et al. Rapid assessment of salivary MMP-8 and periodontal disease using lateral flow immunoassay. **Oral Dis** 2016; 22(7): 681–687. <https://doi.org/10.1111/odi.12521>.
49. Finoti LS, Nepomuceno R, Pigossi SC, Corbi SC, Secolin R, Scarel-Caminaga RM. Association between interleukin-8 levels and chronic periodontal disease: a PRISMA-compliant systematic review and meta-analysis. **Medicine** 2017; 96(22):e6932. <https://doi.org/10.1097/MD.0000000000006932>.
50. Takahashi K, Azuma T, Motohira H, Kinane DF, Kitetsu S. The potential role of interleukin-17 in the immunopathology of periodontal disease. **J Clin Periodontol** 2005; 32(4): 369–374. <https://doi.org/10.1111/j.1600-051X.2005.00676.x>.
51. Noguchi K, Ishikawa I. The roles of cyclooxygenase-2 and prostaglandin E2 in periodontal disease. **Periodontology** 2007; 43: 85–101. <https://doi.org/10.1111/j.1600-0757.2006.00170.x>.

**How to cite this article:** Arroyo E, Oliveira-Alves MG, Chamorro-Petronacci CM, Marichalar-Mendía X, Bravo-López SB, Blanco-Carrión J, Pérez-Sayáns M. Protein-based salivary biomarkers for the diagnosis of periodontal diseases: Systematic review and meta-analysis. *J Taibah Univ Med Sc* 2023;18(4):737–747.