

CASE REPORT

Effectiveness of late coronary angioplasty in restoring sinus rhythm in inferior wall myocardial infarction with complete AV block

Avitso Liegise  | Lanu Naro

Department of Cardiology, Christian Institute of Health Sciences and Research (CIHSR), Dimapur, Nagaland, India

Correspondence

Avitso Liegise, Department of Cardiology, Christian Institute of Health Sciences and Research (CIHSR), Dimapur, Nagaland, India.
Email: avitso2669@gmail.com

Abstract

An elderly gentleman presenting late with inferior wall myocardial infarction and complete heart block underwent revascularization of an occluded proximal right coronary artery more than 4 days after the onset of symptoms and recovered sinus rhythm within 48 h of the procedure. There are no clear guidelines for time to percutaneous coronary intervention (PCI) in late-presenting myocardial infarction with complete atrioventricular blocks (CAVB), and studies looking at outcomes of primary PCI in this situation appear to be scarce. The case presented here is a good example of the relevance of late PCI.

KEYWORDS

complete heart block, inferior wall MI, late coronary angioplasty

1 | BACKGROUND

Patients presenting with myocardial infarction and high-grade AV blocks have worse outcomes, while early coronary interventions have shown improved results. Late revascularization outcomes, however, are not well documented with limited information available for guiding PCI in these situations with persisting complete heart blocks.

2 | CASE PRESENTATION

A gentleman in his early-sixties presented to a remote hospital elsewhere with a history of acute chest discomfort that was associated with light-headedness and shortness of breath. He is a chain smoker, nondiabetic, and nonhypertensive. There was no prior cardiac history.

The electrocardiogram (ECG) showed inferior wall ST elevation myocardial infarction with complete AV block. He was hemodynamically stable with a heart rate of 30 to 35/mt and was referred to our center for further management without any revascularization attempts. He arrived 24 h later still in complete heart block with a heart rate ranging from 40 to 45/mt and persisting ST elevation (Figure 1) with a blood pressure of 140/90 mmHg. A temporary pacemaker was inserted with a plan to do coronary angioplasty. However, the patient became delirious and restless and did not allow the team to proceed. Since he was hemodynamically stable with no ongoing angina, it was decided to delay the procedure a little while.

While in the ward he became breathless and went into distress with low blood pressure despite adequate pacing. He was intubated by the on-call doctor, started on inotropes, and shifted back to the catheterization laboratory for

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.




FIGURE 1 ECG at presentation showing inferior wall ST elevation and complete AV block.




FIGURE 2 Angiogram showing proximally occluded RCA after successful wiring across the occlusion.




FIGURE 3 RCA angiogram after successful recanalization and stenting with two drug-eluting stents.

revascularization. The coronary angiogram showed a single vessel disease with a 100% proximally occluded right coronary artery (Figure 2). Complete revascularization of the vessel was achieved with two drug-eluting stents (Figure 3) and over the next 36–48 h he was slowly taken off inotropic support (noradrenaline). On the second postprocedure day, he recovered sinus rhythm with a first-degree heart block and a PR interval of 280 ms. He was extubated the same day and taken off temporary pacemaker support the following day. The patient was discharged 2 days later maintaining sinus rhythm and a much reduced PR interval of 204 ms (Figure 4). The echocardiogram showed a

mild left ventricular systolic dysfunction with no other mechanical complications.

3 | OUTCOME AND FOLLOW-UP

The patient was reviewed 2 weeks later with the ECG showing sinus rhythm and no evidence of atrioventricular conduction delay (Figure 5). He reported no difficulty with routine activities. Compliance with medications has been good and he was advised to review after a month.




FIGURE 4 Discharge ECG showing sinus rhythm with mildly prolonged PR interval.




FIGURE 5 Follow-up ECG at 2 weeks with normal PR interval.

4 | DISCUSSION

Guidelines state that revascularization should be considered in late-presenting MI's with high-grade AV blocks,¹ but clear recommendations for PCI time in these situations

are lacking. Studies focusing on outcomes in these situations are limited with complete heart block being a bad prognostic marker even in the primary PCI era.² In a retrospective study looking at AV block recovery following late PCI in inferior wall MI with high-grade AV blocks

(82% of them had CAVB), a median time to PCI of 4 days resulted in complete recovery of 1:1 AV conduction for all patients at discharge.³

A similar case report by Liang et al.⁴ suggested that late coronary intervention in a patient presenting with IWMI complicated by complete heart block avoided a potential permanent pacemaker implant. However, in this case, the AV block was developed after presenting late to the hospital, and complete revascularization was achieved within a short span of developing the conduction block.

The AV node tissue has been described as resistant to permanent damage due to its high intracellular glycogen content,⁵ unique blood supply involving adjacent septal perforators, and ability to absorb oxygen and nutrients from the neighboring venous sinusoids.⁶ Conduction system abnormalities in acute posterior-inferior infarction are hence usually transient. However, it has been observed that those with persistence of such disturbance have more extensive damage to the conduction system involving the proximal as well as distal areas.⁷ It is in this group of patients with persistent high-grade AV blocks that revascularization, even if late, may result in good outcomes.

5 | CONCLUSION

1. Complete AV block in inferior wall myocardial infarction is a reversible complication despite late presentation.
2. Late revascularization of a completely occluded RCA should be attempted to potentially avoid permanent pacemaker implantation.

AUTHOR CONTRIBUTIONS

Avitso Liegise: Conceptualization; formal analysis; investigation; writing – original draft. **Lanu Naro:** Data curation; investigation; writing – review and editing.

ACKNOWLEDGMENTS

No funds, grants, or other support were received for this case report.

FUNDING INFORMATION

None.

CONFLICT OF INTEREST STATEMENT

Funding and/or Conflicts of interests/Competing interests: Avitso Liegise and Lanunaro declare that they have

no conflict of interest, and have received no financial support for the publication of this case report.

DATA AVAILABILITY STATEMENT

Data available on request due to privacy/ethical restrictions.

CONSENT

Written informed consent was obtained from the patient to publish this report in accordance with the journal's patient consent policy.

ORCID

Avitso Liegise  <https://orcid.org/0000-0002-9107-566X>

REFERENCES

1. Ibanez B, James S, Agewall S, et al. 2017 ESC guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: the task force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2018;39(2):119-177.
2. Gang UJO, Hvelplund A, Pedersen S, et al. High-degree atrioventricular block complicating ST-segment elevation myocardial infarction in the era of primary percutaneous coronary intervention. *Europace*. 2012;14(11):1639-1645.
3. Fan X, Maharjan P, Liu P, Bai L. Effect of primary PCI on the recovery of atrioventricular block in inferior STEMI patients with late presentation (>12 hours): insights from a single center 10-year experience. *J Investig Med off Publ Am Fed Clin Res*. 2020;68(5):1011-1014.
4. Liang M, Chin J, Pasupati S. Complete heart block in late presentation of inferior STEMI successfully treated with percutaneous coronary intervention. *J Invasive Cardiol*. 2011;23(9):E21 9-E221.
5. Regional distribution of glycogen and phosphorylase in the ventricles of the heart. *Circ Res*. 1964;14(3):202-211.
6. Kawamura Y, Yokoyama H, Kitayama K, et al. Clinical impact of complete atrioventricular block in patients with ST-segment elevation myocardial infarction. *Clin Cardiol*. 2021; 44(1):91-99.
7. Waller BF, Gering LE, Branyas NA, Slack JD. Anatomy, histology, and pathology of the cardiac conduction system—part V. *Clin Cardiol*. 1993;16(7):565-569.

How to cite this article: Liegise A, Naro L.

Effectiveness of late coronary angioplasty in restoring sinus rhythm in inferior wall myocardial infarction with complete AV block. *Clin Case Rep*. 2024;12:e8226. doi:[10.1002/ccr3.8226](https://doi.org/10.1002/ccr3.8226)