

# Defining counselling in contraceptive information and services: outcomes from an expert think tank

# Moazzam Ali 💿 ,<sup>1</sup> Nguyen Toan Tran 💿 <sup>2,3</sup>

<sup>1</sup>Department of Reproductive Health and Research, World Health Organization, Geneva, Switzerland <sup>2</sup>Australian Centre for Public and Population Health Research, Faculty of Health, University of Technology Sydney, Sydney, New South Wales, Australia <sup>3</sup>Faculty of Medicine, University of Geneva, Geneva, Switzerland

#### Correspondence to

Dr Moazzam Ali, Department of Reproductive Health and Research, World Health Organization, Geneva, Switzerland; alimoa@who.int

Received 18 March 2021 Accepted 17 May 2021 Published Online First 14 June 2021



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Ali M, Tran NT. BMJ Sex Reprod Health 2022;48:79–81. As a global public health good, contraception is a core component of the Sustainable Development Goal 3.7 (universal access to sexual and reproductive healthcare services, including family planning). Fundamentally, access to contraceptive information and services is a human right that advances other human right aspects. Quality contraceptive information and services reinforce people's freedom to determine the number and spacing of their children and offer a range of potential benefits encompassing women's empowerment, economic development, education and improved health outcomes, including maternal and child health. However, in low- and middle-income countries, around 218 million women of reproductive age still have an unmet need for contraception in 2019 – meeting this need could drop annually an estimated 111 to 35 million unintended pregnancies, 35 to 10 million unsafe abortions, and 299 000 maternal deaths to 113 000.<sup>1</sup>

Many contraceptive users discontinue their methods or fail to use them optimally.<sup>2 3</sup> Quality contraceptive counselling has the potential to play a key role in supporting individuals select a method that matches their needs and expectations, mitigate any side effects, continue their method, or turn to other options, thereby reducing the unmet need for contraception, among other factors.<sup>4</sup> There is, however, no standard definition of contraceptive counselling, although the centrality of quality counselling is underscored in different frameworks and programmatic and policy recommendations, as illustrated below.

Relevant documents include the Bruce framework, which was published in 1990 and identified six dimensions of quality for family planning services: technical competence, follow-up and continuity mechanisms, and the appropriate constellation of services, in addition to three dimensions that are specifically related to counselling - choice of methods, the information given to clients, and interpersonal relations.<sup>5</sup> Other quality components of contraceptive counselling - given in a client-centred approach - were outlined in recent years, including needs assessment, trust-building with clients, tailored communication. shared decisionmaking (by eliciting and responding to client preferences), method choice, and follow-up.4 <sup>6</sup> Attention has been called to the specific counselling needs of adolescents, such as dual protection against pregnancy and sexually transmitted infections and respect for adolescents' autonomy.<sup>7</sup> As for the World Health Organization (WHO), its 2016 Selected Practice Recommendations for Contraceptive Use include guidelines for counselling content for each contraceptive method (focusing primarily on side effects and dual protection), while its 2018 Global Handbook for Family Planning Providers made further recommendations on interpersonal qualities, including respect and confidentiality.

Counselling can either occur face-toface, using digital technology, or a combination thereof. For example, the definition of the Population Council includes only face-to-face interactions involving a twoway communication between a counsellor and an individual or couple, or a counsellor and a group. The counsellor gives evidence-based information and assists the individual, couple or group to make a decision about behaviour change, taking into account the feelings and concerns



79

of the client.<sup>8</sup> Other authors define counselling as an interactive process between a provider and a client intended to help the client achieve a reproductive health goal.<sup>9</sup>

Drawing from the extant literature on the different dimensions of contraceptive counselling and following a systematic review on the effectiveness of counselling strategies for modern contraceptive methods,<sup>10</sup> a think tank on the topic was held at WHO, Geneva in May 2019. The think tank gathered representatives from academia, implementing agencies and international organisations working in the contraception field (see Acknowledgements) and had the objective of identifying research and guidance gaps and other opportunities for improving the quality of contraceptive counselling. Notably, the discussions culminated in the development of a comprehensive definition of counselling, which can hopefully serve as a benchmark for further development, update and use by academia and implementing agencies. The definition reads as follows:

Contraceptive counselling is defined as the exchange of information on contraceptive methods based on an assessment of the client's needs, preferences, and lifestyle to support decision-making as per the client's intentions. This includes the selection, discontinuation or switching of a contraceptive method. The key principles are based on: coercionfree and informed choice; neutral, understandable and evidence-based information; collaborative and confidential decision-making process; ensuring respectful care, dignity, and choice.

The definition took into consideration the core dimensions articulated in the literature and mirrors the principles that were deliberated by the think tank: a two-way discussion and decision-making process between the client and provider, sharing objective and user-friendly information, and an approach fundamentally grounded in the rights to autonomy, dignity, privacy, respect and participation, among others.

Operationalising this comprehensive definition into clinical services will require a practical balance between the constraints inherent to resourcechallenged settings and achieving programmatic goals. Based on our client-oriented definition, reaching traditional programmatic goals, such as increasing couple-year protection and modern contraceptive prevalence, should be questioned from the perspective of clients' autonomy, freedom from coercion, and satisfaction. However, further research is needed to develop and test metrics that capture our definition dimensions of quality counselling and focus on clients' reproductive autonomy and empowerment in decision-making. Acknowledgements The authors are grateful to the Bill and Melinda Gates Foundation, USA; The International Federation of Gynaecology and Obstetrics (FIGO), UK; Harvard School of Public Health, USA; HRH2030/Chemonics, USA; Ibis Reproductive Health, Institute of Tropical Medicine, Antwerp, Belgium; International Planned Parenthood Federation (IPPF), UK; Johns Hopkins Bloomberg School of Public Health Center for Communication Program (CCP), USA; MSI Reproductive Choices, UK; Population Council, USA; Population Services International (PSI), Tanzania; United Nations Population Fund (UNFPA); University of Cape Town, South Africa; United States Agency for International Development (USAID), USA; World Health Organization (WHO).

**Contributors** MA and NTT jointly wrote this editorial and contributed equally.

**Funding** This work has been supported by the UNDP-UNFPA-UNICEF-WHO-World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), a cosponsored programme executed by the World Health Organization (WHO).

**Disclaimer** This report contains the collective views of an international group of experts, and does not necessarily represent the decisions or the stated policy of the World Health Organization (WHO).

Competing interests None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4. 0/.

## ORCID iDs

Moazzam Ali http://orcid.org/0000-0001-6949-8976 Nguyen Toan Tran http://orcid.org/0000-0001-7134-7878

### REFERENCES

- Guttmacher Institute. Adding it up investing in sexual and reproductive health in low- and middle-income countries. Fact Sheet. July, 2020. Available: https://www.guttmacher.org/ sites/default/files/factsheet/adding-it-up-investing-in-sexualreproductive-health.pdf [Accessed 17 Dec 2020].
- 2 Polis C, Bradley SEK, Bankole A, et al. Contraceptive failure rates in the developing world: an analysis of demographic and health survey data in 43 countries. New York: Guttmacher Institute, 2016. https://www.guttmacher.org/report/ contraceptive-failure-rates-in-developing-world
- 3 Ali MM, Cleland JG, Shah IH, World Health Organization. Causes and consequences of contraceptive discontinuation: evidence from 60 Demographic and Health Surveys. Geneva: World Health Organization, 2012.
- 4 Dehlendorf C, Krajewski C, Borrero S. Contraceptive counseling: best practices to ensure quality communication and enable effective contraceptive use. *Clin Obstet Gynecol* 2014;57:659.
- 5 Bruce J. Fundamental elements of the quality of care: a simple framework. *Stud Fam Plann* 1990;21:61–91.

Twitter Moazzam Ali @Moazzam2000

- 6 Holt K, Dehlendorf C, Langer A. Defining quality in contraceptive counseling to improve measurement of individuals' experiences and enable service delivery improvement. *Contraception* 2017;96:133–7.
- 7 Raidoo S, Kaneshiro B. Contraception counseling for adolescents. Curr Opin Obstet Gynecol 2017;29: 310–5.
- 8 Population Council. A client-centered approach to reproductive health. Islamabad, Pakistan, 2005. Available:

https://www.popcouncil.org/uploads/pdfs/CCA\_ TrainersManual.pdf [Accessed 14 Mar 2021].

- 9 Zapata LB, Tregear SJ, Curtis KM, *et al.* Impact of contraceptive counseling in clinical settings: a systematic review. *Am J Prev Med* 2015;49:S31–45.
- 10 Cavallaro FL, Benova L, Owolabi OO, et al. A systematic review of the effectiveness of counselling strategies for modern contraceptive methods: what works and what doesn't? BMJ Sex Reprod Health 2020;46:254–69.