

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Available online at

# **ScienceDirect**

www.sciencedirect.com

#### Elsevier Masson France





# Correspondence

# Letter on the article "Long-COVID: Cognitive deficits (brain fog) and brain lesions in non—hospitalized patients"



In "LONG-COVID: COGNITIVE DEFICITS (BRAIN FOG) AND BRAIN LESIONS IN NON—HOSPITALIZED PATIENTS", Hugon concluded that "neurological complications of COVID may be detected in moderately affected individuals and extensive ......[1]." For long COVID, brain fog is an important clinical problem [2,3]. In a recent report, female is more affected than male. For the pathogenesis, it is still not conclusive. The brain fog might be associated with immunophathology. Also, after COVID-19, there is an increased blood viscosity [4,5]. The excessive high blood viscosity might affect brain function [6].

#### **Conflict of interest**

None.

### References

- Hugon J. Long-COVID: cognitive deficits (brain fog) and brain lesions in non-hospitalized patients. Presse Med 2021 Oct 27:104090 Online ahead of print. doi: 10.1016/j.lpm.2021.104090.
- [2] Asadi-Pooya AA, Akbari A, Emami A, Lotfi M, Rostamihosseinkhani M, Nemati H, Barzegar Z, Kabiri M, Zeraatpisheh Z, Farjoud-Kouhanjani M, Jafari A, Sasannia F, Ashrafi S, Nazeri M, Nasiri S, Shahisavandi M. Long COVID syndrome-associated brain fog. J Med Virol 2021 Oct 21 Online ahead of print. doi: 10.1002/jmv.27404.

- [3] Tabacof L, Tosto-Mancuso J, Wood J, Cortes M, Kontorovich A, McCarthy D, Rizk D, Rozanski G, Breyman E, Nasr L, Kellner C, Herrera JE, Putrino D. Post-acute COVID-19 syndrome negatively impacts physical function, cognitive function, healthrelated quality of life and participation. Am J Phys Med Rehabil 2021 Oct 20 Online ahead of print. doi: 10.1097/PHM.000000000001910.
- [4] Joob B, Wiwanitkit V. Blood viscosity of COVID-19 patient: a preliminary report. Am J Blood Res 2021 Feb 15;11(1):93–5.
- [5] Joob B, Wiwanitkit V. Expected viscosity after COVID-19 vaccination, hyperviscosity and previous COVID-19. Clin Appl Thromb Hemost 2021;27: 10760296211020833 Jan-Dec.
- [6] Small C, Land AM, Haist SA, Estrada CA, Snyder ED. Managing cognitive load to uncover an unusual cause of syncope: exercises in clinical reasoning. J Gen Intern Med 2016;31(2):247–51 Feb.

Rujittika Mungmunpuntipantip\* Viroj Wiwanitkit Private Academic Consultant, Bangkok Thailand Honorary professor, Dr DY Patil University, Pune, India

\*Corresponding author at: Private Academic Consultant, Bangkok Thailand.

E-mail address: rujittika@gmail.com (R. Mungmunpuntipantip).

Available online 28 April 2022