



# Draft Genome Sequence of *Acidovorax* sp. Strain NB1, Isolated from a Nitrite-Oxidizing Enrichment Culture

Hiroto Ide,<sup>a</sup> Kento Ishii,<sup>a</sup> Hirotsugu Fujitani,<sup>b,c</sup> Satoshi Tsuneda<sup>a,c</sup>

<sup>a</sup>Department of Life Science and Medical Bioscience, Waseda University, Tokyo, Japan

<sup>b</sup>Biomedical Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Ibaraki, Japan

<sup>c</sup>Research Organization for Nano and Life Innovation, Waseda University, Tokyo, Japan

**ABSTRACT** Here, we report the draft genome sequence of *Acidovorax* sp. strain NB1, isolated from an enrichment culture of nitrite-oxidizing bacteria (NOB). Genes involved in denitrification were found in the draft genome of NB1. The closest strain to NB1 based on genomic relatedness is *Acidovorax* sp. strain GW101-3H11, with 91.5% average nucleotide identity.

Nitrogen exists in various forms, and microorganisms mediate its circulation in natural environments and engineered systems. Ammonia (NH<sub>3</sub>) is eventually converted to nitrogen gas (N<sub>2</sub>) through nitrification and denitrification, which are the key processes of the global nitrogen cycle. *Acidovorax* spp., which are of the family Comamonadaceae in the order Betaproteobacteriales, are known as denitrifying bacteria and contribute to biological nitrogen removal in wastewater treatment plants (1–3).

Here, we announce the draft genome sequence of *Acidovorax* sp. strain NB1, isolated from a nitrite-oxidizing bacterium (NOB) enrichment culture, and the genes involved in denitrification. Previously, our group enriched the environmentally important NOB from the genus “*Candidatus Nitrotoga*” in mineral salts medium, in which they likely provided by-products and nitrate to heterotrophic bacteria (4). Among the heterotrophs, *Acidovorax* sp. NB1 was isolated from the “*Candidatus Nitrotoga*” enrichment culture using a nutrient broth (NB) medium gelled by agar.

The pure culture of NB1 was cultivated in the NB medium at 29°C prior to DNA extraction and genome sequencing. The NB1 genome was sequenced with an Illumina MiSeq instrument and QIAseq FX DNA library kit, which generated paired-end read sequences of 301 bp at 263.5× coverage, on average. Platanus\_trim ([http://platanus.bio.titech.ac.jp/platanus\\_trim](http://platanus.bio.titech.ac.jp/platanus_trim)) was used to remove adapters and low-quality reads. The draft genome sequence of NB1 was assembled using Platanus v1.2.4 (5) and comprised 60 contigs with a total genome size of 5,493,953 bp (65.3% G+C content; N<sub>50</sub>, 332.59 kbp). Automated annotation of the coding sequence (CDS) region was performed using DFAST (6), generating 5,009 CDS features, 50 tRNAs, and 6 rRNAs. The genome sequences assigned to the genus *Acidovorax* were downloaded using NCBI genome downloading scripts (<https://github.com/kblin/ncbi-genome-download>), and the average nucleotide identity (ANI) was calculated using the Python module PYANI v0.2.8 (7). The ANI values between NB1 and the sequenced *Acidovorax* genomes were 81.9 to 91.5%. All software was run with default settings.

NB1 has the whole gene set necessary for nitrate reduction (*narK2K1GHJI*), and the gene that regulates their transcription, *narXL*, is located upstream (Fig. 1). The *narXL-narK-narGHJI* operon is conserved in most Betaproteobacteriales genomes (8, 9). Dissimilatory-nitrite-reduction-related genes, such as *nirK* or *nirS*, were not found in the NB1 genome. Other *Acidovorax* strains that have nitrate reduction potential lack dissimilatory nitrite reductases (10–12). The *nosZRDFYLL* operon involved in nitrous

**Citation** Ide H, Ishii K, Fujitani H, Tsuneda S. 2019. Draft genome sequence of *Acidovorax* sp. strain NB1, isolated from a nitrite-oxidizing enrichment culture. Microbiol Resour Announc 8:e00547-19. <https://doi.org/10.1128/MRA.00547-19>.

**Editor** Frank J. Stewart, Georgia Institute of Technology


**Copyright** © 2019 Ide et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Hirotsugu Fujitani, [hirotosugu.fujitani@aist.go.jp](mailto:hirotosugu.fujitani@aist.go.jp), or Satoshi Tsuneda, [stsuneda@waseda.jp](mailto:stsuneda@waseda.jp).

**Received** 6 June 2019

**Accepted** 24 July 2019

**Published** 15 August 2019



**FIG 1** Schematic representation of the nitrate reductase (NAR) and nitrous oxide reductase (NOS) operons of *Acidovorax* sp. NB1. Each arrow represents a CDS with the gene subunit letter identity indicated above the arrow.

oxide ( $N_2O$ ) reduction is present in the NB1 genome, and *nosR*, the regulator gene of *nosZ*, lies downstream of its cluster. The order of nitrous oxide reductases is the same as that of “*Candidatus Magnetobacterium*” spp. of the phylum *Nitrospirae* (13). With respect to nitric oxide (NO) reduction, only a nitric oxide reductase large subunit gene (*norB*) was identified in the NB1 genome. Considering that NB1 has a nitrate reductase gene cluster, reciprocal feeding may occur between NB1 and “*Candidatus Nitrotoga*” spp. in the enrichment culture.

**Data availability.** The draft genome sequence of *Acidovorax* sp. NB1 has been deposited in DDBJ/ENA/GenBank under the accession number [BJHU00000000](https://doi.org/10.1101/2018.08.14.238111). The BioProject number is [PRJDB5517](https://doi.org/10.1093/bioinformatics/btx713), and the BioSample number is [SAMD00074225](https://doi.org/10.1093/bioinformatics/btx713). The associated raw sequence data have been deposited in the NCBI under the accession number [DRP005091](https://doi.org/10.1093/bioinformatics/btx713).

## ACKNOWLEDGMENTS

Genome sequencing and assembly of *Acidovorax* sp. NB1 were supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI grant 16H06279 (to H.F.). The other part of this research was supported by a Grant-in-Aid for Young Scientists (B) from the JSPS (grant 16K18609 to H.F.) and a Grant-in-Aid for JSPS Research Fellowship (grant 18J22997 to K.I.).

## REFERENCES

- Etchebehere C, Errazquin I, Barrandeguy E, Dabert P, Moletta R, Muxi L. 2001. Evaluation of the denitrifying microbiota of anoxic reactors. *FEMS Microbiol Ecol* 35:259–265. <https://doi.org/10.1111/j.1574-6941.2001.tb00811.x>.
- Mechichi T, Stackebrandt E, Fuchs G. 2003. Alicyclophilus denitrificans gen. nov., sp. nov., a cyclohexanol-degrading, nitrate-reducing  $\beta$ -proteobacterium. *Int J Syst Evol Microbiol* 53:147–152. <https://doi.org/10.1099/ijs.0.02276-0>.
- Hoshino T, Terahara T, Tsuneda S, Hirata A, Inamori Y. 2005. Molecular analysis of microbial population transition associated with the start of denitrification in a wastewater treatment process. *J Appl Microbiol* 99:1165–1175. <https://doi.org/10.1111/j.1365-2672.2005.02698.x>.
- Ishii K, Fujitani H, Soh K, Nakagawa T, Takahashi R, Tsuneda S. 2017. Enrichment and physiological characterization of a cold-adapted nitrite-oxidizing *Nitrotoga* sp. from an eelgrass sediment. *Appl Environ Microbiol* 83:e00549-17. <https://doi.org/10.1128/AEM.00549-17>.
- Kajitani R, Toshimoto K, Noguchi H, Toyoda A, Ogura Y, Okuno M, Yabana M, Harada M, Nagayasu E, Maruyama H, Kohara Y, Fujiyama A, Hayashi T, Itoh T. 2014. Efficient de novo assembly of highly heterozygous genomes from whole-genome shotgun short reads. *Genome Res* 24:1384–1395. <https://doi.org/10.1101/gr.170720.113>.
- Tanizawa Y, Fujisawa T, Nakamura Y. 2018. DFAST: a flexible prokaryotic genome annotation pipeline for faster genome publication. *Bioinformatics* 34:1037–1039. <https://doi.org/10.1093/bioinformatics/btx713>.
- Pritchard L, Glover RH, Humphris S, Elphinstone JG, Toth IK. 2016. Genomics and taxonomy in diagnostics for food security: soft-rotting enterobacterial plant pathogens. *Anal Methods* 8:12–24. <https://doi.org/10.1039/C5AY02550H>.
- Price MN, Dehal PS, Arkin AP. 2008. Horizontal gene transfer and the evolution of transcriptional regulation in *Escherichia coli*. *Genome Biol* 9:R4. <https://doi.org/10.1186/gb-2008-9-1-r4>.
- Stewart V. 2003. Nitrate- and nitrite-responsive sensors NarX and NarQ of proteobacteria. *Biochem Soc Trans* 31:1–10. <https://doi.org/10.1042/bst0310001>.
- Huang Y, Li H, Rensing C, Zhao K, Johnstone L, Wang G. 2012. Genome sequence of the facultative anaerobic arsenite-oxidizing and nitrate-reducing bacterium *Acidovorax* sp. strain NO1. *J Bacteriol* 194:1635–1636. <https://doi.org/10.1128/JB.06814-11>.
- Ehsani E, Jauregui R, Geffers R, Jarek M, Boon N, Pieper DH, Vilchez-Vargas R. 2015. First draft genome sequence of the *Acidovorax caeni* sp. nov. type strain R-24608 (DSM 19327). *Genome Announc* 3:e01378-15. <https://doi.org/10.1128/genomeA.01378-15>.
- Vanechoutte M, Janssens MJ, Avesani V, Delmée M, Deschaght P. 2013. Description of *Acidovorax wautersii* sp. nov. to accommodate clinical isolates and an environmental isolate, most closely related to *Acidovorax avenae*. *Int J Syst Evol Microbiol* 63:2203–2206. <https://doi.org/10.1099/ijs.0.046102-0>.
- Lin W, Jogler C, Schüller D, Pan Y. 2011. Metagenomic analysis reveals unexpected subgenomic diversity of magnetotactic bacteria within the phylum *Nitrospirae*. *Appl Environ Microbiol* 77:323–326. <https://doi.org/10.1128/AEM.01476-10>.