

Clinical characteristics and molecular epidemiology of children with meningitis in Tehran, Iran: a prospective study

A. Pormohammad¹, S. Lashkarbolouki², T. Azimi^{3,4}, P. Gholizadeh⁵, N. Bostanghadiri¹, H. Safari⁶, S. Armin⁷, N. Mohtavinejad⁸ and F. Fallah¹

1) Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran, 2) Department of Biology, Tehran North Branch, Islamic Azad University, Tehran, Iran, 3) Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran, 4) Students Scientific Research Center, Tehran University of Medical Sciences, Tehran, Iran, 5) Department of Medical Microbiology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran, 6) Health Promotion Research Center, school of public health, Iran University of Medical Sciences, Tehran, Iran, 7) Pediatric Infections Research Center, Research Institute for Children Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran and 8) Department of Radio Pharmacy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

Abstract

The molecular epidemiology of meningitis in children is unclear in Iran, and data are scarce. We aimed to characterize its clinical and paraclinical features as well as to determine the distribution of genotype/capsular types of common bacterial meningitis agents in children in Iran. All children suspected to have meningitis aged 4 days to 15 years were enrolled onto a prospective cross-sectional study from January 2015 to September 2017. Diagnostic values of clinical features, cerebrospinal fluid and serum parameters were evaluated independently and in combination with each other by multivariate logistic regression to develop a diagnostic rule. Genotype/capsular types of all the isolates were determined by targeting serotype-specific genes with uniplex or multiplex PCR. Among 119 patients suspected of having meningitis, 43 had bacterial meningitis, 19 aseptic and one tuberculous; and there were 56 nonmeningitis cases (NMC). Presentation of four features at the same time—cerebrospinal fluid white blood cell count, protein, polymorphonuclear leukocytes and serum C-reactive protein—revealed 100% sensitivity and 86.4% specificity for diagnosis of bacterial meningitis. *Haemophilus influenzae* type b (60%), *Streptococcus pneumoniae* serotype 3 (28.5%) and *Neisseria meningitidis* B (63.5%) were the most prevalent serotypes. This study demonstrated that a well-designed combination of clinical and paraclinical features is useful, but these combinations are not good enough to be relied on as stand-alone exclusionary tests for the diagnosis of bacterial meningitis. In addition, public immunization of infants with the most prevalent bacterial meningitis serotypes is recommended.

© 2019 Published by Elsevier Ltd.

Keywords: Children, diagnosis, genotype, meningitis, molecular epidemiology

Original Submission: 8 March 2019; **Revised Submission:** 18 July 2019; **Accepted:** 20 August 2019

Article published online: 29 August 2019

Corresponding author: F. Fallah, Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

E-mails: pormohammadali@yahoo.com (A. Pormohammad), falah@sbmu.ac.ir (F. Fallah)

Introduction

Meningitis is a medical emergent illness which leads to devastating neurologic sequelae and is almost always life-threatening

without treatment [1,2]. Its prognosis depends heavily on rapid diagnosis and early sufficient treatment [3]. Children and neonates are at a high risk of neurologic disorders, especially meningitis [4]. There are many different types of meningitis, including bacterial, viral, fungal, parasitic, noninfectious and aseptic meningitis. Here we use the term 'aseptic meningitis' to describe both confirmed and presumed cases of viral meningitis [5]. Many studies previously suggested cerebrospinal fluid (CSF) analysis and clinical assessments to differentiate between the various types of meningitis [3,4,6–10]. However, now researchers pay less attention to that in favor of distinguishing suspected cases of meningitis from other infections (e.g.

sinusitis, encephalitis, septicaemia) and neurologic disorders that may present similarly (e.g. hydrocephalus, seizure). Thus, well-designed studies are essential to accurately and rapidly distinguish suspected cases of meningitis by using clinical and laboratory findings that add potential value to standard laboratory tests [3,11].

A systematic search performed in our recent meta-analysis indicated that the distribution of common bacterial meningitis agent genotypes or capsular types is unclear in Iran—information essential for vaccine and prevention strategies [12]. Furthermore, some additional factors such as having poor laboratory facilities, as well as in a region of the world with high rates of advanced HIV- or tuberculosis-infected patients and drug-resistant pathogens make the diagnosis and prevention of meningitis complicated in Iran [13]. These issues create an exceptional challenge for clinicians in terms of the management of meningitis [13,14]. The 7-valent pneumococcal conjugate vaccine has become commercially available; however, it is not a mandatorily prescribed vaccine in Iran, although the conjugate polysaccharide vaccine against *Haemophilus influenzae* type b (Hib) is [12].

In this study, we focused on characteristics of clinical features and CSF findings solely and in combination with each other in children and neonates with suspected meningitis to create a multivariable diagnosis model for the diagnosis of meningitis. In addition, we aimed to determine the distribution of genotype/capsular types of common bacterial meningitis agents in children in Iran for vaccine and prevention strategies.

Materials and methods

Ethics statement

The ethics committee of Shahid Beheshti University of Medical Sciences approved this study (IR.SBMU.RETECH.REC.1395.212).

Study population and study design

We included all consecutive children (defined as patients <16 years of age) who were suspected of having meningitis in Baqiyatallah and Mofid paediatric hospitals (territorial hospitals in Tehran) in a retrospective cross-sectional study from March 2015 to March 2017. In a masked study, all of the clinical diagnoses, molecular assays, and CSF and serum analyses were performed by separate personnel unknown to each other. All included patients were completely vaccinated according to Iran's vaccine schedule.

Patients with suspected meningitis had any of the following criteria: (a) fever with signs of meningeal irritation, bulging fontanelles, unexplained irritability, persistent nausea and

vomiting, petechial rash or purpura; (b) unexplained alteration of consciousness with or without fever; (c) fever with no localizing signs in an ill or toxic child; and (d) complex febrile seizure with or without fever. Inclusion criteria were age <16 years old and suspected diagnosis of meningitis. Exclusion criteria were neurosurgical procedure performed before the onset of meningitis, presence of a shunt within the central nervous system or presence of a known immunodeficiency (Fig. 1). Duplicate samples from the same patient were excluded.

Conventional bacteriologic diagnostic tests such as smear staining and culture tests were performed for CSF and blood samples according to the laboratory methods for the diagnosis of meningitis reported by the World Health Organization [15] and the US Centers for Disease Control and Prevention guidelines (<https://www.cdc.gov/meningitis/lab-manual/index.html>) (Fig. 2).

On the basis of clinical and laboratory combination tests, confirmed patients were categorized into three groups: bacterial meningitis, aseptic meningitis and NMC. Clinical status was evaluated with the Glasgow Outcome Scale, which ranges from 1 to 5, as follows: 1 = death; 2 = persistent vegetative state (the patient is unable to interact with the environment); 3 = severe disability; 4 = moderate disability; 5 = good recovery [4]. Severe or moderate disability was defined as any of the following conditions: muscle weakness and immobility in one or more limbs; hearing loss; microcephaly; spasticity hydrocephalus; and seizure disorder. For the purpose of this analysis, a good outcome was defined as a score of 5, and a poor outcome was a score of 1 to 4. All demographic characteristics, serum and CSF findings were recorded for each patient.

DNA extraction

Upon arrival at the laboratory, 200 to 400 µL of each CSF and serum sample was extracted according to the Roche High Pure template DNA kit manual (Roche, Basel, Switzerland).

PCR survey

Conventional PCR for species identification was performed with primers specific for amplifying the genes *CtrA* (*Neisseria meningitidis*), *lytA* (*Streptococcus pneumoniae*) and *hpd* (*H. influenzae*) for all samples from patients suspected to have meningitis. PCR was performed as described previously [16]. Standard strains were obtained from the Persian Type Culture Collection (PTCC) and Microbial Collection of Pasteur Institute of Iran, *H. influenzae* Type Cultures (ATCC 35056), *N. meningitidis* (PTCC 1760) and *S. pneumoniae* (PTCC 1800 and PTCC 1240). Specificity of the three primer sets was determined by using genomic DNA from bacteria and viruses which are most likely to be present in CSF samples, such as




FIG. 1. Flow diagram of patient selection.

group B *Streptococcus*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Enterobacter*, *Staphylococcus aureus*, herpes simplex virus, Epstein-Barr virus, cytomegalovirus and varicella-zoster virus. Also, we prepared human genomic DNA from the microbiology department of Shahid Beheshti University. We used Taq DNA Polymerase 2× Master Mix Red (Ampliqon, Odense, Denmark) and optimized it in a gradient cyler (Mastercycler Gradient; Eppendorf, Hamburg, Germany). Primers, PCR assembly and cycling conditions are described in [Supplementary Table S1](#).

Molecular genotyping by multiplex PCR

First, all of the confirmed bacterial meningitis CSF samples were examined with minor modifications designed to cover most of the *N. meningitidis*, *S. pneumoniae* and *H. influenzae* predominant serotypes reported in Asia and Africa [17,18]. Accordingly,

each reaction was designed as uniplex or multiplex for targeting the serotype-specific genes of individual serotypes; it included an internal positive control ([Supplementary Table S1](#)).

Statistical analysis

All statistical analyses were performed by SPSS 16.0 software (IBM, Armonk, NY, USA) and MedCalc 12.1.4 (MedCalc Software, Ostend, Belgium). The chi-square test, Fisher exact test, Student *t* test or Mann-Whitney *U* test were used for comparing categorical and continuous variables and for comparisons between groups in nonnormally distributed variables. We determined a cutoff point for each scale variable (biomarkers) to differentiate the bacterial and aseptic meningitis from other similar disorders by using the receiver operating characteristic curve. Likewise, for each biomarker, sensitive specific values with 95% confidence interval were calculated on




FIG. 2. *Haemophilus influenzae*, *Neisseria meningitidis* and *Streptococcus pneumoniae* colonies in upper images, and Gram staining in lower pictures.

the basis of the chosen cutoff point. Results are presented as mean \pm standard deviation, or number and percentage. We used logistic regression to examine the accuracy of association of various biomarkers in distinguishing bacterial and aseptic meningitis from the other three groups. All statistical tests were two tailed with $p < 0.05$.

Results

Among 119 patients suspected to have meningitis, 43 were confirmed as bacterial, 19 as aseptic and one as tuberculous, as well as 56 NMC, by using a combination of clinical and

TABLE 1. Clinical features and laboratory findings of study patients

Characteristic	Bacterial meningitis (n = 43)	Aseptic meningitis (n = 19)	NMC (n = 56)	Total (n = 118)	p
Male sex	25 (58)	10/17 (52.6)	32 (57.7)	67/116 (56.3)	0.91
Age group					0.17
<1 month	9 (21)	0	4/55 (5.3)	13/115 (11.3)	
1 month to 1 year	20 (46.5)	11/17 (64.7)	25/55 (44.7)	56/115 (48.7)	
1–7 years	9 (21)	4/17 (23.5)	19/55 (36.8)	32/115 (27.8)	
7–16 years	5 (11.6)	2/17 (11.8)	7/55 (13.2)	14/115 (12.2)	
Clinical features					
Fever	41 (95.3)	18 (95)	30 (53.6)	89 (75.5)	>0.005
Seizure	12 (28)	6 (31.5)	7/55 (12.7)	25/117 (21.4)	0.09
Sleepy	11/18 (61)	5/8 (62.5)	9/12 (75)	25/38 (65.8)	0.71
Hydrocephalus	3 (7)	1/17 (6)	0	4/115 (3.5)	0.14
Anorexia	4/35 (11.5)	4/13 (31)	6/36 (16.7)	14/84 (16.7)	0.28
Nausea, vomiting	19 (44.2)	12/18 (66.7)	18/56 (32)	49/117 (42)	0.03
Cough	6 (14)	5/18 (28)	15/55 (27.3)	26/116 (22.4)	0.24
Diarrhea	5 (11.5)	3/18 (16.7)	7/55 (12.7)	15/116 (13)	0.86
Poor feeding	5 (11.5)	0	9/55 (16.4)	14/115 (12.2)	0.19
Headache	7 (16.3)	4/17 (23.5)	8 (14.3)	19/116 (16.5)	0.66
Coryza	1/29 (3.4)	4/12 (33.3)	2/22 (9)	7/63 (11)	0.02
Neck stiffness	1/25 (4)	0	1/14 (7)	2/50 (4)	0.66
Rash	5 (11.5)	1/17 (6)	1/56 (1.8)	7/116 (6)	0.12
Daze	3 (7)	1 (5.3)	3 (5.4)	7/118 (6)	0.93
Double vision	2/7 (28.6)	0	3/6 (50)	5/14 (35.7)	0.53
Birth, term	7/9 (77.8)	6/6 (100)	11/11 (100)	24/26 (92.3)	0.13
Parent, relatives	20/30 (66.7)	7/10 (70)	24/34 (70.6)	51/74 (69)	0.94
Surgery history	1 (2.3)	0	2 (3.6)	3 (2.5)	0.7
Outcome, good	27/33 (82)	10/12 (83.3)	38/42 (90.5)	75/87 (86)	0.53
Lab findings					
Blood culture	3 (7)	0	3/55 (5.5)	6/115 (5.2)	0.02
CSF culture	3 (7)	0	0	3/115 (2.6)	0.1
CSF smear	2 (4.7)	0	0	2/118 (1.7)	0.31
Blood smear	1 (2.3)	0	0	1/116 (1)	0.63
Urine culture	3 (7)	0	1/55 (1.8)	4/113 (3.5)	0.26

Data are presented as n (%) or n/N (%).

CSF, cerebrospinal fluid; NMC, nonmeningitis cases.

TABLE 2. Serum findings in study patients

Characteristic	Bacterial meningitis (n = 43)			Aseptic meningitis (n = 19)			NMC (n = 56)			p
	n	Mean ± SD	Range	n	Mean ± SD	Range	n	Mean ± SD	Range	
CSF RBC	29	3178 ± 15753	0–85000	8	587 ± 1386	1–4000	38	108 ± 304	0–1500	0.44
ESR (mm/h)	25	33.8 ± 37.3	2–85	12	23 ± 20	5–65	29	32.5 ± 27.27	2–95	0.45
Blood WBC	36	16285 ± 15285	3300–88000	18	12365 ± 4880	4400–22000	52	6415 ± 3901	1100–18000	>0.001 ^a
Blood Poly	28	57.25 ± 19.87	23–91	16	53.68 ± 15.3	30–85	33	53.7 ± 22.3	20–95	0.76
Blood lymphocytes	28	39.6 ± 19.3	6–75	16	41.56 ± 15.3	14–70	32	41.5 ± 21	5–80	0.91
Blood Na	34	134.2 ± 2.7	129–139	12	134.2 ± 2.8	130–139	33	135.8 ± 7	125–165	0.37
Blood K	34	4.6 ± 0.76	3–7	12	4.5 ± 0.37	4–5	32	4.1 ± 1.2	0–8	0.12
CRP	28	64 ± 64.3	1–149	15	16.9 ± 12.7	1–45	28	14.64 ± 25.15	1–84	>0.001 ^a
Body temperature	43	38.3 ± 0.75	37–40	19	38.4 ± 0.84	37–40	56	37.7 ± 1	36–40	0.002

CRP, C-reactive protein; ESR, erythrocyte sedimentation rate; NMC, nonmeningitis cases; Poly, polymorphonuclear leukocytes; RBC, red blood cell; WBC, white blood cell count.
^aLeven test was statistically significant; we used Welch analysis of variance.

laboratory findings (Fig. 1). Bacterial meningitis was confirmed by the following positive tests or combination of tests: 28 PCR (seven *S. pneumoniae*, 11 *N. meningitidis*, ten *H. influenzae*), three CSF culture and PCR (two *S. pneumoniae* and one *Acinetobacter baumannii*), one blood culture and PCR (*S. pneumoniae*), one CSF microscopy examination (Gram-positive coccobacillus), one CSF microscopy examination and blood smear (Gram-negative diplococci) and one blood smear and BACTEC (Gram-negative coccobacillus). Seven meningitis cases were confirmed by a combination of clinical and CSF findings. Moreover, aseptic meningitis was confirmed using the combination of clinical and CSF findings. In addition, 67 patients (56%) were boys and 52 (44%) were girls, with a minimum age of 4 days and a maximum age of 15 years. Fever was the most frequent feature in patients

with bacterial and aseptic meningitis, with 41 (95.3%) of 43 and 18 (95%) of 19 respectively (Table 1).

Clinical features

In terms of the clinical features, body temperature, nausea/vomiting and coryza/snivel were statistically significant for differentiation from each other of bacterial meningitis, aseptic meningitis and NMC (Table 1). Furthermore, *post hoc* analysis showed that body temperature was significant for differentiation of bacterial and aseptic meningitis from NMC (Supplementary Table S2).

Serum findings

Only serum C-reactive protein (CRP) and blood white blood cell count (WBC) were significant for differentiating between

TABLE 3. CSF findings in study patients

Characteristic	Bacterial meningitis (n = 43)			Aseptic meningitis (n = 19)			p
	n	Mean ± SD	Range	n	Mean ± SD	Range	
CSF RBC	29	3178 ± 15 753	0–85 000	8	587 ± 1386	1–4000	0.65
CSF WBC (cell/mm ³)	43	1809 ± 4430	230–21 200	19	324 ± 458	20–2050	0.03 ^a
CSF poly (%)	37	64.3 ± 26.7	3–95	18	29.6 ± 21.7	1–85	>0.001
CSF lymph (%)	37	33 ± 25	2–90	18	70.3 ± 22	15–99	>0.001
CSF Pro	41	144.7 ± 131	35–670	14	54.3 ± 64	10–260	0.01
CSF Glu	41	42.6 ± 21.7	10–116	14	61.6 ± 20	39–112	0.01
CSF:blood glucose ratio	15	0.53 ± 0.34	0–1.22	5	0.37 ± 0.28	0–0.66	0.28

CSF, cerebrospinal fluid; Glu, glucose; Poly, polymorphonuclear leukocytes; Pro, protein; RBC, red blood cell; WBC, white blood cell count.
^aLeven test was statistically significant; we used Welch analysis of variance.

TABLE 4. Accuracy of CSF and serum findings for differentiation of bacterial meningitis from aseptic meningitis

Biomarker	AUC	Cutoff values	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)
CSF WBC	0.881	420	81.4	84.2	92	66
CSF Pro	0.880	79	80.5	78.6	91	57
CSF poly	0.789	50	78.4	88.9	93	66
CSF Glu	0.784	51.5	71.4	73.2	47	88
Serum CRP	0.894	32	78.3	80	88	66.4
Blood WBC	0.476	10 250	61	49	70	38
CSF WBC; Pro, Poly and serum CRP	0.977	—	100	86.4	80	100
CSF WBC and Pro, Glu, Poly and serum CRP	1	—	100	100	100	100

AUC, area under the plasma concentration vs. time curve; CRP, C-reactive protein; Glu, glucose; NPV, negative predictive value; Poly, polymorphonuclear leukocytes; PPV, positive predictive value; Pro, protein.

bacterial meningitis, aseptic meningitis and NMC (Table 2). Moreover, *post hoc* analysis indicated that serum CRP and blood WBC were significant for differentiation of bacterial and aseptic meningitis from NMC (Supplementary Tables S3 and S4).

CSF findings

As shown in Table 3, amounts of CSF WBC, and percentage of polymorphonuclear leukocytes (Poly), lymphocytes, protein (Pro) and glucose (Glu) were significant for differentiation of bacterial and aseptic meningitis from each other.

Accuracy of CSF and serum findings

Table 4 shows the biomarkers with the highest sensitivity, specificity, area under the plasma concentration vs. time curve (AUC) and cutoff values for differentiation of bacterial meningitis from aseptic meningitis. Except serum CRP, with 78.3% sensitivity and 80% specificity with a cutoff value of 32, most of

the serum findings had low sensitivity and specificity for distinguishing bacterial and aseptic meningitis from each other. However, CSF findings such as WBC, Pro, Poly and Glu had high accuracy, with 0.881, 0.880, 0.789 and 0.784 AUC respectively for differentiation of bacterial meningitis from aseptic meningitis. Logistic regression analysis showed that the presentation of four features at the same time (CSF WBC, Pro, Poly and serum CRP) revealed 100% sensitivity and 86.4% specificity (Fig. 3), and association of five features (CSF WBC, Pro, Poly, Glu and serum CRP) showed 100% sensitivity and 100% specificity for differentiation of bacterial and aseptic meningitis from each other (Fig. 4 and Table 4).

Molecular genotyping

Of 28 PCR-positive isolates, ten (35.7%) were *H. influenzae*, seven (25%) *S. pneumoniae* and 11 (40%) *N. meningitidis*. Table 5 shows the distribution of *H. influenzae*, *S. pneumoniae* and *N. meningitidis* serotypes on the basis of patient age group. Hib (60%), *S. pneumoniae* serotype 3 (28.5%) and *Neisseria meningitidis* B (NmB) (63.5%) were the most prevalent serotypes in *H. influenzae*, *S. pneumoniae* and *N. meningitidis* respectively.




FIG. 3. ROC plot, accuracy of best biomarkers for differentiation of bacterial meningitis from aseptic meningitis. Left upper corner shows accurate biomarker with high sensitivity and specificity. ROC, receiver operating characteristic.

Discussion

Meningitis is an emergency disease with a high rate of mortality and morbidity in Iran, especially in children. The molecular epidemiology of meningitis in children is unclear in Iran and is rarely performed. We aimed to characterize clinical and para-clinical features as well as the distribution of genotype/capsular types of common bacterial meningitis agents in children in Iran.

Early diagnosis of meningitis is essential to improve the prognosis of the disease. Physicians find it difficult to diagnose the various forms of meningitis and separate meningitis from other disorders with similar presentations [13,19]. PCR has a wide range of specificity (57–94%) and sensitivity (30–95%) in diagnosing the cause of meningitis [20–24]. In this study, 72% of the cases of bacterial meningitis were PCR positive, which




FIG. 4. ROC plot, accuracy of four features (CSF WBC, Pro, Poly and serum CRP) association at same time for differentiation of bacterial meningitis from aseptic meningitis. Left upper corner shows accurate biomarker with high sensitivity and specificity. CRP, C-reactive protein; CSF, cerebrospinal fluid; ROC, receiver operating characteristic; WBC, white blood cell count.

could be due to the low copy numbers of the pathogen’s nucleic acid in CSF. However, nucleic acid amplification methods have some limitations because of their high cost and the need for special laboratory facilities, which makes it particularly difficult to use in developing countries [6,13,25]. Moreover, although CSF culture is the reference standard for the diagnosis of bacterial meningitis, only 4% to 70% of meningitis cases are positive by CSF culture [10,13,21,24–27]. Our results showed that only 7% of the bacterial meningitis cases were culture positive. Because the Mofid paediatric hospital is a referral hospital, most meningitis cases are referred to it from other hospitals. Patients routinely receive antibiotic therapy before a lumbar puncture is performed; this could be a reason for the low meningitis culture-positive cases in our study. Studies have indicated that standard microbiology diagnostics have far different sensitivities for meningitis [6,13,24], which we also found in our study. A combination of clinical and laboratory features is therefore essential to diagnose patients suspected to have meningitis [28]. Previously, many studies suggested CSF analysis and clinical assessments to differentiate between the various types of meningitis [3,4,6–10,29]. However, when other similar disorders are considered, assessment of clinical signs and symptoms should be interpreted carefully because of overlapping clinical features.

Our results showed that only fever was a significant clinical feature in distinguishing bacterial and aseptic meningitis from each other. In regard to serum findings, previous studies showed that increasing the CRP level in serum could be helpful in the

TABLE 5. Distribution of *Haemophilus influenzae*, *Streptococcus pneumoniae* and *Neisseria meningitidis* genotype/capsular types based on patient age group

Organism	Genotype/capsular type	Age group				Total (n = 115)
		< 1 month (n = 13)	1 month to 1 year (n = 56)	1–7 years (n = 32)	7–16 years (n = 14)	
<i>H. influenzae</i>	Hia	0	0	0	0	0
	Hib	1	2	2	1	6 (60%)
	Hic	0	0	1	1	2 (20%)
	Hid	0	0	0	0	0
	Hie	0	0	0	0	0
	Hif	0	0	0	0	0
	Nontypeable ^a	0	2	0	0	2 (20%)
	Total	1 (10%)	4 (40%)	3 (30%)	2 (20%)	10 (100%)
<i>S. pneumoniae</i>	3	0	1	1	0	2 (28.5%)
	23F	0	1	0	0	1 (14.3%)
	19F	0	1	0	0	1 (14.3%)
	19A	1	0	0	0	1 (14.3%)
	11	0	1	0	0	1 (14.3%)
	14	0	0	0	1	1 (14.3%)
	Total	1 (14.3%)	4 (57%)	1 (14.3%)	1 (14.3%)	7
	<i>N. meningitidis</i>	NmB	2	3	1	1
NmC		0	1	1	1	3 (27%)
NmA		0	0	0	0	0
NmW135		0	0	0	0	0
NmY		0	0	0	0	0
NmX		0	0	0	0	0
Nontypeable ^a		1	0	0	0	1 (9%)
Total		3 (27%)	4 (36.5%)	2 (18%)	2 (18%)	11 (100%)

^aNontypeable refers to isolates that did not react with study primers.

differentiation of bacterial meningitis from aseptic meningitis, with 70% to 100% sensitivity and 90% to 100% specificity [6,30]. We found that a serum CRP with a cutoff value of 32 mg/ml had 78.3% sensitivity and 80% specificity for differentiation of bacterial meningitis from aseptic meningitis; however, in a similar study setting, Dashti et al. [6] reported that CRP with cutoff value of 57 mg/mL had 91% sensitivity and 10% specificity in differentiating bacterial meningitis from viral meningitis. Moreover, logistic regression analysis demonstrated that assessment of five biomarkers (CSF WBC, Pro, Poly, Glu and serum CRP) could have 1 AUC for differentiation of bacterial meningitis from viral meningitis; however, Dashti et al. reported 0.994 AUC for the association of three biomarkers (CRP and erythrocyte sedimentation rate; CSF absolute neutrophil count; and CSF lactate) in differentiating bacterial meningitis from viral meningitis [6].

The range of CSF WBC count reported in bacterial meningitis (mean, 1500–3143 cells/mL) was significantly higher than that in viral meningitis (mean, 159–539 cell/mL) [6,12,31,32]. Our results also demonstrated that the mean CSF count was 1809 cells/mL for bacterial meningitis and 324 cells/mL for aseptic meningitis. It should be noted that the marked difference in CSF WBC counts and lower cell counts may be because our study sample included immunocompromised children as well as immunocompetent children with pneumococcus and meningococcus meningitis [6,31]. The left upper corner of the receiver operating characteristic plot shows a higher sensitivity and specificity of biomarkers in distinguishing between bacterial and aseptic meningitis (Figs. 3 and 4). Consistent with previous reports, serum CRP, CSF Poly, CSF WBC and CSF Pro, in order, have the highest accuracy for bacterial meningitis (Figs. 3 and 4) [6,14,32,33].

@@Our study showed that Hib (60%), serotype 3 (28.5%) and NmB (63.5%) were the most prevalent serotypes in *H. influenzae*, *S. pneumoniae* and *N. meningitidis* respectively. Previously, several studies reported that serotypes 11, 14, 23F, 15B/15C, 18C, 19F and 19A for *S. pneumoniae*, serogroups NmB, NmC *N. meningitidis*, and type b and c *H. influenzae* represent the causative agent in approximately 70% to 85% of children with bacterial meningitis, especially in nonvaccine regions [34–38]. Therefore, the distribution of genotype/capsular types of common agents causing bacterial meningitis in children in Iran are not different from other regions of the world.

The main limitation of the present study was that the incidence of meningitis in children in Iran is less considered in this study. For future direction, additional and further well-designed studies are necessary to accurately apply new biomarkers to assess suspected cases of meningitis which will add potential value to standard laboratory tests. It is also necessary to perform comprehensive molecular epidemiology studies from different regions of Iran.

In conclusion, we demonstrated that a well-designed combination of clinical and paraclinical features could be useful to diagnose bacterial meningitis, but these features are not good enough to be relied on as stand-alone, exclusionary tests. In addition, public immunization of infants with the most prevalent bacterial meningitis serotypes is recommended.

Conflict of Interest

None declared.

Acknowledgements

The research reported herein was extracted from the PhD thesis written by Ali Pormohammad, School of Medicine, Shahid Beheshti University of Medical Sciences (Registration 191). Supported in part by the Shahid Beheshti University of Medical Sciences.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.nmni.2019.100594>.

References

- [1] Van de Beek D, Cabellos C, Dzupova O, Esposito S, Klein M, Kloek A, et al. ESCMID guideline: diagnosis and treatment of acute bacterial meningitis. *Clin Microbiol Infect* 2016;22:S37–62.
- [2] Pormohammad A, Abasi E, Azimi T, Hosseini S, Safari H, Nasiri MJ, et al. Global estimate of *Neisseria meningitidis* serogroups proportion in invasive meningococcal disease: a systematic review and meta-analysis. *Microb Pathog* 2019;134:103571.
- [3] Nazir M, Wani WA, Malik MA, Mir MR, Ashraf Y, Kawoosa K, et al. Cerebrospinal fluid lactate: a differential biomarker for bacterial and viral meningitis in children. *J Pediatr (Rio J)* 2018;94:88–92.
- [4] Tan J, Kan J, Qiu G, Zhao D, Ren F, Luo Z, et al. Clinical prognosis in neonatal bacterial meningitis: the role of cerebrospinal fluid protein. *PLoS One* 2015;10:e0141620.
- [5] Martin NG, Iro MA, Sadarangani M, Goldacre R, Pollard AJ, Goldacre MJ. Hospital admissions for viral meningitis in children in England over five decades: a population-based observational study. *Lancet Infect Dis* 2016;16:1279–87.
- [6] Dashti AS, Alizadeh S, Karimi A, Khalifeh M, Shoja SA. Diagnostic value of lactate, procalcitonin, ferritin, serum-C-reactive protein, and other biomarkers in bacterial and viral meningitis: a cross-sectional study. *Medicine (Baltimore)* 2017;96:e7637.
- [7] Pormohammad A, Riahi SM, Nasiri MJ, Fallah F, Aghazadeh M, Doustdar F, et al. Diagnostic test accuracy of adenosine deaminase for tuberculous meningitis: a systematic review and meta-analysis. *J Infect* 2017;74:545–54.

- [8] McGill F, Griffiths MJ, Solomon T. Viral meningitis: current issues in diagnosis and treatment. *Curr Opin Infect Dis* 2017;30:248–56.
- [9] Glimåker M, Sjölin J, Åkesson S, Naucler P. Lumbar puncture performed promptly or after neuroimaging in acute bacterial meningitis in adults: a prospective national cohort study evaluating different guidelines. *Clin Infect Dis* 2017;66:321–8.
- [10] Van de Beek D, De Gans J, Spanjaard L, Weisfelt M, Reitsma JB, Vermeulen M. Clinical features and prognostic factors in adults with bacterial meningitis. *N Engl J Med* 2004;351:1849–59.
- [11] Zimmerli W. How to differentiate bacterial from viral meningitis. Amsterdam: Springer; 2005.
- [12] Hourri H, Pormohammad A, Riahi SM, Nasiri MJ, Fallah F, Dabiri H, et al. Acute bacterial meningitis in Iran: systematic review and meta-analysis. *PLoS One* 2017;12:e0169617.
- [13] Brouwer MC, Thwaites GE, Tunkel AR, van de Beek D. Dilemmas in the diagnosis of acute community-acquired bacterial meningitis. *Lancet* 2012;380(9854):1684–92.
- [14] Scarborough M, Thwaites GE. The diagnosis and management of acute bacterial meningitis in resource-poor settings. *Lancet Neurol* 2008;7:637–48.
- [15] World Health Organization; US Centers for Disease Control and Prevention. Laboratory methods for the diagnosis of meningitis caused by *Neisseria meningitidis*, *Streptococcus pneumoniae*, and *Haemophilus influenzae*. WHO manual. 2nd ed. Geneva: World Health Organization; 2011 Available at: https://www.who.int/csr/resources/publications/meningitis/WHO_CDS_CSR_EDC_99_7_EN/en/.
- [16] Wang X, Theodore MJ, Mair R, Trujillo-Lopez E, du Plessis M, Wolter N, et al. Clinical validation of multiplex real-time PCR assays for detection of bacterial meningitis pathogens. *J Clin Microbiol* 2012;50:702–8.
- [17] Tzanakaki G, Tsopanomalou M, Kesanopoulos K, Matzourani R, Sioumala M, Tabaki A, et al. Simultaneous single-tube PCR assay for the detection of *Neisseria meningitidis*, *Haemophilus influenzae* type b and *Streptococcus pneumoniae*. *Clin Microbiol Infect* 2005;11:386–90.
- [18] Pai R, Gertz RE, Beall B. Sequential multiplex PCR approach for determining capsular serotypes of *Streptococcus pneumoniae* isolates. *J Clin Microbiol* 2006;44:124–31.
- [19] Agarwal AK, Bansal S, Nand V. A hospital based study on estimation of adenosine deaminase activity (ADA) in cerebrospinal fluid (CSF) in various types of meningitis. *J Clin Diagn Res* 2014;8:73.
- [20] Abdeldaim GM, Strålin K, Korsgaard J, Blomberg J, Welinder-Olsson C, Herrmann B. Multiplex quantitative PCR for detection of lower respiratory tract infection and meningitis caused by *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Neisseria meningitidis*. *BMC Microbiol* 2010;10:310.
- [21] Rafi W, Chandramuki A, Mani R, Satishchandra P, Shankar SK. Rapid diagnosis of acute bacterial meningitis: role of a broad range 16S rRNA polymerase chain reaction. *J Emerg Med* 2010;38:225–30.
- [22] Rath PM, Schoch B, Adamzik M, Steinmann E, Buer J, Steinmann J. Value of multiplex PCR using cerebrospinal fluid for the diagnosis of ventriculostomy-related meningitis in neurosurgery patients. *Infection* 2014;42:621–7.
- [23] Rothman R, Ramachandran P, Yang S, Hardick A, Wwon H, Kecojevic A, et al. Use of quantitative broad-based polymerase chain reaction for detection and identification of common bacterial pathogens in cerebrospinal fluid. *Acad Emerg Med* 2010;17:741–7.
- [24] Sarookhani M-R, Ayazi P, Alizadeh S, Foroughi F, Sahmani A, Adineh M. Comparison of 16S rDNA-PCR amplification and culture of cerebrospinal fluid for diagnosis of bacterial meningitis. *Iran J Pediatr* 2010;20:471.
- [25] Ghotaslou R, Farajnia S, Yeganeh F, Abdoli-Oskouei S, Rezaee MA, Barzegar M. Detection of acute childhood meningitis by PCR, culture and agglutination tests in Tabriz, Iran. *Acta Med Iran* 2012;50:192.
- [26] Wagner K, Springer B, Pires VP, Keller PM. Pathogen identification by multiplex LightMix real-time PCR assay in patients with meningitis and culture-negative cerebrospinal fluid specimens. *J Clin Microbiol* 2018;56. e01492-17.
- [27] Morrissey S, Nielsen M, Ryan L, Al Dhanhani H, Meehan M, McDermott S, et al. Group B streptococcal PCR testing in comparison to culture for diagnosis of late onset bacteraemia and meningitis in infants aged 7–90 days: a multi-centre diagnostic accuracy study. *Eur J Clin Microbiol Infect Dis* 2017;36:1317–24.
- [28] Waghdhare S, Kalantri A, Joshi R, Kalantri S. Accuracy of physical signs for detecting meningitis: a hospital-based diagnostic accuracy study. *Clin Neurol Neurosurg* 2010;112:752–7.
- [29] Jebamalar AA, Prabhat, Balakrishnapillai AK, Parmeswaran N, Dhiman P, Rajendiran S. Cerebrospinal fluid ferritin and albumin index: potential candidates for scoring system to differentiate between bacterial and viral meningitis in children. *Biomarkers* 2016;21:424–8.
- [30] Shimetani K, Shimetani M, Mori N. Levels of three inflammation markers, C-reactive protein, serum amyloid A protein and procalcitonin, in the serum and cerebrospinal fluid of patients with meningitis. *Scand J Clin Lab Invest* 2001;61:567–74.
- [31] Águeda S, Campos T, Maia A. Prediction of bacterial meningitis based on cerebrospinal fluid pleocytosis in children. *Braz J Infect Dis* 2013;17:401–4.
- [32] Bonsu BK, Harper MB. Differentiating acute bacterial meningitis from acute viral meningitis among children with cerebrospinal fluid pleocytosis: a multivariable regression model. *Pediatr Infect Dis J* 2004;23:511–7.
- [33] Negri B, Kelleher KJ, Wald ER. Cerebrospinal fluid findings in aseptic versus bacterial meningitis. *Pediatrics* 2000;105:316–9.
- [34] Stein-Zamir C, Abramson N, Zentner G, Shoob H, Valinsky L, Block C. Invasive meningococcal disease in children in Jerusalem. *Epidemiol Infect* 2008;136:782–9.
- [35] Jafri RZ, Ali A, Messonnier NE, Tevi-Benissan C, Durrheim D, Eskola J, et al. Global epidemiology of invasive meningococcal disease. *Popul Health Metr* 2013;11:17.
- [36] Sridhar S, Greenwood B, Head C, Plotkin SA, Sáfiadi MA, Saha S, et al. Global incidence of serogroup B invasive meningococcal disease: a systematic review. *Lancet Infect Dis* 2015;15:1334–46.
- [37] Nuorti JP, Whitney CG, Centers for Disease Control and Prevention (CDC). Prevention of pneumococcal disease among infants and children: use of 13-valent pneumococcal conjugate vaccine and 23-valent pneumococcal polysaccharide vaccine: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep* 2010;59(RR-11):1–18.
- [38] Monge S, Hahné SJ, de Melker HE, Sanders EA, van der Ende A, Knol MJ. Effectiveness of the DTPa-HBV-IPV/Hib vaccine against invasive *Haemophilus influenzae* type b disease in The Netherlands (2003–16): a case-control study. *Lancet Infect Dis* 2018;18:749–57.