



Identification and genotyping of a new subtype of bovine viral diarrhea virus 1 isolated from cattle with diarrhea

Bin Tian¹ · Dongjie Cai² · Weiqiang Li³ · Qinglong Bu³ · Mingshu Wang¹ · Gang Ye³ · Jianzhu Liu⁴ · Ya Wang² · Liping Gou² · Jun Yi⁵ · Zhicai Zuo²

Received: 16 September 2020 / Accepted: 21 December 2020 / Published online: 13 February 2021
© The Author(s) 2021

Abstract

In 2019, diarrhea cases occurred on cattle farms in Qionglai and Guang'an, Sichuan Province. Two out of 20 (10%) serum and nasal swab samples were positive when tested using a bovine viral diarrhea virus (BVDV) antigen-capture ELISA kit. Two non-cytopathic strains of BVDV were isolated and named QL1903 and GA190608, respectively. The nucleotide sequences of the genomes of the two isolates were 89.52% identical. Phylogenetic analysis based on the 5'-UTR sequence revealed that the BVDV isolate QL1903 belonged to BVDV subtype 1b, whereas isolate GA190608 clustered with strains HN1814, EN-19, and BJ09_26 in a separate branch, which has tentatively been classified as a new genetic subtype, "1v".

Bovine viral diarrhea virus (BVDV) is an immunosuppressive virus that is widely distributed throughout the world and poses an economic risk to the cattle industry [1]. It is responsible for a wide spectrum of clinical syndromes in cattle, including respiratory disease, reproductive dysfunction, immunosuppression, persistent infection, and mucosal disease [2]. BVDV, border disease virus (BDV) and classical swine fever virus (CSFV) belong to the genus *Pestivirus* within the family *Flaviviridae* [3]. Based on antigenic

and genetic differences, two genotypes of BVDV have been identified: BVDV-1 and BVDV-2. Another group, referred to as "BVDV-3" or "atypical pestiviruses", whose members are quite different from the above two genotypes, includes the strains Th/04_KhonKaen, D32/00-'HoBi', and others [4–6]. In addition, based on their ability to cause a cytopathic effect, BVDV-1, BVDV-2, and BVDV-3 isolates are categorized as either cytopathogenic (CP) or noncytopathogenic (NCP) [7, 8]. So far, 21 subgenotypes (BVDV-1a to BVDV-1u) have been reported worldwide [9, 10]. The BVDV genome is about 12.3 kb in length and comprises a 5' untranslated region (UTR), a 3'-UTR, and one open reading frame [11]. Due to the highly conserved nature of the 5'-UTR, the sequence of this region is widely used to determine the species and genotype of new pestivirus strains [12].

In March and June of 2019, calves on two cattle farms in Qionglai and Guang'an, Sichuan Province, developed clinical signs including diarrhea, anorexia, blood and intestinal mucosa in feces, and emaciation. Their condition did not improve significantly after antibiotic treatment, and they were clinically diagnosed as having a suspected BVDV infection. In order to determine the cause of diarrhea, 20 samples (10 serum samples and 10 nasal swabs) were collected from sick calves. The clinical manifestations of severe diarrhea in sick calves are shown in Fig. S1. All of the samples were examined for the presence of the viral E^{rns} glycoprotein using a BVDV Antigen Test Kit (IDEXX Laboratories), and two of the 20 samples were confirmed to be positive. The two BVDV-positive samples were processed

Handling Editor: Tim Skern.

Bin Tian, Dongjie Cai and Weiqiang Li contributed equally to this work.

✉ Zhicai Zuo
zzcjl@126.com

¹ Institute of Preventive Veterinary Medicine, Sichuan Agricultural University, Chengdu 611130, China

² Department of Veterinary Clinical Sciences, College of Veterinary Medicine, Sichuan Agricultural University, Chengdu 611130, China

³ Ministry of agriculture "Veterinary medicine and veterinary technology" Sichuan scientific experimental observation station, Sichuan Agricultural University, Chengdu 611130, China

⁴ College of Veterinary Medicine, Shandong Agricultural University, Tai'an 271018, China

⁵ Sichuan Academy of Animal Science, Chengdu 610066, China

for analysis by RT-PCR. Viral RNA was isolated using an EasyPure Viral DNA/RNA Kit (TransGen Biotech, China). *Trans Taq* HiFi PCR SuperMix (TransGen Biotech, China) and the primers 324 and 326 [13] were used for 5'-UTR amplification. This resulted in bright positive bands migrating above a 250-bp marker band (Fig. S2). A 288-bp region corresponding to the 5-UTR was sequenced by Sangon Biotech (Shanghai).

In order to isolate the virus and identify its biotype, the BVDV-positive samples were inoculated onto a monolayer of Madin-Darby bovine kidney (MDBK) cells in DMEM (Gibco, Life Technologies, USA) containing fetal bovine serum (FBS; Hyclone, Life Technologies USA) without bovine viral diarrhea virus antibodies or antigens. Cells were cultured at 37°C with 5% CO₂, and morphological changes in the cells were monitored each day. After one week of incubation, the lysates of the third-generation cells were collected to detect BVDV by RT-PCR. Two isolates (QL1903 and GA190608) were obtained that could be passaged stably in MDBK cells. QL1903 was isolated from a nasal swab sample from a calf in Qionglai. GA190608 was isolated from a serum sample from a calf in Guang'an. No cytopathic effect was observed in MDBK cells after inoculation with QL1903 or GA190608, suggesting that these two BVDV isolates belong to the NCP biotype. An indirect immunofluorescence assay performed as described previously [14] showed green staining in the cytoplasm of cells infected with these isolates (Fig. 1).

To investigate the evolutionary relationship of the new isolates to BVDV isolates, phylogenetic analysis was

performed based on 5'-UTR sequences available in the GenBank database. Nucleotide sequence comparisons were performed using NCBI BLAST online. Alignment and manual editing were performed using Molecular Evolutionary Genetics Analysis (MEGA) software, version 7. All of the sequences were aligned with BVDV reference strains retrieved from GenBank, including BVDV-1, BVDV-2, and BVDV-3, as well as CSFV and BDV. Phylogenetic trees were constructed using MEGA software as described previously [15]. The 5'-UTR sequence of the QL1903, and GA190608 isolates have been deposited in the GenBank database under accession numbers MN849041 and MT933204.

Phylogenetic analysis showed that the isolates QL1903 and GA190608 clustered with BVDV-1 isolates (Fig. 2A). The nucleotide sequences of the two isolates were 89.52% identical to each other. The QL1903 isolate shared 94.72%–98.78% nucleotide sequence identity with the other BVDV-1 strains. QL1903 clustered with strains CC13B and HJ-1 in a branch (Fig. 2B) representing subgenotype 1b. QL1903 shared 94.09% and 95.93% nucleotide sequence identity with strains CC13B and HJ-1, respectively. QL1903 shared the highest nucleotide sequence identity (95.93%) with strain HJ-1 (KU756226), which was isolated in Heilongjiang Province in 2010. In the past years, BVDV subtype 1b has been detected frequently in Chinese bovine and porcine species [16, 17]. GA190608 shared 89.52%–98.37% nucleotide sequence identity with the other BVDV-1 strains and clustered with HN1814, EN-19, and BJ09_26 in a branch of the phylogenetic tree (Fig. 2B) that has not been assigned a genotype. GA190608 shared 98.67%, 98.76%,




Fig. 1 Immunofluorescence staining of BVDV in MDBK cells. The control group was not inoculated with BVDV-positive samples. The QL1903 group was inoculated with BVDV QL1903 samples. The

GA190608 group was inoculated with BVDV GA190608 samples. BVDV antigens in the cytoplasm were stained green. Scale bar, 100 μ m




Fig. 2 Phylogenetic tree constructed by the neighbor-joining method based on partial nucleotide sequences of the BVDV 5'-UTR. The QL1903 and GA190608 isolates are indicated by a black triangle.

and 99.56% nucleotide sequence identity with HN1814, EN-19, and BJ09_26, respectively. GA190608 shared the highest nucleotide sequence identity (99.56%) with strain BJ09_26. According to data from GenBank, strain HN1814 was isolated from cattle in Henan Province in January 2020, EN-19 strain was isolated from ear notch samples from dairy cattle in China in 2017, and strain BJ09_26 (HQ116551) was isolated in Beijing in 2009. Since these three BVDV strains have not been assigned a subgenotype, we tentatively labelled the branch containing GA190608, HN1814, EN-19, and BJ09_26 as "BVDV-1v". Although these strains have not been assigned a genotype designation, they have been circulating recently in different areas of China. This may be related to the rapid development of the cattle industry and the transport of cattle from various places. This study is the first to report the emergence of the new subgenotype BVDV-1v, which can enrich the BVDV strains database and provide early warning to cattle disease control agencies.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00705-021-04990-7>.

Funding This work was supported by the Program for the National Key Research and Development Program of China (Grant/Award Number: 2018YFD0501800); Sichuan Science and Technology Program (Grant/Award Number: 2018NZ0002/2019YFQ0012); and Sichuan

Another 51 sequences from pestiviruses reference strains (BVDV-1, BVDV-1, BVDV-3, CSFV, BDV) were retrieved from the GenBank database.

Beef Cattle Innovation Team of National Modern Agricultural Industry Technology System, (Grant/Award Number: 035Z2036).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethics statement Bovine blood and nasal swab sampling were conducted as a part of disease diagnosis based on the domestic animal infectious disease control law in China, and the samples were permitted to be used in this study by the cattleman.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ridpath JF (2010) Bovine viral diarrhoea virus: global status. *Vet Clin N Am Food Anim Pract* 26(1):105–121. <https://doi.org/10.1016/j.cvfa.2009.10.007>
- Blome S, Beer M, Wernike K (2017) New leaves in the growing tree of pestiviruses. *Adv Virus Res* 99:139–160. <https://doi.org/10.1016/bs.aivir.2017.07.003>
- Smith DB, Meyers G, Bukh J, Gould EA, Becher P (2017) Proposed revision to the taxonomy of the genus Pestivirus, family Flaviviridae. *J Gen Virol*. <https://doi.org/10.1099/jgv.0.000873>
- Liu L, Xia H, Wahlberg N, Belák S, Baule C (2009) Phylogeny, classification and evolutionary insights into pestiviruses. *Virology* 385(2):351–357. <https://doi.org/10.1016/j.virol.2008.12.004>
- Mehmet ÖT, Hakan A (2019) Increased genetic diversity of bvdv strains circulating in eastern anatolia, turkey: first detection of bvdv-3 in turkey. *Tro Anim Health Pro* 51:1953–1961. <https://doi.org/10.1007/s11250-019-01901-6>
- Bauermann FV, Ridpath JF (2015) HoBi-like viruses—the typical 'atypical bovine pestivirus. *Anim Health Res Rev*. <https://doi.org/10.1017/S146625231500002X>
- Moirá B, Luigi B, Monica G, Mario DMG (2018) Complete genome sequences of a cytophatic/noncytophatic pair of bovine viral diarrhoea virus subtype 1a viruses. *Arch Virol* 163:1–2. <https://doi.org/10.1007/s00705-018-3959-7>
- Colitti B, Nogarol C, Giacobini M, Capucchio MT, Bertolotti L (2019) Compartmentalized evolution of Bovine Viral Diarrhoea Virus type 2 in an immunotolerant persistently infected cow. *Sci Rep* 9(1):15460. <https://doi.org/10.1038/s41598-019-52023-w>
- Donoso A, Inostroza F, Celedón M, Pizarro-Lucero J (2018) Genetic diversity of Bovine Viral Diarrhoea Virus from cattle in Chile between 2003 and 2007. *BMC Vet Res* 14(1):314. <https://doi.org/10.1186/s12917-018-1641-7>
- Kadir YI, Gizem A, Paul B (2017) Variability and global distribution of subgenotypes of bovine viral diarrhoea virus. *Viruses* 9(6):128. <https://doi.org/10.3390/v9060128>
- Lanyon S, Hill F, Reichel M, Brownlie J (2014) Bovine viral diarrhoea: pathogenesis and diagnosis. *Vet J* 199(2):201–209. <https://doi.org/10.1016/j.tvjl.2013.07.024>
- Giangaspero M, Harasawa R (2007) Numerical taxonomy of the genus pestivirus based on palindromic nucleotide substitutions in the 5' untranslated region. *J Virol Methods* 146(1–2):375–388. <https://doi.org/10.1016/j.jviromet.2007.07.009>
- Factor C, Yus E, Eiras C, Sanjuan ML, Cerviño M, Arnaiz I, Diéguez FJ (2016) Genetic diversity of bovine viral diarrhoea viruses from the Galicia region of Spain. *Vet Record Open* 3(1):e000196. <https://doi.org/10.1136/vetreco-2016-000196>
- Cai D, Song Q, Duan C, Wang S, Wang J, Zhu Y (2018) Enhanced immune responses to E2 protein and DNA formulated with ISA 61 VG administered as a DNA prime–protein boost regimen against bovine viral diarrhoea virus. *Vaccine* 36(37):5591–5599. <https://doi.org/10.1016/j.vaccine.2018.07.054>
- Cai D, Song Q, Wang J, Zhu Y (2016) Genomic characterization of three bovine viral diarrhoea virus isolates from cattle. *Arch Virol* 161(12):1–4. <https://doi.org/10.1007/s00705-016-3055-9>
- Deng M, Ji S, Fei W, Sohail R, He C, Chen Y, Chen H, Guo A, Zhou D (2015) Prevalence study and genetic typing of bovine viral diarrhoea virus (BVDV) in four bovine species in China. *PLoS ONE* 10(4):e0121718. <https://doi.org/10.1371/journal.pone.0121718>
- Deng Y, Sun CQ, Cao SJ, Lin T, Yuan SS, Zhang HB, Zhai SL, Huang L, Shan TL, Zheng H (2012) High prevalence of bovine viral diarrhoea virus 1 in Chinese swine herds. *Vet Microbiol* 159:3–4. <https://doi.org/10.1016/j.vetmic.2012.04.023>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.