

POSTER PRESENTATION

Open Access

A double blind clinical trial evaluating the relative pharmacokinetics and bioavailability of oral creatine monohydrate when combined with either isomaltulose or dextrose in healthy adult males

Douglas S Kalman*, Samantha Feldman, Howard I Schwartz, Diane R Krieger

From International Society of Sports Nutrition: 9th Annual ISSN Conference and Expo Clearwater, FL, USA. 22-23 June 2012

Background

Isomaltulose (6-0- α -D-glucopyranosyl-D-fructose) is a low-glycemic, low-insulinemic disaccharide that is absorbed more slowly than conventional sugars (monosaccharides). In sports nutrition, creatine monohydrate is often combined with dextrose (a monosaccharide) for the purpose of enhanced absorption and cellular uptake.

Methods

In a prospective, randomized, double blind, active-comparator-controlled, parallel group pilot study, 30 male subjects, age 27.0 ± 4.6 years, with BMI of 24.75 ± 1.99 kg/m² and a body surface area (BSA) of 1.953 ± 0.075 m², were randomly assigned to ingest 3 grams of creatine monohydrate (CM) in combination with isomaltulose (ISO) or dextrose (DEX) in 1 of 3 concentrations (5 gm liquid, 17 gm capsules or 50 gm liquid). Rate of absorption (t_{Max}) and overall absorption (from BSA adjusted AUC_{0-8h} and C_{Max}) of CM was determined via changes in serum creatine over an 8-hour test period. Blood was collected at baseline and 0.5, 1, 2.5, 4 and 8 hours post ingestion with efficacy endpoints including C_{Max} , t_{Max} , AUC_{0-8h} and λ_{Elim} derived from normalized concentration vs. time curves for serum creatine (AUC by trapezoidal integration). Serum creatine levels were normalized by BSA using the Mosteller formula. For PK parameters, paired Student t test (or Wilcoxon if non-

normally distributed) was used and for categorical variables, Fisher Exact test (or Chi-Square if necessary) was used. Statistics were calculated by R v2.14.0 (www.r-project.org).

Results

For the 17 gm concentrations, ISO had a significantly higher C_{Max} than DEX (18.1 ± 1.5 vs 12 ± 1.6 mg/dl*m²; $p < 0.001$) and for the 50 gm concentrations, the C_{Max} trended higher for ISO than DEX (19.1 ± 6.4 vs 13.1 ± 3.3 mg/dl*m²; $p = 0.099$). The AUC for the 50 gm concentration was significantly higher for ISO than DEX (54.6 ± 9.2 vs 40.3 ± 10 ; $p = 0.046$). The 17 gm (1.9 ± 0.8 hrs) and 50 gm (1.3 ± 0.7 hrs) concentrations were associated with larger t_{Max} , which trended toward significance over the 5 gm concentration (1 ± 0 hrs) for ISO ($p = 0.078$) and was not significant for DEX. For all 3 concentrations, the C_{Max} and AUC were significantly higher for ISO than DEX (17.8 ± 4.7 vs 13.5 ± 2.8 mg/dl*m² and 50.8 ± 17.1 vs 38.8 ± 10.3 ; $p = 0.005$ and $p = 0.027$ respectively).

Conclusions

CM appears to be absorbed more efficiently when combined with ISO over DEX supported by a significantly higher C_{max} for the 17 g concentration and a significantly higher AUC for the 50 g concentration. The 17 and 50 gm formulations appear to be superior to the 5 gm concentration. ISO appears to be a beneficial carbohydrate for facilitating the delivery of creatine to the body.

* Correspondence: dkalman@miamiresearch.com
Miami Research Associates, 6141 Sunset Drive Suite 301, Miami, FL 33143, USA

Acknowledgements

Hong Kong Life Sciences Company Limited. Wanchai, Hong Kong.

Published: 19 November 2012

doi:10.1186/1550-2783-9-S1-P14

Cite this article as: Kalman *et al.*: A double blind clinical trial evaluating the relative pharmacokinetics and bioavailability of oral creatine monohydrate when combined with either isomaltulose or dextrose in healthy adult males. *Journal of the International Society of Sports Nutrition* 2012 **9**(Suppl 1):P14.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

