



OPEN ACCESS

REVIEW

A unifying theory for cognitive abnormalities in functional neurological disorders, fibromyalgia and chronic fatigue syndrome: systematic review

Tiago Teodoro,^{1,2,3} Mark J Edwards,^{1,2} Jeremy D Isaacs^{1,2}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jnnp-2017-317823>).

¹Neurosciences Research Centre, Molecular and Clinical Sciences Research Institute, St George's, University of London, London, UK

²Department of Neurology, St George's University Hospitals NHS Foundation Trust, London, UK

³Instituto de Medicina Molecular Faculdade de Medicina, Universidade de Lisboa & Serviço de Neurologia Hospital de Santa Maria, Lisboa, Portugal

Correspondence to

Dr Jeremy D Isaacs, Department of Neurology, St George's Hospital, London SW17 0QT, UK; jeremy.isaacs@nhs.net

Received 12 December 2017

Revised 24 March 2018

Accepted 10 April 2018

Published Online First 7 May 2018

ABSTRACT

Background Functional cognitive disorder (FCD) describes cognitive dysfunction in the absence of an organic cause. It is increasingly prevalent in healthcare settings yet its key neuropsychological features have not been reported in large patient cohorts. We hypothesised that cognitive profiles in fibromyalgia (FM), chronic fatigue syndrome (CFS) and functional neurological disorders (FNDs) would provide a template for characterising FCD.

Methods We conducted a systematic review of studies with cognition-related outcomes in FM, CFS and FND.

Results We selected 52 studies on FM, 95 on CFS and 39 on FND. We found a general discordance between high rates of subjective cognitive symptoms, including forgetfulness, distractibility and word-finding difficulties, and inconsistent objective neuropsychological deficits. Objective deficits were reported, including poor selective and divided attention, slow information processing and vulnerability to distraction. In some studies, cognitive performance was inversely correlated with pain, exertion and fatigue. Performance validity testing demonstrated poor effort in only a minority of subjects, and patients with CFS showed a heightened perception of effort.

Discussion The cognitive profiles of FM, CFS and non-cognitive FND are similar to the proposed features of FCD, suggesting common mechanistic underpinnings. Similar findings have been reported in patients with mild traumatic brain injury and whiplash. We hypothesise that pain, fatigue and excessive interoceptive monitoring produce a decrease in externally directed attention. This increases susceptibility to distraction and slows information processing, interfering with cognitive function, in particular multitasking. Routine cognitive processes are experienced as unduly effortful. This may reflect a switch from an automatic to a less efficient controlled or explicit cognitive mode, a mechanism that has also been proposed for impaired motor control in FND. These experiences might then be overinterpreted due to memory perfectionism and heightened self-monitoring of cognitive performance.

INTRODUCTION

Practitioners in several settings report an increase in patients presenting with non-dementia cognitive symptoms in recent years.^{1–3} Notably, a range of benign cognitive symptoms are experienced by between 5% and 32% of healthy young adults.⁴ The term functional cognitive disorder (FCD) (or functional memory disorder) has been proposed

to describe the subjective experience of cognitive dysfunction in the absence of underlying brain pathology.^{5–8} In common with historical practice in other functional neurological disorders (FNDs), the approach of cognitive specialists towards FCD has largely been to ‘rule out’ organic causes, rather than making an affirmative diagnosis on the basis of distinctive clinical features.⁹

Typical symptoms of FCD include forgetting tasks while preparing to execute them, inability to retrieve overlearned memories, disrupted flow of thoughts and conversations, prospective memory failures and absent-mindedness.^{5–9} These burdensome symptoms are believed to occur in the absence of relevant objective deficits on neuropsychological assessment, that is, a discrepancy between subjective experience and objective performance, which is a canonical feature of FNDs.^{8,10} However, neuropsychological data on large cohorts of patients with FCD have not been reported. FCD is not a trivial disorder, with many patients continuing to experience distressing cognitive symptoms for months to years.⁸

Theoretically, FCD is proposed to be underpinned by psychological factors such as memory perfectionism, overinterpretation of attentional lapses and heightened self-monitoring for cognitive errors.^{5–6,11} However, the empirical evidence supporting this model is modest.

FCD is considered part of the spectrum of FND (also known as conversion disorder).⁶ FNDs in general are characterised by both ‘internal’ inconsistency and incongruity with other neurological disorders (‘external’ inconsistency).¹² With respect to FCD, internal inconsistency can be demonstrated, for example, by patients being able to provide detailed accounts of their memory lapses, while distinctive phenomenological differences with organic cognitive disorders constitute external inconsistency.¹³

People with FND most commonly present with abnormalities of movement control (eg, weakness, tremor and dystonic posturing) or attacks that resemble epilepsy (dissociative seizures/non-epileptic attacks). Theories regarding the underlying neurobiology of FND suggest a central role for misdirected attention towards movement/sensation.¹⁴ Of note, people with FND presenting with movement disorder or dissociative seizures commonly report cognitive difficulties.⁹

There is ongoing debate regarding the relevance of the growing pathophysiological understanding of

Check for updates

To cite: Teodoro T, Edwards MJ, Isaacs JD. *J Neurol Neurosurg Psychiatry* 2018;**89**:1308–1319.

FND to people who have chronic pain or chronic fatigue, who are typically diagnosed with fibromyalgia (FM) or chronic fatigue syndrome (CFS). FM and CFS can be regarded as functional somatic syndromes, a group of conditions affecting different body systems, but which are argued to have overlapping clinical and aetiological characteristics.¹⁵ In our clinical practice, it is very common to see patients with FND and comorbid FM and/or CFS. Importantly, FM and CFS are also associated with subjective cognitive difficulties, sometimes referred to as 'brain fog'.^{6 16–18}

The relationship between archetypal FCD and cognitive symptoms in the context of FM, CFS and FND remains unclear. Furthermore, the extent to which cognitive difficulties in these conditions are separate or overlapping disorders has not previously been studied. However, the extensive literature on cognitive symptomatology in these conditions affords an opportunity to compare them and to consider whether the cognitive symptoms of FND, FM, and CFS provide a template for understanding the emerging concept of FCD.

We conducted a systematic review of cognitive findings in patients with FM, CFS and FND, in order to define their key neuropsychological characteristics. Our hypothesis was that the cognitive profiles associated with functional motor, fatigue and pain syndromes would be similar to each other and to that proposed for isolated FCD. This would support the existence of a shared causal mechanism contributing to cognitive symptoms in these conditions with important implications for diagnosis and treatment.

METHODS

We performed a systematic review of studies with cognition-related outcomes in FND, FM and CFS. The protocol was registered with the PROSPERO database (CRD42017072044). Outcomes included self-reported cognitive difficulties and deficits on objective neuropsychological assessment.

More specifically, we planned to analyse and compare evidence regarding: cognitive symptoms; objective neuropsychological abnormalities (memory, attention and executive functions, information processing, language and social cognition); consistency on repeat assessment; performance validity testing (PVT); and concordance between self-reported symptoms and objective cognitive deficits. These domains were defined based on our aims, previous knowledge and preliminary review of the literature, including relevant review articles.

We searched PubMed and hand-searched reference lists for articles published in English language. JDI, MJE and TT defined search strategy and selection criteria. TT searched PubMed, hand-searched relevant reference lists and selected studies for inclusion in the final qualitative analysis. Further details regarding study identification (including 'search terms') can be found in the online (supplementary PRISMA diagrams for FND, FM and CFS studies).

Study quality was assessed based on: use of a control group, inclusion of PVT and blinding of neuropsychological assessment. A summary of the methodology and main findings of each selected study for each condition can also be found in the online (supplementary summary tables from FND, FM and studies).

RESULTS: EVIDENCE FOR ABNORMAL COGNITION IN FM, CFS AND FND

For our final qualitative analysis (table 1), we selected 39 studies on FND including functional movement disorders (FMDs) and non-epileptic attacks (NEAs), 52 studies on FM and 95

on CFS. Further details regarding study selection, including PRISMA flow diagrams, can be found in the online supplementary materials. Study characteristics were highly heterogeneous. In general, the evidence was scarcer for FND, including only one study assessing the neuropsychological profile of patients with FMD.¹⁹ We characterise each study in more detail in the supplementary materials (supplementary summary tables from FND, FM and studies). Here we present a summary of the main findings, organised by topic and condition (table 1).

Cognitive symptoms

Fibromyalgia

Patients with FM reported cognitive difficulties including forgetfulness, distractibility, speech/language difficulties and disorganised thinking.^{20–24} The proportion of patients reporting cognitive symptoms ranged from around 50% to 90%.^{20 23 24} This prevalence was higher than in healthy controls²¹ or controls with rheumatic disorders.^{22–24} Contributing factors included pain, fatigue, depressed mood and poor sleep.²²

Chronic fatigue syndrome

Patients with CFS often experienced cognitive symptoms, including poor memory, impaired attention and word-finding and reasoning difficulties.^{25–39} The proportion of affected patients ranged from around 70 up to 90%.^{25–28} Cognitive difficulties were more prevalent than in healthy subjects.^{24 27} Coexisting fatigue,^{28 38 40} depression,^{26 28 38 40 41} anxiety^{26 31 38 41} and pain²⁹ contributed to but did not entirely account for the severity of cognitive symptoms.²⁶

Functional neurological disorder

A relatively small number of studies have investigated cognitive symptoms in patients with FND (including FMD¹⁹ and NEA^{42–45}), with only one¹⁹ enrolling healthy controls. Patients with FMD reported more cognitive complaints than healthy subjects (50% vs 9.1%).¹⁹ Patients who had NEA reported more prominent cognitive difficulties in memory,⁴² attention/concentration⁴² and language^{42 43} domains than controls with epilepsy. Cognitive symptoms were correlated with the presence of depressed mood^{42 44} and post-traumatic stress disorder (PTSD).⁴⁵

Objective neuropsychological abnormalities

Memory

Fibromyalgia

Selective memory impairment was observed in tests that included stimulus competition,⁴⁶ did not allow information rehearsal,⁴⁷ were vulnerable to distraction⁴⁸ and/or required sustained effort.⁴⁹ Grisart *et al*⁵⁰ reported an impairment of explicit memory contrasting with a preservation of automatic memory processes. A further study reported abnormalities in implicit memory.⁵¹

Abnormal working memory was correlated with pain^{48 52 53} and anxiety⁴⁸ but not with depression.^{48 52 53} Episodic memory impairment was correlated with pain^{51 52} and depression,⁴⁹ but other studies did not find a link with comorbidities.^{50 54}

Chronic fatigue syndrome

Working memory was often reported to be abnormal only in more effortful tasks, and this pattern was attributed to poor attention and slow information processing.^{34 55–58}

Memory registration and consolidation were predominantly impaired relative to memory retrieval,^{31 59–61} which might reflect disruption of information acquisition and encoding by distraction.⁵⁹ However, poor retrieval has also been described.^{62 63}

Table 1 Summary of main findings per condition and domain

Parameter	Functional cognitive disorder***	Fibromyalgia	Chronic fatigue syndrome	Functional neurological disorder†
Cognitive symptoms	Memory Language (word finding)	Attention/concentration Memory Language (word finding)	Attention/concentration Memory Language (word finding) Reasoning	Attention/concentration Memory Language (word finding)
Objective neuropsychological deficits				
Memory	Vulnerability to distraction Explicit worse than implicit memory*		Impaired registration and consolidation* Working memory disruption by abnormal attention/information processing*	Working memory**
Attention/concentration	Selective/divided attention* Bias towards emotionally negative information**		Divided attention Executive function of attention** Bias towards threatening stimuli*	Attention** Attentional bias towards social threatening stimuli**
Executive functions	Cognitive inhibition**			
Information processing			Slow	Slow**
Language	Verbal fluency*		Verbal fluency**	
Social cognition	Alexithymia** Recognising others' emotions*			Alexithymia* Affect expression and recognition**
Consistency on repeat assessment			Increased performance variability	
Factors related with neuropsychological deficits	Pain*		Fatigue**	Psychoopathology**
Discrepancies between symptoms and objective deficits	Symptoms>objective deficits	Symptoms>objective deficits	Symptoms>objective deficits	Symptoms>objective deficits**
Performance validity/effort testing	A minority fail validity testing	Only a minority fail validity testing*	Rare failures on validity testing*	Overall, only a minority fail validity testing
Neurobiology of cognitive impairment	No structural damage	No structural damage* Dysfunction of a fronto-parieto-temporal network involved in attention, memory and executive functions, as well in emotion and pain processing	No structural damage* Dysfunction of the working memory network	No structural damage
Others	Memory perfectionism Overinterpretation of attentional lapses Heightened self-monitoring for cognitive errors Low memory self-efficacy		Heightened perception of effort	

* , some evidence but inconsistent between studies; ** , some evidence but particularly inconsistent and/or scarce; ***Proposed features.

†Non-epileptic attacks and functional movement disorders.

Predominant impairment of verbal relative to visual memory has been described in some studies.^{28 64–66} However, other studies observed that both verbal and visual domains were affected,^{67–69} while others failed to demonstrate significant impairment of any memory domain.^{70–75}

Functional neurological disorder

When compared with healthy subjects, patients with FMD had an isolated deficit on one verbal memory test,¹⁹ and patients who had NEA had a selective impairment of spatial working memory, associated with poor attention.⁷⁶ In comparison with normative data, patients who had NEA demonstrated memory abnormalities,^{77 78} including impairment of working memory associated with poor attention and processing speed.⁷⁹ However, normal memory performance has also been reported.⁸⁰ Patients who had NEA performed similarly,^{80–85} similarly/better^{43 79} or better^{44 86 87} than patients with epileptic seizures (ESs).

Memory impairment was correlated with longer disease duration⁸⁷ and psychopathology,⁸⁷ including PTSD (along with substance abuse and use of psychopharmacological medication),⁴⁵ low mood⁴⁴ and anxiety.⁴³

In summary, the available evidence is heterogeneous but does not support a global impairment of memory in FM, CFS or FND. However, some specific abnormalities have been described, which might reflect the effects of increased vulnerability to distraction on memory performance.

Attention and executive function

Attention, executive function and working memory are closely interlinked. Indeed, storage and processing of items in working memory depends on attentional mechanisms. Moreover, executive function includes the ability to manage sources of distraction.¹⁷

Fibromyalgia

Abnormal attention has been related to an increased vulnerability of memory and other cognitive processes to distraction.^{46 48 52 88–90} However, some studies argue against a generalised impairment of attention.^{91–95}

Miró *et al*⁸⁸ observed a selective impairment of ‘alerting’ and ‘executive control’ but not of ‘orienting’ components of attention (Attentional Network Test-Interactions). Impaired cognitive inhibition has been related to an inability to filter distracting information, underpinning the vulnerability to distraction.^{46 47 96} However, a specific impairment in cognitive inhibition was not confirmed in two studies.^{97 98} Overall, the evidence for abnormal executive function is mixed.^{91 93–95 99 100}

Pain might exhaust attentional resources, thereby increasing susceptibility to distraction. An association between performance on cognitive tests probing attention and pain⁸⁹ and measures of central sensitisation⁹⁰ has been described. However, another study did not confirm this relationship.⁸⁸ Additionally, attentional bias towards emotionally negative information may capture additional processing resources.¹⁰¹

Chronic fatigue syndrome

Patients with CFS do not appear to show generalised abnormalities of attention. However, prominent impairment of divided attention has been reported.¹⁰² Abnormalities on the Paced Auditory Serial Addition Test (PASAT) and Paced Visual Serial Addition Test have been repeatedly observed in CFS.^{31 57 62 63 69 75 103–106} Abnormal performance on these tests may also reflect poor divided attention.¹⁰⁷

Impaired divided attention might account for poor performance on more effortful attentional tasks.^{34 108 109} Patients with CFS might be particularly prone to distraction by irrelevant stimuli, as suggested by impairment on the Stroop interference task.^{27 32 110} However, this finding was not confirmed in all studies.^{25 35 111 112} Attentional bias to threat^{113 114} and towards emotionally negative information¹¹⁵ have been observed. However, others did not confirm a bias towards illness-related information.^{116 117}

A selective impairment of executive function of attention has been described.¹¹⁸ Abnormal executive attention has been related to cause slowing of information processing¹¹⁸ and an attentional bias to threat.^{113 114} Other studies have observed abnormalities in executive functions.^{60 66 71 75 119 120} Finally, impairments of selective and sustained attention and cognitive inhibition have been reported in subjects with comorbid FM.^{121 122}

Functional neurological disorder

In comparison with healthy subjects, patients who had NEA, but not those with FMD, have been reported to have poor attention.^{19 76 81} This was associated with abnormal working memory,⁷⁶ while executive functions were generally normal.^{76 81} In comparison with normative data, patients who had NEA demonstrated abnormalities in attention (associated with poor processing speed and working memory).^{42 77 79} In comparison with patients with ES, patients who had NEA performed similarly on tests of attention^{43 44 80 83 85} and executive function,^{80 82 87} with just one study reporting worse performance on attention.⁷⁹

Overall, evidence across FM, CFS and FND suggests an impairment of selective and/or divided attention.

Information processing

Fibromyalgia

Evidence for impaired information processing as probed by PASAT is mixed, with some studies reporting abnormalities^{46 48} and others normal performance.^{93 94} On 10 measures of information processing speed, patients with FM only performed worse than controls with memory complaints in Stroop colour and word naming; this was proposed to reflect delayed lexical access.⁹³ Some studies did not confirm slower information processing or psychomotor speed.^{52 94 95} Reduced processing speed in mental arithmetic has been correlated with pain and opiate medication but not with depression, anxiety, fatigue and sleep disturbance.¹²³

Chronic fatigue syndrome

Impaired information processing, as measured using PASAT, has been repeatedly reported.^{31 57 62 63 69 103–106} Information processing is related to attention, as it is thought to depend on how fast the attentional system operates. Notably, there was an association between poor divided attention and an impairment of information processing in more challenging settings, where patients need to allocate attention resources to multiple tasks.^{31 55} However, several authors failed to confirm an impairment of information processing^{36 39 41 71 119 124} and/or psychomotor slowing.^{25 39 64} When present, poor information processing was independent of depression,^{31 57 62 63 69 103 106} and the contribution of fatigue was inconsistent.^{31 106 109 125}

Functional neurological disorder

In comparison with healthy subjects, patients who had NEA, but not those with FMD, showed poor information processing.^{19 81} In comparison with normative data, patients who had NEA also

showed slow processing speed, along with poor attention and working memory.⁷⁹ Patients who had NEA performed similarly to patients with ES in tests where information processing plays a role.^{43 79 81 83 85} On tests of psychomotor speed, patients who had NEA performed less well than healthy subjects⁸¹ and similarly to^{81 87} or worse⁸² than patients with ES.

Language and other cognitive domains

Fibromyalgia

Verbal fluency provides a measure of how quickly and efficiently stored knowledge about words can be accessed.¹⁷ Patients with FM showed abnormal verbal fluency (and/or 'verbal knowledge').^{52 91 93 126} Delayed lexical access has been proposed to contribute to difficulties finding words or understanding word meaning.¹²⁶ It has been linked with psychomotor slowing⁹⁹ and with coexisting depression.⁴⁹ However, other studies did not support these associations.^{52 92 94}

Chronic fatigue syndrome

Evidence regarding abnormal verbal fluency is conflicting, with some studies reporting impairment,^{125 127 128} others only selective abnormalities^{34 129} and others essentially normal performance.^{38 39 64 73 102 110 130} CFS twins performed similarly to non-fatigued twins and healthy subjects in tests of language and reasoning.¹¹² Performance on the Boston Naming Test has been consistently normal,^{60 127 128} and several other studies report normal higher cognitive functions, including language⁶³ and abstract reasoning,^{103 130 131} although this last finding is not uniform.¹⁰⁹ Language impairments including poor verbal fluency have been related to abnormal executive functioning but not with coexisting psychiatric disorders or use of psychoactive medication.¹²⁸

Functional neurological disorder

Patients with FMD and healthy subjects performed similarly on verbal fluency tests.¹⁹ In comparison with normative data, patients who had NEA showed either naming and/or fluency abnormalities,^{42 78 79} or normal performances on a composite language score including naming, verbal comprehension and vocabulary.⁸⁰ In comparison with patients with ES, patients who had NEA showed similar^{79 82 83} or better^{43 44 78 81 85 87} performance on language assessment (including confrontational naming, verbal fluency, verbal comprehension and/or writing speed). Longer disease duration was correlated with poorer scores in a language composite including naming, fluency, reading and comprehension.⁸⁷

Social cognition

Social cognition refers to the ability to decipher information about the intentions and affective states of others. This capacity is crucial for the implementation of appropriate behaviour during social interactions. We did not find studies assessing social cognition in CFS.

Fibromyalgia

Patients with FM showed impaired recognition of others' emotions, impaired representation of other people's affective mental states and higher levels of alexithymia but normal empathy quotients.⁹¹ These deficits were independent of executive function and depression/anxiety. Poor emotional face recognition has been associated with pain and, to a lesser degree, with depression, anxiety and alexithymia.¹³² Abnormal social cognition has been related to interoceptive impairment,¹³³ as well as

with an interference of nociception with cognitive and emotional processing, due to the partial overlap of brain networks involved in pain processing and face recognition.

Functional neurological disorder

Alexithymia is characterised by an inability to identify emotions at a cognitive level and has been repeatedly demonstrated in FND,^{76 134 135} although not uniformly.¹³⁶ Patients who had NEA showed worse affect expression and recognition,⁴³ impaired ability to switch between emotion and non-emotion face categorisation,¹³⁷ positive attentional bias towards socially threatening stimuli¹³⁸ and an overall greater intensity of emotional experience and diminished positive emotional behaviour on exposure to pleasant, neutral and unpleasant pictures.¹³⁹ As an exception, normal facial expression recognition has also been described.¹³⁵

Consistency on repeat assessment

Chronic fatigue syndrome

Patients with CFS showed an increased intraindividual variability on performance on repeated neuropsychological testing.¹⁴⁰ Significant variability of self-reported mental fatigue was also observed on repeated assessment.¹⁴¹

These findings can be regarded as evidence for internal inconsistency, which is considered a cardinal feature of functional neurological symptoms in general.⁶

Discrepancies between self-reported cognitive difficulties and objective assessment

Fibromyalgia

Although self-reported difficulties are prominent, evidence for objective cognitive impairment is far more contradictory, suggesting an overestimation of cognitive problems. Unfortunately, only a handful of studies directly compared symptoms with objective performance. In two studies, prominent memory symptoms were disproportionate to objective impairment,^{48 94} particularly after controlling for fatigue, pain and depression.⁹⁴ However, another observed a correlation between memory difficulties and objective performance.¹⁴² Cognitive symptoms have been associated with objective abnormalities in attention, processing speed, memory and executive functions.^{52 98}

Chronic fatigue syndrome

Prominent reported difficulties in memory,^{25 38 124} attention/concentration,^{25 124} information processing¹²⁴ and general cognitive function^{30 38 131 143} contrast with little or no deficit on objective assessment.^{25 30 38 124 131 143} Notably, prominent symptoms have been associated with impairment in only a *minority* of patients.²⁸ Others observed a restricted pattern of neuropsychological impairment, involving attention/concentration, information processing and/or memory domains in the context of multiple, multimodal cognitive symptoms.^{26 63 73} Finally, a correlation between subjective and objective deficits has also been described, involving memory, attention and processing speed.^{27 31–34 55 108 111 144}

'Metamemory' corresponds to self-knowledge about one's own memory capabilities. Impossibly high standards of personal performance might contribute to discrepancies between self-reported and objective abnormalities. This has been related to a tendency to underestimate actual performance relative to expectations of 'normal' performance.³⁵ Contrary to this hypothesis, two studies observed normal estimations of memory abilities and corresponding objective performance.^{30 60}

Functional neurological disorder

A discrepancy between prominent symptoms and relatively normal objective performance has been observed in a few studies.^{19 43 44} However, in one of these, this dissociation was observed in only one among several cognitive domains.⁴³ Moreover, another study found discrepancies in patients who had NEA and in those with ES.⁴²

Performance validity testing (PVT)

PVT was rarely included in test batteries assessing cognition in patients with FM or CFS. This is important in determining the cause of any underperformance on neuropsychological assessment. However, the few studies including PVT did not demonstrate a generalised lack of effort.

Fibromyalgia

Only a minority of patients with FM demonstrated suboptimal effort in the few studies that included PVT. However, the majority of participants performed normally on validity testing.^{94 145-148}

The percentage of patients with FM passing PVT ranged from 63% in Johnson-Greene *et al*¹⁴⁸ and Brooks *et al*,¹⁴⁷ which included subjects receiving or seeking disability benefits or with possible secondary gains, to up to 100% in Iverson *et al*,¹⁴⁵ where patients were not involved in disability benefits assessments. In Gervais *et al*,¹⁴⁶ 65% patients with FM who were on or seeking benefits performed in the normal range on PVT, and this percentage increased to 96% among those not seeking or receiving benefits. Poor performance on PVT was correlated with prominent cognitive symptoms,¹⁴⁶ Millon Clinical Multi-axial Inventory-III scores, including somatoform, depression and anxiety subscales¹⁴⁷ and pain and poor sleep.¹⁴⁸ In one study, patients with FM showed prominent memory complaints but normal objective performance after excluding participants who demonstrated poor effort and controlling for pain, fatigue and depression.⁹⁴

Chronic fatigue syndrome

In Busichio *et al*⁷⁵ the Test of Memory Malingering did not reveal lack of effort in any patient with CFS. In Constant *et al*,⁶⁸ only 2/25 patients with CFS scored under the consistency threshold on PVT. Notably, these studies reported objective impairments of attention, information processing, memory and motor speed.^{68 75} A couple of other studies reported lack of evidence of poor motivation or effort.^{67 71 118 140 149}

A different but related observation is a heightened perception of effort or fatigue after repeated or more challenging cognitive testing, which has been correlated with worse performance on cognitive testing.^{36 131 149 150}

Functional neurological disorder

Performance validity testing in studies of patients with FND has yielded heterogeneous results. Several studies reported worse performance than in controls,^{19 81 151 152} while others reported performance that was normal⁷⁹ or similar to neurological disease control groups.¹⁵¹

The percentage of patients passing on PVT ranged from 48.8%¹⁵³ to 100%,⁷⁹ with most studies suggesting that only a minority proportion of patients with FND fail effort testing.^{151 154 155} Increased failure in comparison with controls was reported by Drane *et al*¹⁵³ but not confirmed by Dodrill⁸² and other authors.^{19 155} Additionally, subjects with FND rarely scored below chance on forced-choice tests,^{81 152 154} which suggests that malingering is uncommon, at least in those patients

with FND who are recruited to observational studies. Finally, the observation of 'isolated' effort deficits, contrasting with normal scores in many other neuropsychological tests also requiring 'effort',¹⁹ raises the hypothesis that some tests of performance validity may be measuring a specific cognitive deficit rather than incomplete effort.

DISCUSSION

FND, FM and CFS appear to be associated with similar, prominent cognitive symptoms, including forgetfulness, distractibility and word-finding difficulties. In contrast, generalised neuropsychological deficits have not been demonstrated.

Nevertheless, our systematic analysis of the literature reveals several specific abnormalities in FM, CFS and, to a lesser extent, FND. Poor selective and divided attention and slow information processing are reported across these disorders (table 1). Memory symptoms might therefore reflect increased vulnerability to distraction, which interferes with memory registration.

Coexisting psychopathology is not robustly associated with cognitive difficulties.^{26 31 48 52 53 57 62 63 69 103 106 123 128} However, both subjective cognitive symptoms and objective cognitive deficits correlate with pain in FM and with mental or physical exertion and fatigue in CFS.

Overall, these conditions show a mismatch between prominent cognitive symptoms and, at best, only selective objective abnormalities. Importantly, the few studies including PVT reported poor effort in only a small proportion of patients, with the exception of patients with FM receiving or seeking disability benefits, where one-third failed PVT.

The cognitive profiles of FM, CFS and non-cognitive FND share several features with the recently proposed FCD (table 1), including a disproportion between self-reported memory and word-finding difficulties and objective neuropsychological deficits and a strong bias towards symptoms with a likely attentional basis.

Some key features proposed to underpin FCD, including memory perfectionism, overinterpretation of attentional lapses and heightened self-monitoring for cognitive errors, have not been widely investigated in these disorders. However, in the CFS literature, there is some evidence for unrealistic expectations of cognitive performance,³⁴ while in FM, there are well-described alterations in interoceptive awareness.¹³² These findings suggest the possibility of common risk factors, but empirical evidence confirming the relevance of these findings remains sparse.

Interestingly, neuropsychological assessment in FCD has not shown deficits in straightforward tests of attention,^{8 10} suggesting that it is a lack of attentional reserve rather than inattention per se that underpins the subjective experience of cognitive impairment. To date, however, the more complex tests of attention used in the literature on FND, CFS and FM have not been reported in an FCD cohort.

Reduced attentional reserve could increase susceptibility to distraction and slow information processing. Heightened distractibility and attentional lapses could then contribute to the subjective experience of memory impairment (figure 1). Abnormal attention might have a disproportionate impact on real-life multitasking, but be more difficult to measure using standard neuropsychological evaluation.¹⁰²

Excessive attention towards the body, severe pain and fatigue might underlie the decrease in externally directed attention. In the context of alexithymia, difficulties in recognising somatic markers of emotions might promote body monitoring and further exhaust attentional resources (figure 1).




Figure 1 Proposed mechanisms underpinning cognitive difficulties across the spectrum of functional cognitive disorders spectrum.

Memory perfectionism and heightened self-monitoring of cognitive processes and errors might then lead to overinterpretation of such attentional lapses in daily life which, in turn, could drive advice-seeking from healthcare professionals (figure 1). Interestingly, the ability to ‘normalise’ somatic experiences inversely correlates with health seeking behaviour in FM.¹⁵⁶ Perfectionism may be relevant in FM and CFS. Women with FM do not have greater perfectionist traits overall, but perfectionism correlates with poor health status.¹⁵⁷ In CFS, self-critical perfectionism has been shown to predict fatigue and pain levels.¹⁵⁸ However, the concept of memory perfectionism as a core feature of, or risk factor for, FCD needs further elucidation.

Another interesting observation is the heightened perception of effort after repeated or more challenging cognitive testing in patients with CFS.^{36 131 149 150} Heightened perception of effort is proposed to reflect an interpretive bias for somatic information,¹¹⁶ a greater operational demand to control and monitor cognitive processes¹⁵⁹ or a sense of insufficiency resulting from unrealistically high expectations about one’s performance, leading to an overestimation of exertion and fatigue.³⁵

If present across this group of disorders, a heightened perception of cognitive effort may also contribute to the experience of illness, due to the extreme effort invested in otherwise trivial daily tasks. The experience of illness would then be decoupled, to a degree, from objective cognitive performance. Therefore, abnormal effort may well constitute a piece in the jigsaw of FCD.

Cognitive effort can be regarded as a cost of *controlled* cognitive processes.¹⁶⁰ A switch from an *automatic* to a *controlled* mode of cognitive processing may therefore underlie the increase in cognitive effort. Notably, this would mirror what is observed in FMD, where a switch from an automatic to an explicit (‘conscious’) mode of motor control impairs motor performance.¹⁶¹

In addition, the hypothetical switch from an automatic to a highly controlled (and effortful) cognitive operational mode

could be a maladaptive attempt to compensate for ‘memory failures’ that were invested with pathological significance. Paradoxically, an exaggerated investment of effort could also contribute to the failure on performance validity testing by a minority of patients. Indeed, motivation depends on a cost–benefit analysis,¹⁶² and a ‘high effort’ operational mode could lead to earlier disengagement due to unacceptably high performance costs.

Although our systematic review focused on FND, FM and CFS, there are other conditions where prominent cognitive difficulties contrast with inconsistent neuropsychological deficits and an absence of structural lesions on conventional neuroimaging.

Postconcussion syndrome (PCS) is characterised by a constellation of ‘non-specific’ neuropsychiatric symptoms following a concussion or mild traumatic brain injury. Similarly to FND, FM and CFS, patients report poor concentration, mental slowness, forgetfulness and word-finding difficulties.¹⁶³

Neuropsychological assessment in PCS does not confirm widespread cognitive deficits.¹⁶⁴ However, specific abnormalities in selective and divided attention, information processing and working memory have been described, again mirroring observations in FND, FM and CFS.^{165–168} This similarity was corroborated by direct comparisons between PCS and CFS.⁶¹

Whiplash syndrome follows a hyperflexion, hyperextension or hyperlateroversion of the neck without external head injury or loss of consciousness. Patients typically describe chronic pain involving neck/shoulders. Cognitive difficulties and other associated symptoms are similar to PCS.^{169–171} Notwithstanding the controversy regarding the inconsistency of objective neuropsychological deficits, a meta-analysis reported objective abnormalities in attention, working memory, immediate and delayed recall, visuomotor tracking and cognitive flexibility.¹⁷²

Functional imaging studies in the disorders investigated here, as well as in PCS, have described dysfunction of a fronto-parieto-temporal network postulated to be involved in attention,

memory and executive functions as well as in emotion and pain processing.^{36 173–179} This would provide a neurobiological basis for the proposed interaction between attention-related cognitive functions and abnormalities with pain and emotional processing as well as movement control.

Interestingly, negative expectations regarding the consequences of a head injury and recall bias have both been implicated in self-reported symptoms after concussion.^{180 181} This gains particular relevance given the proposed role of abnormal predictions or expectations in the pathogenesis of functional neurological symptoms.¹⁴

Limitations

Evidence for the presence and characteristics of cognitive abnormalities in FNDs, FM and CFS is remarkably heterogeneous.

Sample sizes were usually small, and a control group of healthy subjects was often lacking. In Vercoulen *et al*,²⁸ objective cognitive impairment in a minority of patients with CFS was sufficient to drive between-group differences towards statistical significance. In these disorders, population heterogeneity means that small sample sizes and the use of average scores (rather than, eg, the proportion of individuals performing below threshold) might conceivably lead to false-positive results.

Notably, cognition in FND has been far less studied than in FM and CFS. Moreover, among 39 studies on FND, 36 focused on NEA, 1 in neurological ‘medically unexplained symptoms’¹⁵⁴ and only 2 on FMD.^{19 134} Although we decided to use the more encompassing term ‘FND’, available evidence does not allow us to conclude much about the cognitive profile of FMD. However, it would be reasonable to expect a degree of overlap among FND subtypes.

Patients with FM, CFS and FND are often treated with medications such as benzodiazepines or opioids, which can have cognitive side effects. Unfortunately, the possible contribution of medication to cognitive performance was infrequently assessed. Although blinding in neuropsychological evaluations was rare, the increased availability of computerised evaluations might have helped reduce observer bias.

Despite the ongoing discussion about the discordance between subjective and objective abnormalities, relatively few studies investigated discrepancies within the same sample. In addition, PVT was rarely included in neuropsychological assessment, despite the pervasive belief that patients ‘exaggerate’ their symptoms, do not apply sufficient effort in cognitive testing or may be seeking secondary gains.

The variety of methods used for PVT was striking. Furthermore, the definition of ‘poor effort’ was heterogeneous, including: scoring below chance in forced-choice tests, scoring below/above a prespecified cut-off proposed to distinguish ‘poor effort’ and scoring lower than controls in PVT. It is important to highlight that poor effort does not necessarily equate to malingering or a factitious disorder. For example, patients with FM are frequently affected by performance anxiety, and one can imagine that it may interfere with falsely ‘challenging’ effort tasks. Another contributor could be ‘cogniphobia’, a concept akin to ‘kinesiophobia’ that reflects a reluctance to engage in any cognitive task, due to a fear that doing so may aggravate other symptoms.¹⁸²

Finally, we acknowledge that there is ongoing debate about the aetiology and diagnosis of CFS and FM, in particular the relevance or not of ‘organic’ and ‘non-organic’ factors.^{183 184}

There is considerable debate, particularly with regard to CFS, about how to define and diagnose the condition, and indeed whether there may be a number of different conditions present within groups of patients diagnosed with CFS. We accept that these debates are important and as yet unresolved and that heterogeneity in the CFS and FM populations reviewed here might have reduced the power of studies to identify consistent deficits.

CONCLUSION

Given the considerable heterogeneity in the literature and its methodological shortcomings, further research is warranted before reaching firm conclusions about the nature of cognitive difficulties in FND, CFS and FMD or the existence of an overarching ‘functional cognitive disorder spectrum’, also encompassing conditions such as PCS.

However, the available evidence does not support the existence of separate cognitive disorders in CFS, FM and FND. Rather, these conditions share common cognitive symptoms with an emphasis on attentional dysfunction. The newly emerging concept of FCD shares similar symptoms and hypothesised mechanistic underpinnings. Our understanding of these cognitive disorders and the development of novel interventions might be facilitated by regarding them as related conditions whose cognitive symptoms fall within the FCD spectrum, rather than as discrete entities. While evidence-based treatment for FCD remains lacking, interventions for fatigue, pain and functional non-cognitive neurological symptoms are available.^{185–187} Therefore, practitioners assessing patients with FCD should look for these features, which may be missed if the consultation is exclusively focused on cognitive complaints. Clinical trials in FCD might reasonably also include subjects with CFS, FM or FND-related cognitive difficulties, which will generate larger sample sizes and lead to faster development of evidence-based treatment for this increasingly prevalent and distressing disorder.

Correction notice Since this paper was first published online the third affiliation has been updated.

Contributors The study was conceived by JDI and MJE; TT performed data acquisition and wrote the first draft, which was revised by JDI and MJE; all authors contributed to study design, data interpretation and approving the final version of the manuscript.

Funding This work was supported by the Wellcome Trust (204809/Z/16/Z), Medical Research Council and Fundação para a Ciência e Tecnologia (SFRH/SINTD/95267/2013).

Competing interests None declared.

Patient consent Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement There is no additional unpublished data from this study.

Open access This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Menon R, Lerner AJ. Use of cognitive screening instruments in primary care: the impact of national dementia directives (NICE/SCIE, National Dementia Strategy). *Fam Pract* 2011;28:272–6.
- Lerner A. Impact of the National Dementia Strategy in a neurology-led memory clinic: 5-year data. *Clin Med* 2014;14:216.

- 3 Bell S, Harkness K, Dickson JM, *et al.* A diagnosis for £55: what is the cost of government initiatives in dementia case finding. *Age Ageing* 2015;44:344–5.
- 4 McCaffrey RJ, Bauer L, Palav AA, O'Bryant S, *et al.* eds. *Practitioner's Guide to Symptom Base Rates in the General Population* [Internet. New York: Springer-Verlag, 2006. <http://www.springer.com/us/book/9780387267579>.
- 5 Pennington C, Hayre A, Newson M, *et al.* Functional cognitive disorder: a common cause of subjective cognitive symptoms. *J Alzheimers Dis* 2015;48(s1):S19–S24.
- 6 Stone J, Pal S, Blackburn D, *et al.* Functional (psychogenic) cognitive disorders: a perspective from the neurology clinic. *J Alzheimers Dis* 2015;48(s1):S5–S17.
- 7 Delis DC, Wetter SR. Cogniform disorder and cogniform condition: proposed diagnoses for excessive cognitive symptoms. *Arch Clin Neuropsychol* 2007;22:589–604.
- 8 Schmidtko K, Pohlmann S, Metternich B. The syndrome of functional memory disorder: definition, etiology, and natural course. *Am J Geriatr Psychiatry* 2008;16:981–8.
- 9 Pennington C, Newson M, Hayre A, *et al.* Functional cognitive disorder: what is it and what to do about it? *Pract Neurol* 2015;15:436–44.
- 10 Wakefield SJ, Blackburn DJ, Harkness K, *et al.* Distinctive neuropsychological profiles differentiate patients with functional memory disorder from patients with amnesic-mild cognitive impairment. *Acta Neuropsychiatr* 2018;30:1–7.
- 11 Mol ME, van Boxtel MP, Willems D, *et al.* Do subjective memory complaints predict cognitive dysfunction over time? A six-year follow-up of the Maastricht Aging Study. *Int J Geriatr Psychiatry* 2006;21:432–41.
- 12 Carson A, Hallett M, Stone J. Assessment of patients with functional neurologic disorders. *Handb Clin Neurol* 2016;139:169–88.
- 13 Reuber M, Blackburn DJ, Elyse C, *et al.* An interactional profile to assist the differential diagnosis of neurodegenerative and functional memory disorders. *Alzheimer Dis Assoc Disord* 2018:1.
- 14 Edwards MJ, Adams RA, Brown H, *et al.* A Bayesian account of 'hysteria'. *Brain* 2012;135(11):3495–512.
- 15 Wessely S, Nimnuan C, Sharpe M. Functional somatic syndromes: one or many? *The Lancet* 1999;354:936–9.
- 16 Griem J, Stone J, Carson A, *et al.* Psychologic/functional forms of memory disorder. *Handb Clin Neurol* 2016;139:407–17.
- 17 Glass JM. Review of cognitive dysfunction in fibromyalgia: a convergence on working memory and attentional control impairments. *Rheum Dis Clin North Am* 2009;35:299–311.
- 18 Cvejic E, Birch RC, Vollmer-Conna U. Cognitive dysfunction in chronic fatigue syndrome: a review of recent evidence. *Curr Rheumatol Rep* 2016;18:24.
- 19 Heintz CE, van Tricht MJ, van der Salm SM, *et al.* Neuropsychological profile of psychogenic jerky movement disorders: importance of evaluating non-credible cognitive performance and psychopathology. *J Neurol Neurosurg Psychiatry* 2013;84:862–7.
- 20 Mease PJ, Arnold LM, Crofford LJ, *et al.* Identifying the clinical domains of fibromyalgia: contributions from clinician and patient Delphi exercises. *Arthritis Rheum* 2008;59:952–60.
- 21 Bennett RM, Jones J, Turk DC, *et al.* An internet survey of 2,596 people with fibromyalgia. *BMC Musculoskelet Disord* 2007;8:27.
- 22 Williams DA, Clauw DJ, Glass JM. Perceived cognitive dysfunction in fibromyalgia syndrome. *J Musculoskelet Pain* 2011;19:66–75.
- 23 Leavitt F, Katz RS, Mills M, *et al.* Cognitive and dissociative manifestations in fibromyalgia. *J Clin Rheumatol* 2002;8:77–84.
- 24 Katz RS, Heard AR, Mills M, *et al.* The prevalence and clinical impact of reported cognitive difficulties (fibrofog) in patients with rheumatic disease with and without fibromyalgia. *J Clin Rheumatol* 2004;10:53–8.
- 25 Fiedler N, Kipen HM, DeLuca J, *et al.* A controlled comparison of multiple chemical sensitivities and chronic fatigue syndrome. *Psychosom Med* 1996;58:38–49.
- 26 McDonald E, Cope H, David A. Cognitive impairment in patients with chronic fatigue: a preliminary study. *J Neurol Neurosurg Psychiatry* 1993;56:812–5.
- 27 Smith AP, Behan PO, Bell W, *et al.* Behavioural problems associated with the chronic fatigue syndrome. *Br J Psychol* 1993;84:411–23.
- 28 Vercoulen JH, Bazelmans E, Swanink CM, *et al.* Evaluating neuropsychological impairment in chronic fatigue syndrome. *J Clin Exp Neuropsychol* 1998;20:144–56.
- 29 Schmalzing KB, Betterton KL. Neurocognitive complaints and functional status among patients with chronic fatigue syndrome and fibromyalgia. *Quality of Life Research* 2016;25:1257–63.
- 30 Lakein DA, Fantie BD, Grafman J, *et al.* Patients with chronic fatigue syndrome and accurate feeling-of-knowing judgments. *J Clin Psychol* 1997;53:635–45.
- 31 Michiels V, de Gucht V, Cluydts R, *et al.* Attention and information processing efficiency in patients with Chronic Fatigue Syndrome. *J Clin Exp Neuropsychol* 1999;21:709–29.
- 32 Thomas M, Smith A. An investigation into the cognitive deficits associated with chronic fatigue syndrome. *Open Neurol J* 2009;3:13–23.
- 33 Ickmans K, Meeus M, Kos D, *et al.* Cognitive performance is of clinical importance, but is unrelated to pain severity in women with chronic fatigue syndrome. *Clin Rheumatol* 2013;32:1475–85.
- 34 Joyce E, Blumenthal S, Wessely S. Memory, attention, and executive function in chronic fatigue syndrome. *J Neurol Neurosurg Psychiatry* 1996;60:495–503.
- 35 Metzger FA, Denney DR. Perception of cognitive performance in patients with chronic fatigue syndrome. *Ann Behav Med* 2002;24:106–12.
- 36 Schmalzing KB, Lewis DH, Fiedelak JL, *et al.* Single-photon emission computerized tomography and neurocognitive function in patients with chronic fatigue syndrome. *Psychosom Med* 2003;65:129–36.
- 37 Wessely S, Powell R. Fatigue syndromes: a comparison of chronic "postviral" fatigue with neuromuscular and affective disorders. *J Neurol Neurosurg Psychiatry* 1989;52:940–8.
- 38 Cope H, Pernet A, Kendall B, *et al.* Cognitive functioning and magnetic resonance imaging in chronic fatigue. *Br J Psychiatry* 1995;167:86–94.
- 39 Blackwood SK, MacHale SM, Power MJ, *et al.* Effects of exercise on cognitive and motor function in chronic fatigue syndrome and depression. *Journal of Neurology, Neurosurgery & Psychiatry* 1998;65:541–6.
- 40 Attree EA, Arroll MA, Dancey CP, *et al.* Psychosocial factors involved in memory and cognitive failures in people with myalgic encephalomyelitis/chronic fatigue syndrome. *Psychol Res Behav Manag* 2014;7:67–76.
- 41 Gotts ZM, Ellis JG, Deary V, *et al.* The association between daytime napping and cognitive functioning in chronic fatigue syndrome. *PLoS One* 2015;10:e0117136.
- 42 Fargo JD, Scheffert BK, Szaflarski JP, *et al.* Accuracy of self-reported neuropsychological functioning in individuals with epileptic or psychogenic nonepileptic seizures. *Epilepsy & Behavior* 2004;5:143–50.
- 43 Prigatano GP, Kirlin KA. Self-appraisal and objective assessment of cognitive and affective functioning in persons with epileptic and nonepileptic seizures. *Epilepsy & Behavior* 2009;14:387–92.
- 44 Breier JL, Fuchs KL, Brookshire BL, *et al.* Quality of life perception in patients with intractable epilepsy or pseudoseizures. *Arch Neurol* 1998;55:660–5.
- 45 Myers L, Zeng R, Perrine K, *et al.* Cognitive differences between patients who have psychogenic nonepileptic seizures (PNESs) and posttraumatic stress disorder (PTSD) and patients who have PNESs without PTSD. *Epilepsy Behav EB* 2014;37:82–6.
- 46 Leavitt F, Katz RS. Distraction as a key determinant of impaired memory in patients with fibromyalgia. *J Rheumatol* 2006;33:127–32.
- 47 Leavitt F, Katz RS. Normalizing memory recall in fibromyalgia with rehearsal: a distraction-counteracting effect. *Arthritis & Rheumatism* 2009;61:740–4.
- 48 Grace GM, Nielson WR, Hopkins M, *et al.* Concentration and memory deficits in patients with fibromyalgia syndrome. *J Clin Exp Neuropsychol* 1999;21:477–87.
- 49 Landrø NI, Stiles TC, Sletvold H. Memory functioning in patients with primary fibromyalgia and major depression and healthy controls. *J Psychosom Res* 1997;42:297–306.
- 50 Grisart J, Van der Linden M, Masquelier E. Controlled processes and automaticity in memory functioning in fibromyalgia patients: relation with emotional distress and hypervigilance. *J Clin Exp Neuropsychol* 2002;24:994–1009.
- 51 Duschek S, Werner NS, Winkelmann A, *et al.* Implicit memory function in fibromyalgia syndrome. *Behav Med* 2013;39:11–16.
- 52 Park DC, Glass JM, Minear M, *et al.* Cognitive function in fibromyalgia patients. *Arthritis & Rheumatism* 2001;44:2125–33.
- 53 Dick BD, Verrier MJ, Harker TK, *et al.* Disruption of cognitive function in Fibromyalgia Syndrome. *Pain* 2008;139:610–6.
- 54 Canovas R, Leon I, Roldan MD, *et al.* Virtual reality tasks disclose spatial memory alterations in fibromyalgia. *Rheumatology* 2009;48:1273–8.
- 55 Michiels V, Cluydts R, Fischler B. Attention and verbal learning in patients with chronic fatigue syndrome. *J Int Neuropsychol Soc* 1998;4:456–66.
- 56 Dobbs BM, Dobbs AR, Kiss I. Working memory deficits associated with chronic fatigue syndrome. *J Int Neuropsychol Soc* 2001;7:285–93.
- 57 DeLuca J, Christodoulou C, Diamond BJ, *et al.* Working memory deficits in chronic fatigue syndrome: differentiating between speed and accuracy of information processing. *J Int Neuropsychol Soc* 2004;10:101–9.
- 58 Marshall PS, Forstot M, Callies A, *et al.* Cognitive slowing and working memory difficulties in chronic fatigue syndrome. *Psychosom Med* 1997;59:58–66.
- 59 Johnson SK, DeLuca J, Diamond BJ, *et al.* Memory dysfunction in fatiguing illness: examining interference and distraction in short-term memory. *Cogn Neuropsychiatry* 1998;3:269–85.
- 60 Sandman CA, Barron JL, Nackoul K, *et al.* Memory deficits associated with chronic fatigue immune dysfunction syndrome. *Biol Psychiatry* 1993;33:618–23.
- 61 Tiersky LA, Cicerone KD, Natelson BH, *et al.* Neuropsychological functioning in chronic fatigue syndrome and mild traumatic brain injury: a comparison. *The Clinical Neuropsychologist* 1998;12:503–12.
- 62 DeLuca J, Christodoulou C, Diamond BJ, *et al.* The nature of memory impairment in chronic fatigue syndrome. *Rehabil Psychol* 2004;49:62–70.
- 63 DeLuca J, Johnson SK, Beldowicz D, *et al.* Neuropsychological impairments in chronic fatigue syndrome, multiple sclerosis, and depression. *J Neurol Neurosurg Psychiatry* 1995;58:38–43.
- 64 Riccio M, Thompson C, Wilson B, *et al.* Neuropsychological and psychiatric abnormalities in myalgic encephalomyelitis: A preliminary report. *Br J Clin Psychol* 1992;31(Pt 1):111–20.
- 65 Neu D, Kajosch H, Peigneux P, *et al.* Cognitive impairment in fatigue and sleepiness associated conditions. *Psychiatry Res* 2011;189:128–34.

- 66 Claypoole KH, Noonan C, Mahurin RK, et al. A twin study of cognitive function in chronic fatigue syndrome: the effects of sudden illness onset. *Neuropsychology* 2007;21:507–13.
- 67 Michiels V, Cluydts R, Fischler B, et al. Cognitive functioning in patients with chronic fatigue syndrome. *J Clin Exp Neuropsychol* 1996;18:666–77.
- 68 Constant EL, Adam S, Gillain B, et al. Cognitive deficits in patients with chronic fatigue syndrome compared to those with major depressive disorder and healthy controls. *Clin Neurol Neurosurg* 2011;113:295–302.
- 69 DeLuca J, Johnson SK, Ellis SP, et al. Cognitive functioning is impaired in patients with chronic fatigue syndrome devoid of psychiatric disease. *J Neurol Neurosurg Psychiatry* 1997;62:151–5.
- 70 Wearden A, Appleby L. Cognitive performance and complaints of cognitive impairment in chronic fatigue syndrome (CFS). *Psychol Med* 1997;27:81–90.
- 71 Grafman J, Schwartz V, Dale JK, et al. Analysis of neuropsychological functioning in patients with chronic fatigue syndrome. *J Neurol Neurosurg Psychiatry* 1993;56:684–9.
- 72 Johnson SK, DeLuca J, Fiedler N, et al. Cognitive functioning of patients with chronic fatigue syndrome. *Clin Infect Dis* 1994;18 Suppl 1(Supplement_1):S84–S85.
- 73 Gaudino EA, Coyle PK, Krupp LB. Post-Lyme syndrome and chronic fatigue syndrome. Neuropsychiatric similarities and differences. *Arch Neurol* 1997;54:1372–6.
- 74 Tiersky LA, Matheis RJ, Deluca J, et al. neuropsychological functioning, and mood in chronic fatigue syndrome (CFS): relationship to psychiatric disorder. *J Nerv Ment Dis* 2003;191:324–31.
- 75 Busichio K, Tiersky LA, Deluca J, et al. Neuropsychological deficits in patients with chronic fatigue syndrome. *J Int Neuropsychol Soc* 2004;10:278–85.
- 76 O'Brien FM, Fortune GM, Dicker P, et al. Psychiatric and neuropsychological profiles of people with psychogenic nonepileptic seizures. *Epilepsy & Behavior* 2015;43:39–45.
- 77 Kalogjera-Sackellares D, Sackellares JC. Intellectual and neuropsychological features of patients with psychogenic pseudoseizures. *Psychiatry Res* 1999;86:73–84.
- 78 Reuber M, Fernández G, Helmstaedter C, et al. Evidence of brain abnormality in patients with psychogenic nonepileptic seizures. *Epilepsy & Behavior* 2002;3:249–54.
- 79 Strutt AM, Hill SW, Scott BM, et al. A comprehensive neuropsychological profile of women with psychogenic nonepileptic seizures. *Epilepsy & Behavior* 2011;20:24–8.
- 80 Walterfang M, Choi Y, O'Brien TJ, et al. Utility and validity of a brief cognitive assessment tool in patients with epileptic and nonepileptic seizures. *Epilepsy & Behavior* 2011;21:177–83.
- 81 Binder LM, Kindermann SS, Heaton RK, et al. Neuropsychologic impairment in patients with nonepileptic seizures. *Arch Clin Neuropsychol* 1998;13:513–22.
- 82 Dodrill CB. Do patients with psychogenic nonepileptic seizures produce trustworthy findings on neuropsychological tests? *Epilepsia* 2008;49:691–5.
- 83 Turner K, Piazzini A, Chiesa V, et al. Patients with epilepsy and patients with psychogenic non-epileptic seizures: video-EEG, clinical and neuropsychological evaluation. *Seizure* 2011;20:706–10.
- 84 Lelliott PT, Fenwick P. Cerebral pathology in pseudoseizures. *Acta Neurol Scand* 1991;83:129–32.
- 85 Hill SW, Gale SD. Neuropsychological characteristics of nonepileptic seizure semiological subgroups. *Epilepsy & Behavior* 2011;22:255–60.
- 86 Sackellares JC, Giordani B, Berent S, et al. Patients with pseudoseizures: intellectual and cognitive performance. *Neurology* 1985;35:116–9.
- 87 Locke DEC, Berry DTR, Fakhoury TA, et al. Relationship of indicators of neuropathology, psychopathology, and effort to neuropsychological results in patients with epilepsy or psychogenic non-epileptic seizures. *J Clin Exp Neuropsychol* 2006;28:325–40.
- 88 Miró E, Martínez MP, Sánchez AI, et al. Men and women with fibromyalgia: relation between attentional function and clinical symptoms. *Br J Health Psychol* 2015;20:632–47.
- 89 Dick B, Eccleston C, Crombez G. Attentional functioning in fibromyalgia, rheumatoid arthritis, and musculoskeletal pain patients. *Arthritis Rheum* 2002;47:639–44.
- 90 Coppieters I, Ickmans K, Cagnie B, et al. Cognitive Performance Is Related to Central Sensitization and Health-related Quality of Life in Patients with Chronic Whiplash-Associated Disorders and Fibromyalgia. *Pain Physician* 2015;18:E389–401.
- 91 Di Tella M, Castelli L, Colonna F, et al. Theory of mind and emotional functioning in fibromyalgia syndrome: an investigation of the relationship between social cognition and executive function. *PLoS One* 2015;10:e0116542.
- 92 Cherry BJ, Zettel-Watson L, Shimizu R, et al. Cognitive performance in women aged 50 years and older with and without fibromyalgia. *J Gerontol B Psychol Sci Soc Sci* 2014;69:199–208.
- 93 Leavitt F, Katz RS. Speed of mental operations in fibromyalgia: a selective naming speed deficit. *J Clin Rheumatol Pract Rep Rheum Musculoskelet Dis* 2008;14:214–8.
- 94 Suhr JA. Neuropsychological impairment in fibromyalgia: relation to depression, fatigue, and pain. *J Psychosom Res* 2003;55:321–9.
- 95 Walitt B, Roebuck-Spencer T, Bleiberg J, et al. Automated neuropsychiatric measurements of information processing in fibromyalgia. *Rheumatol Int* 2008;28:561–6.
- 96 Verdejo-García A, Lopez-Torrecillas F, Calandre EP, et al. Executive function and decision-making in women with fibromyalgia. *Archives of Clinical Neuropsychology* 2009;24:113–22.
- 97 Veldhuijzen DS, Sondaal SFV, Oosterman JM. Intact cognitive inhibition in patients with fibromyalgia but evidence of declined processing speed. *The Journal of Pain* 2012;13:507–15.
- 98 Tesio V, Torta DME, Colonna F, et al. Are fibromyalgia patients cognitively impaired? objective and subjective neuropsychological evidence. *Arthritis Care Res* 2015;67:143–50.
- 99 de Melo LF, Da-Silva SL. Neuropsychological assessment of cognitive disorders in patients with fibromyalgia, rheumatoid arthritis, and systemic lupus erythematosus. *Rev Bras Reumatol* 2012;52:181–8.
- 100 Cuevas-Toro Ana M^a, López-Torrecillas F, Díaz-Batanero M^a Carmen, et al. Neuropsychological function, anxiety, depression and pain impact in fibromyalgia patients. *Span J Psychol* 2014;17:E78.
- 101 Duschek S, Werner NS, Limbert N, et al. Attentional bias toward negative information in patients with fibromyalgia syndrome. *Pain Medicine* 2014;15:603–12.
- 102 Ross S, Fantie B, Straus SF, et al. Divided attention deficits in patients with chronic fatigue syndrome. *Appl Neuropsychol* 2001;8:4–11.
- 103 DeLuca J, Johnson SK, Natelson BH. Information processing efficiency in chronic fatigue syndrome and multiple sclerosis. *Arch Neurol* 1993;50:301–4.
- 104 Johnson SK, DeLuca J, Diamond BJ, et al. Selective impairment of auditory processing in chronic fatigue syndrome: a comparison with multiple sclerosis and healthy controls. *Percept Mot Skills* 1996;83:51–62.
- 105 DeLuca J, Johnson SK, Ellis SP, et al. Sudden vs gradual onset of chronic fatigue syndrome differentiates individuals on cognitive and psychiatric measures. *J Psychiatry Res* 1997;31:83–90.
- 106 Johnson SK, Lange G, DeLuca J, et al. The effects of fatigue on neuropsychological performance in patients with chronic fatigue syndrome, multiple sclerosis, and depression. *Appl Neuropsychol* 1997;4:145–53.
- 107 Tombaugh TN. A comprehensive review of the Paced Auditory Serial Addition Test (PASAT). *Arch Clin Neuropsychol* 2006;21:53–76.
- 108 Capuron L, Welberg L, Heim C, et al. Cognitive dysfunction relates to subjective report of mental fatigue in patients with chronic fatigue syndrome. *Neuropsychopharmacology* 2006;31:1777–84.
- 109 Smith AP, Borysiewicz L, Pollock J, et al. Acute fatigue in chronic fatigue syndrome patients. *Psychol Med* 1999;29:283–90.
- 110 Santamarina-Perez P, Eiroa-Orosa FJ, Freniche V, et al. Length of illness does not predict cognitive dysfunction in chronic fatigue syndrome. *Appl Neuropsychol* 2011;18:216–22.
- 111 Ray C, Phillips L, Weir WR. Quality of attention in chronic fatigue syndrome: subjective reports of everyday attention and cognitive difficulty, and performance on tasks of focused attention. *Br J Clin Psychol* 1993;32(Pt 3):357–64.
- 112 Mahurin RK, Claypoole KH, Goldberg JH, et al. Cognitive processing in monozygotic twins discordant for chronic fatigue syndrome. *Neuropsychology* 2004;18:232–9.
- 113 Hou R, Moss-Morris R, Risdale A, et al. Attention processes in chronic fatigue syndrome: attentional bias for health-related threat and the role of attentional control. *Behav Res Ther* 2014;52:9–16.
- 114 Hou R, Moss-Morris R, Bradley BP, et al. Attentional bias towards health-threat information in chronic fatigue syndrome. *J Psychosom Res* 2008;65:47–50.
- 115 Creswell C, Chalder T. Underlying self-esteem in chronic fatigue syndrome. *J Psychosom Res* 2002;53:755–61.
- 116 Moss-Morris R, Petrie KJ. Experimental evidence for interpretive but not attention biases towards somatic information in patients with chronic fatigue syndrome. *Br J Health Psychol* 2003;8(Pt 2):195–208.
- 117 Martin M, Alexeeva I. Mood volatility with rumination but neither attentional nor interpretation biases in chronic fatigue syndrome. *Br J Health Psychol* 2010;15(Pt 4):779–96.
- 118 Togo F, Lange G, Natelson BH, et al. Attention network test: assessment of cognitive function in chronic fatigue syndrome. *J Neuropsychol* 2015;9:1–9.
- 119 Cook DB, Nagelkirk PR, Peckerman A, et al. Exercise and cognitive performance in chronic fatigue syndrome. *Med Sci Sports Exerc* 2005;37:1460–7.
- 120 Neu D, Mairesse O, Montana X, et al. Dimensions of pure chronic fatigue: psychophysical, cognitive and biological correlates in the chronic fatigue syndrome. *Eur J Appl Physiol* 2014;114:1841–51.
- 121 Ickmans K, Meeus M, De Koning M, et al. Can recovery of peripheral muscle function predict cognitive task performance in chronic fatigue syndrome with and without fibromyalgia? *Phys Ther* 2014;94:511–22.
- 122 Ickmans K, Meeus M, De Koning M, et al. Associations Between Cognitive Performance and Pain in Chronic Fatigue Syndrome: Comorbidity with Fibromyalgia Does Matter. *Pain Physician* 2015;18:e635–e636.
- 123 Reyes Del Paso GA, Pulgar A, Duschek S, et al. Cognitive impairment in fibromyalgia syndrome: the impact of cardiovascular regulation, pain, emotional disorders and medication. *Eur J Pain* 2012;16:421–9.
- 124 Short K, McCabe M, Tooley G. Cognitive functioning in chronic fatigue syndrome and the role of depression, anxiety, and fatigue. *J Psychosom Res* 2002;52:475–83.

- 125 Claypoole K, Mahurin R, Fischer ME, *et al.* Cognitive compromise following exercise in monozygotic twins discordant for chronic fatigue syndrome: fact or artifact? *Appl Neuropsychol* 2001;8:31–40.
- 126 Leavitt F, Katz RS. Cognitive dysfunction in fibromyalgia: slow access to the mental lexicon. *Psychol Rep* 2014;115:828–39.
- 127 Lawrie SM, MacHale SM, Cavanagh JT, *et al.* The difference in patterns of motor and cognitive function in chronic fatigue syndrome and severe depressive illness. *Psychol Med* 2000;30:433–42.
- 128 Marcel B, Komaroff AL, Fagioli LR, *et al.* Cognitive deficits in patients with chronic fatigue syndrome. *Biol Psychiatry* 1996;40:535–41.
- 129 Daly E, Komaroff AL, Bloomingdale K, *et al.* Neuropsychological function in patients with chronic fatigue syndrome, multiple sclerosis, and depression. *Appl Neuropsychol* 2001;8:12–22.
- 130 Krupp LB, Sliwinski M, Masur DM, *et al.* Cognitive functioning and depression in patients with chronic fatigue syndrome and multiple sclerosis. *Arch Neurol* 1994;51:705–10.
- 131 Beaumont A, Burton AR, Lemon J, *et al.* Reduced cardiac vagal modulation impacts on cognitive performance in chronic fatigue syndrome. *PLoS One* 2012;7:e49518.
- 132 Weiß S, Winkelmann A, Duschek S. Recognition of facially expressed emotions in patients with fibromyalgia syndrome. *Behav Med* 2013;39:146–54.
- 133 Duschek S, Montoro CI, Reyes Del Paso GA. Diminished Interoceptive Awareness in Fibromyalgia Syndrome. *Behav Med* 2017;43:100–7.
- 134 Demartini B, Petrochilos P, Ricciardi L, *et al.* The role of alexithymia in the development of functional motor symptoms (conversion disorder). *J Neurol Neurosurg Psychiatry* 2014;85:1132–7.
- 135 Schönenberg M, Jusyte A, Höhnle N, *et al.* Theory of mind abilities in patients with psychogenic nonepileptic seizures. *Epilepsy Behav* 2015;53:20–4.
- 136 Stewart RS, Lovitt R, Stewart RM. Are hysterical seizures more than hysteria? A research diagnostic criteria, DMS-III, and psychometric analysis. *Am J Psychiatry* 1982;139:926–9.
- 137 Gul A, Ahmad H. Cognitive deficits and emotion regulation strategies in patients with psychogenic nonepileptic seizures: a task-switching study. *Epilepsy Behav* 2014;32:108–13.
- 138 Bakvis P, Roelofs K, Kuyk J, *et al.* Trauma, stress, and preconscious threat processing in patients with psychogenic nonepileptic seizures. *Epilepsia* 2009;50:1001–11.
- 139 Roberts NA, Bursellon MH, Weber DJ, *et al.* Emotion in psychogenic nonepileptic seizures: responses to affective pictures. *Epilepsy Behav* 2012;24:107–15.
- 140 Fuentes K, Hunter MA, Strauss E, *et al.* Intraindividual variability in cognitive performance in persons with chronic fatigue syndrome. *Clin Neuropsychol* 2001;15:210–27.
- 141 Wallman KE, Morton AR, Goodman C, *et al.* Reliability of physiological, psychological and cognitive variables in chronic fatigue syndrome and the role of graded exercise. *J Sports Sci Med* 2005;4:463–71.
- 142 Glass JM, Park DC, Minear M, *et al.* Memory beliefs and function in fibromyalgia patients. *J Psychosom Res* 2005;58:263–9.
- 143 Kane RL, Gantz NM, DiPino RK. Neuropsychological and psychological functioning in chronic fatigue syndrome. *Neuropsychiatry Neuropsychol Behav Neurol* 1997;10:25–31.
- 144 Siegel SD, Antoni MH, Fletcher MA, *et al.* Impaired natural immunity, cognitive dysfunction, and physical symptoms in patients with chronic fatigue syndrome: preliminary evidence for a subgroup? *J Psychosom Res* 2006;60:559–66.
- 145 Iverson GL, Le Page J, Koehler BE, *et al.* Test of Memory Malingering (TOMM) scores are not affected by chronic pain or depression in patients with fibromyalgia. *Clin Neuropsychol* 2007;21:532–46.
- 146 Gervais RO, Russell AS, Green P, *et al.* Effort testing in patients with fibromyalgia and disability incentives. *J Rheumatol* 2001;28:1892–9.
- 147 Brooks L, Johnson-Greene D, Lattie E, *et al.* The relationship between performances on neuropsychological symptom validity testing and the MCMI-III in patients with fibromyalgia. *Clin Neuropsychol* 2012;26:816–31.
- 148 Johnson-Greene D, Brooks L, Ference T. Relationship between performance validity testing, disability status, and somatic complaints in patients with fibromyalgia. *Clin Neuropsychol* 2013;27:148–58.
- 149 Schmalzing KB, DiClementi JD, Cullum CM, *et al.* Cognitive functioning in chronic fatigue syndrome and depression: a preliminary comparison. *Psychosom Med* 1994;56:383–8.
- 150 Vollmer-Conna U, Wakefield D, Lloyd A, *et al.* Cognitive deficits in patients suffering from chronic fatigue syndrome, acute infective illness or depression. *British Journal of Psychiatry* 1997;171:377–81.
- 151 van Beilen M, Griffioen BT, Gross A, *et al.* Psychological assessment of malingering in psychogenic neurological disorders and non-psychogenic neurological disorders: relationship to psychopathology levels. *Eur J Neurol* 2009;16:1118–23.
- 152 Binder LM, Salinsky MC, Smith SP. Psychological correlates of psychogenic seizures. *J Clin Exp Neuropsychol* 1994;16:524–30.
- 153 Drane DL, Williamson DJ, Stroup ES, *et al.* Cognitive impairment is not equal in patients with epileptic and psychogenic nonepileptic seizures. *Epilepsia* 2006;47:1879–86.
- 154 Kemp S, Coughlan AK, Rowbottom C, *et al.* The base rate of effort test failure in patients with medically unexplained symptoms. *J Psychosom Res* 2008;65:319–25.
- 155 Cragar DE, Berry DT, Fakhoury TA, *et al.* Performance of patients with epilepsy or psychogenic non-epileptic seizures on four measures of effort. *Clin Neuropsychol* 2006;20:552–66.
- 156 Gulec H. Normalizing attributions may contribute to non-help-seeking behavior in people with fibromyalgia syndrome. *Psychosomatics* 2008;49:212–7.
- 157 Molnar DS, Flett GL, Sadava SW, *et al.* Perfectionism and health functioning in women with fibromyalgia. *J Psychosom Res* 2012;73:295–300.
- 158 Kempke S, Luyten P, Claes S, *et al.* Self-critical perfectionism and its relationship to fatigue and pain in the daily flow of life in patients with chronic fatigue syndrome. *Psychol Med* 2013;43:995–1002.
- 159 Lawrie SM, MacHale SM, Power MJ, *et al.* Is the chronic fatigue syndrome best understood as a primary disturbance of the sense of effort? *Psychol Med* 1997;27:995–9.
- 160 Shenhav A, Musslick S, Lieder F, *et al.* Toward a Rational and Mechanistic Account of Mental Effort. *Annu Rev Neurosci* 2017;40:99–124.
- 161 Pareés I, Kassaveti P, Saifee TA, *et al.* Failure of explicit movement control in patients with functional motor symptoms. *Mov Disord* 2013;28:517–23.
- 162 Pessiglione M, Vinckier F, Bouret S, *et al.* Why not try harder? Computational approach to motivation deficits in neuro-psychiatric diseases. *Brain J Neurol* 2017.
- 163 Lishman WA. Physiogenesis and psychogenesis in the 'post-concussional syndrome'. *British Journal of Psychiatry* 1988;153:460–9.
- 164 Ruff RM. Mild traumatic brain injury and neural recovery: rethinking the debate. *NeuroRehabilitation* 2011;28:167–80.
- 165 Bohnen N, Jolles J, Twijnstra A. Neuropsychological deficits in patients with persistent symptoms six months after mild head injury. *Neurosurgery* 1992;30:692–5.
- 166 Bohnen N, Twijnstra A, Jolles J. Performance in the Stroop color word test in relationship to the persistence of symptoms following mild head injury. *Acta Neurol Scand* 1992;85:116–21.
- 167 Cicerone KD, Azulay J. Diagnostic utility of attention measures in postconcussion syndrome. *Clin Neuropsychol* 2002;16:280–9.
- 168 Hinton-Bayre AD, Geffen G, McFarland K. Mild head injury and speed of information processing: a prospective study of professional rugby league players. *J Clin Exp Neuropsychol* 1997;19:275–89.
- 169 Ferrari R, Russell AS, Carroll LJ, *et al.* A re-examination of the whiplash associated disorders (WAD) as a systemic illness. *Ann Rheum Dis* 2005;64:1337–42.
- 170 Robinson JP, Burwinkle T, Turk DC. Perceived and actual memory, concentration, and attention problems after whiplash-associated disorders (grades I and II): prevalence and predictors. *Arch Phys Med Rehabil* 2007;88:774–9.
- 171 Haldorsen T, Waterloo K, Dahl A, *et al.* Symptoms and cognitive dysfunction in patients with the late whiplash syndrome. *Appl Neuropsychol* 2003;10:170–5.
- 172 Kessels RP, Aleman A, Verhagen WL, *et al.* Cognitive functioning after whiplash injury: a meta-analysis. *J Int Neuropsychol Soc* 2000;6:271–8.
- 173 Seo J, Kim SH, Kim YT, *et al.* Working memory impairment in fibromyalgia patients associated with altered frontoparietal memory network. *PLoS One* 2012;7:e37808.
- 174 Glass JM, Williams DA, Fernandez-Sanchez ML, *et al.* Executive function in chronic pain patients and healthy controls: different cortical activation during response inhibition in fibromyalgia. *J Pain* 2011;12:1219–29.
- 175 Martinsen S, Flodin P, Berrebi J, *et al.* Fibromyalgia patients had normal distraction related pain inhibition but cognitive impairment reflected in caudate nucleus and hippocampus during the Stroop Color Word Test. *PLoS One* 2014;9:e108637.
- 176 Caseras X, Mataix-Cols D, Giampietro V, *et al.* Probing the working memory system in chronic fatigue syndrome: a functional magnetic resonance imaging study using the n-back task. *Psychosom Med* 2006;68:947–55.
- 177 Lange G, Steffener J, Cook DB, *et al.* Objective evidence of cognitive complaints in Chronic Fatigue Syndrome: a BOLD fMRI study of verbal working memory. *NeuroImage* 2005;26:513–24.
- 178 Fischler B, D'Haenen H, Cluydts R, *et al.* Comparison of 99m Tc HMPAO SPECT scan between chronic fatigue syndrome, major depression and healthy controls: an exploratory study of clinical correlates of regional cerebral blood flow. *Neurocytobiology* 1996;34:175–83.
- 179 Smits M, Dippel DW, Houston GC, *et al.* Postconcussion syndrome after minor head injury: brain activation of working memory and attention. *Hum Brain Mapp* 2009;30:2789–803.
- 180 Mittenberg W, DiGiulio DV, Perrin S, *et al.* Symptoms following mild head injury: expectation as aetiology. *J Neurol Neurosurg Psychiatry* 1992;55:200–4.
- 181 Ferguson RJ, Mittenberg W, Barone DF, *et al.* Postconcussion syndrome following sports-related head injury: expectation as etiology. *Neuropsychology* 1999;13:582–9.
- 182 Suhr J, Spickard B. Pain-related fear is associated with cognitive task avoidance: exploration of the cogniphobia construct in a recurrent headache sample. *Clin Neuropsychol* 2012;26:1128–41.
- 183 Sluka KA, Clauw DJ. Neurobiology of fibromyalgia and chronic widespread pain. *Neuroscience* 2016;338:114–29.

- 184 Sharif K, Watad A, Bragazzi NL, *et al.* On chronic fatigue syndrome and nosological categories. *Clin Rheumatol* 2018.
- 185 White PD, Goldsmith KA, Johnson AL, *et al.* PACE trial management group. Comparison of adaptive pacing therapy, cognitive behaviour therapy, graded exercise therapy, and specialist medical care for chronic fatigue syndrome (PACE): a randomised trial. *Lancet* 2011;377:823–36.
- 186 Nielsen G, Buszewicz M, Stevenson F, *et al.* Randomised feasibility study of physiotherapy for patients with functional motor symptoms. *J Neurol Neurosurg Psychiatry* 2017;88:484–90.
- 187 Goldstein LH, Chalder T, Chigwedere C, *et al.* Cognitive-behavioral therapy for psychogenic nonepileptic seizures: a pilot RCT. *Neurology* 2010;74:1986–94.