

Therapeutic deep brain stimulation worsening dysprosody in Parkinson's disease – an unexplored entity

Maria Shoaib¹

Muhammad Taimoor Khan²

¹Department of Medicine, Dow Medical College, Karachi, Pakistan;

²Department of Neurology, Charleston Area Medical Center/ West Virginia University, Charleston, WV, USA

Dear editor

We read this article, “Altered emotional recognition and expression in patients with Parkinson's disease” by Jin et al¹ with great interest and appreciate the novel information provided on altered emotional processing in pre-deep brain stimulation (DBS) Parkinson's disease (PD) patients and we would like to add our feedback on the role of DBS on nonmotor and emotional components of PD.

PD is considered a movement disorder primarily presenting with motor symptoms like resting tremors, bradykinesia, rigidity, and postural instability. However, several nonmotor symptoms such as cognitive impairment, dementia, depression, anxiety, and apathy are also noted. Nonmotor symptoms may even dominate the clinical picture as disease progresses.²

The subthalamic nucleus (STN), in particular basal ganglia and amygdala, has an important role in the recognition of basic facial emotions both from face and music. Results from both pathophysiological and neuroimaging studies suggested this, as well as the fact that its involvement in PD frequently impairs the recognition of disgust and fear emotions, followed by sadness and anger.³

DBS treats motor symptoms, such as tremor, rigidity and bradykinesia, but may worsen certain nonmotor symptoms, including articulation of words and expressing emotions, resulting in dysprosody in the patients. The clinical evaluation of speech after 1 and 3 years of STN-DBS in PD patients showed a significant deterioration of speech. Articulation rather than voice was most frequently affected, with a distinct dysarthria. The mechanisms by which DBS exacerbates dysarthria while improving other symptoms could be partly attributed to medially placed electrodes and high amplitude of stimulation, but this is still unclear.⁴ However, due to obvious ethical and experimental boundaries in human studies, recent observations that hypokinetic or hyperkinetic dysarthria may be worsened by otherwise therapeutic DBS are largely unexplored.

King et al⁵ presented a rodent model of Parkinsonian dysarthria that is worsened by DBS, thus providing a stepping stone for future research. They proposed this study using the existing 6-hydroxydopamine rodent model of Parkinsonism to observe hypokinetic dysarthria and its exacerbation by DBS. Stimulation of the STN aggravated these Parkinsonian vocalization symptoms. Vocalization complexity and rates, word durations, and individual sound bandwidths were all detrimentally compounded

Correspondence: Maria Shoaib
Department of Medicine, Dow Medical College, 176/Y Block 2 PECHS Karachi, Karachi, Pakistan
Tel +92 333 328 9909
Email syedamariashoaib@gmail.com

after DBS. The rats generated scant and shorter vocalizations that were less acoustically complex than those under the diseased condition.

This model will provide a framework for exploration into the electrophysiological mechanisms of Parkinsonian hypokinetic dysarthria and its exacerbation after DBS. Knowing how DBS therapy can intensify dysarthria may lead researchers to devise alternative approaches to treat motor and speech signs together. Such explorations are essential to minimize this side effect of DBS that worsens dysarthria, exacerbating Parkinsonian speech deficits that can greatly diminish patients' emotional recognition, expression, and quality of life.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Jin Y, Mao Z, Ling Z, Xu X, Zhang Z, Yu X. Altered emotional recognition and expression in patients with Parkinson's disease. *Neuropsychiatr Dis Treat*. 2017;13:2891–2902.
2. Sollinger AB, Goldstein FC, Lah JJ, Levey AI, Factor SA. Mild cognitive impairment in Parkinson's disease: subtypes and motor characteristics. *Parkinsonism Relat Disord*. 2010;16(3):177–180.
3. Saenz A, Doé de Maindreville A, Henry A, Labbey S, Bakchine S, Ehrlé N. Recognition of facial and musical emotions in Parkinson's disease. *Eur J Neurol*. 2013;20(3):571–577.
4. Tripoliti E. *Effects of Deep Brain Stimulation on Speech in Patients with Parkinson's Disease and Dystonia* [dissertation]. London: UCL (University College London); 2011.
5. King NO, Anderson CJ, Dorval AD. Deep brain stimulation exacerbates hypokinetic dysarthria in a rat model of Parkinson's disease. *J Neurosci Res*. 2016;94(2):128–138.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Neuropsychiatric Disease and Treatment 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Neuropsychiatric Disease and Treatment editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS,

and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>