

Autotransplantation: A biological treatment alternative for a patient after traumatic dental injury

Meenakshi Vishwanath^{a,b}
Nandakumar Janakiraman^c
Hamed Vaziri^d
Ravindra Nanda^a
Flavio Uribe^a

^aDivision of Orthodontics, School of Dental Medicine, University of Connecticut, Farmington, CT, USA

^bDepartment of Growth and Development, Orthodontic Section, College of Dentistry, University of Nebraska Medical Center, Lincoln, NE, USA

^cDivision of Orthodontics, School of Dentistry, University of Louisville, Louisville, KY, USA

^dPrivate Practice, Spring, TX, USA

Traumatic dental injury is considered a public dental health problem because of a high childhood incidence, high treatment costs, and prolonged treatment time. Although management guidelines for traumatized teeth have been outlined, tooth loss following trauma is occasionally unavoidable. Here, we describe the successful interdisciplinary management of a traumatized central incisor in an 11-year old boy that was extracted because of a poor prognosis and restored by the autotransplantation of an immature donor tooth into the site. The patient underwent orthodontic treatment in order to close the donor site space and bring the autotransplanted tooth to an ideal position. Postorthodontic treatment radiographs and photographs revealed an esthetic and functional natural tooth replacing the lost tooth. The findings from this case suggest that autotransplantation offers unique advantages as a treatment modality for the restoration of missing teeth, particularly in growing children.

[Korean J Orthod 2018;48(2):125-130]

Key words: Transplantation, Orthodontic mini-implant, Growth and development, Dental trauma, Autotransplantation

Received January 17, 2017; Revised June 1, 2017; Accepted July 17, 2017.

Corresponding author: Meenakshi Vishwanath.

Assistant Professor, Department of Growth and Development, Orthodontic Section, College of Dentistry, University of Nebraska Medical Center, 4000 East Campus Loop South, Lincoln, NE 68583-0740, USA.

Tel +1-402-472-1302 **e-mail** minnie.vishwanath@unmc.edu

The authors report no commercial, proprietary, or financial interest in the products or companies described in this article.

© 2018 The Korean Association of Orthodontists.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Traumatic dental injury (TDI) is a public dental health problem because of a high frequency of occurrence, a high incidence during childhood, and associated treatment costs. Moreover, it often carries a lifelong treatment burden for the patient.¹ Specific population studies have shown that trauma to the oral region can account for 5% of all injuries, with the highest risk in children aged 0 to 12 years.² Guidelines for the management of traumatized teeth have been clearly outlined³; however, despite the dentist's best effort, tooth loss following trauma is sometimes inevitable. In order to select the best treatment option for patients with a lost anterior tooth, the clinician must consider several factors such as the patient's growth status, success rates and costs of various treatment options, patient preferences, and, most importantly, treatment options that will enable long-term rehabilitation.

Tooth autotransplantation to replace lost or missing teeth has been attempted for several centuries. Even early on, it became clear that there were two critically important factors for success. These included the use of an immature donor tooth for transplantation and the immediate transplantation of the immature tooth into the recipient site following extraction.⁴ Some of the early published works on autotransplantation appeared in the 1950s,^{5,6} followed by the case series by Slagsvold and Bjercke.⁷ Extensive animal studies and long-term prospective studies by Andreasen et al.⁸⁻¹⁰ have served as stepping stones for the development of this treatment as a viable alternative for tooth substitution.

In a growing individual, the replacement of a lost an-

terior tooth by autotransplantation becomes a logical choice because it allows for continuous skeletal growth and the preservation of a vital periodontium.¹¹ Dental implants, on the other hand, require long periods of space maintenance until the cessation of growth. Once the biological principles behind successful autotransplantation have been understood and the indications have been defined, it can be considered an extremely successful treatment modality with significant savings in time and cost compared with implants.¹²

Here we report the successful outcome of autotransplantation used in conjunction with orthodontic treatment for the replacement of a lost central incisor in a growing child who experienced trauma to the anterior maxilla. The uniqueness of this case is further emphasized by the unconventional decision to use the donor tooth from the arch that already had a congenitally missing tooth.

ETIOLOGY AND DIAGNOSIS

An 11-year-old boy was referred to the orthodontic clinic from the endodontic clinic for the interdisciplinary management of a traumatized anterior tooth. Dental history revealed a recent bicycle accident that resulted in trauma to the maxillary right central incisor (#11). Immediately after the injury was sustained, the patient was rushed to Pediatric Dentistry Clinic of University of Connecticut, where examination demonstrated a suspected coronal–radicular fracture as well as mobility of tooth #11. The maxillary anterior teeth were splinted with fiber-reinforced composite, and the patient was referred to the Endodontic clinic for further management.




Figure 1. Pretreatment photographs and imaging findings for an 11-year-old boy with a traumatized maxillary central incisor. A, Intraoral photograph shows the crown of the maxillary right central incisor with a vertical fracture line extending onto the labial surface. B, Intraoral photograph shows the crown of the maxillary right central incisor with a vertical fracture line extending onto the palatal surface. C, Cone-beam computed tomography (CBCT) reveals the presence of a horizontal fracture at the cemento-enamel junction/coronal portion of the root of the maxillary right central incisor. D, A panoramic radiograph reconstructed from the CBCT data reveals the presence of a vertical crown fracture as well as the root morphology and developmental stage of the mandibular right second premolar.

The splint was removed after 3 weeks, when clinical examination revealed a fracture line extending onto the labial and palatal surfaces of tooth #11 (Figure 1A and 1B). For further investigation, cone-beam computed tomography (CBCT) was performed, which confirmed the presence of a vertical crown fracture and also revealed a horizontal fracture at the cementoenamel junction/coronal portion of the root (Figure 1C). Because the prognosis of the fractured incisor was poor, the interdisciplinary team recommended the extraction of tooth #11.

TREATMENT OBJECTIVES

The primary objective was to determine a treatment plan that involved temporary or permanent replacement of tooth #11, which was lost because of trauma, as well as the restoration of esthetics in the maxillary anterior region. The secondary objective was to maintain the maxillary incisor inclination and/or obtain good functional occlusion.

TREATMENT ALTERNATIVES

The following treatment alternatives were considered.

1. Dental implantation for the replacement of tooth #11. This would involve initial orthodontic treatment to manage the space of tooth #11 and position the adjacent tooth roots ideally for the future implant and fixed retention across the edentulous space to prevent tipping of the adjacent roots,

along with a temporary pontic for tooth #11 for esthetics and space maintenance.

2. Autotransplantation of a mandibular premolar (#45) into the region of the extracted tooth, followed by orthodontic treatment to close all spaces and obtain good functional occlusion. Restorative and esthetic recontouring of the autotransplanted tooth would be performed at the end of orthodontic treatment.

The patient's parents and the interdisciplinary team opted for the second alternative, because it was considered to provide the best long-term prognosis and would consequently prove the most beneficial for the patient.

TREATMENT PROGRESS

The fractured central incisor was extracted and the recipient site was prepared to receive the donor tooth. Subsequently, tooth #45 was atraumatically extracted, placed in the recipient site, and splinted with a 0.020-inch stainless steel wire that was bent passively to fit the labial contour of the tooth and was extended to be engaged into the bracket slots on the adjacent maxillary incisors (Figure 2A and 2B). The patient reported to the orthodontic clinic 4 weeks after the autotransplantation procedure. At that time, orthodontic examinations and records were completed. Intraorally, the patient was in the late mixed dentition stage with a congenitally missing mandibular central incisor, Class 1 molar relationship, and normal overjet and overbite. The panoramic radiograph showed alveolar bone remodeling around the




Figure 2. Extraction and autotransplantation of the fractured maxillary central incisor. A, Atraumatic extraction of the mandibular right second premolar (donor tooth). B, Autotransplantation of the mandibular right second premolar into the socket of the maxillary right central incisor. The autotransplanted tooth is splinted to the adjacent incisors using a 0.020-inch stainless steel wire. C, Pretreatment panoramic radiograph shows alveolar bone remodeling around the autotransplanted tooth (tooth #45 in the socket of tooth #11) and the edentulous space of the donor tooth. D, A stereolithographic model of the donor tooth.

autotransplanted tooth and the edentulous space of the donor tooth (Figure 2C). Cephalometric analysis showed a mild Class III skeletal relationship (A point, nasion, B point [ANB] = 0°), an upper incisor inclination of 105°, and a nasolabial angle of 104°.

Specific orthodontic treatment objectives involved closure of the donor tooth space and movement of the autotransplanted tooth into the ideal position so that it could be restored to achieve the best esthetic outcome. The final plan involved the orthodontic closure of all spaces to eliminate the need for future dental implants, followed by restorative build-ups.

Three months after autotransplantation, 0.022-inch MBT™ prescription metal brackets were bonded on the maxillary and mandibular teeth and initial archwires were

placed. To achieve absolute protraction of the mandibular first molar, which was required to achieve a Class III functional occlusion, a mini-implant (2.0 × 9 mm Lomas; Mondeal Medical Systems GmbH, Donau, Germany) was placed on the alveolar ridge distal to the mandibular right first premolar (Figure 3).

RESULTS

The total treatment duration was 36 months. The autotransplanted tooth was esthetically restored during the finishing phase of the orthodontic treatment; this also helped in achieving good functional occlusion at the end of the orthodontic treatment (Figure 4A). The patient and his parents were pleased with the




Figure 3. Orthodontic treatment in progress. A, Autotransplanted tooth bonded and aligned with the contralateral central incisor. B, A temporary anchorage device is placed on the ridge for closure of the donor space through protraction of the right mandibular molar. C, Maxillary arch held passively to allow for eruption of maxillary canines bilaterally. Space closure in progress in the mandibular arch.



Figure 4. Intraoral photographs and a panoramic radiograph obtained after autotransplantation and orthodontic treatment. A, Intraoral photographs show functional occlusion and the postrestorative findings for the autotransplanted tooth (tooth #45 in the socket of tooth #11). B, Post-treatment panoramic radiograph showing complete root formation of the autotransplanted tooth.

results, considering the interdisciplinary approach had resulted in the esthetically pleasing replacement of the lost central incisor. The post-treatment panoramic radiograph showed that the root formation had progressed to achieve a full length (Figure 4B). Post-orthodontic evaluation of the autotransplanted tooth demonstrated all signs of a successful transplant,¹³ which include a good crown to root ratio, no external root resorption or mobility, and, most importantly, an appearance that is identical to the counterpart, with a matching gingival contour and emergence profile.

DISCUSSION

We reported the successful outcome of autotransplantation used in conjunction with orthodontic treatment for the replacement of a lost central incisor in a growing child who experienced trauma to the anterior maxilla. The novelty of our case is that, autotransplantation was done in a patient who did not meet the typical criteria. Traditionally, autotransplantation is considered, when extraction in one of the arches is indicated as part of orthodontic treatment planning; as in, there is requirement for additional space so as to relieve crowding or address proclined teeth. Our case demonstrates that, even if the strictest criteria for autotransplantation are not met, it should not deter the orthodontic professionals from wanting to use this option as we are in a generation where we can use tools like three-dimensional CBCT and temporary anchorage devices to ensure a more predictable outcomes.

The treatment of TDI in a young patient is often unpredictable, complicated, expensive, and can continue throughout his or her life; therefore, the treatment of choice should minimize these undesired consequences.¹⁴ The treatment options include removable or conventional fixed prostheses, dental implant-supported crowns, autotransplantation, and simple orthodontic space closure with tooth substitution.¹⁵

Currently, an implant-supported restoration is considered the gold standard for the conservative replacement of a lost tooth in adult patients.¹⁵ In a growing child, this treatment modality generally necessitates space maintenance until facial growth is completed. During this time, there is continued alveolar bone remodeling in terms of both width and height; this may require alveolar bone grafting procedures and consequently increase the total cost of treatment. Furthermore, a long-term study demonstrated that the cessation of vertical growth is not very predictable, with mature adults exhibiting vertical steps after the placement of anterior implant restorations to the same extent as patients in the late adolescent stage.¹⁶ Autotransplantation of teeth can be an excellent treatment alternative to overcome the above

concerns in young adolescents. The successful transplantation of a premolar along with healthy surrounding tissues in our patient confirms that this treatment modality can be recommended for the replacement of missing maxillary incisors in young patients.

Successful tooth autotransplantation results in the maintenance of a viable periodontal ligament and, consequently, the potential for bone induction and the capacity for continuous eruption during growth, leading to the re-establishment of a normal alveolar process.¹¹ Because autotransplantation is a technique- and operator-sensitive procedure, it has not been established as a standard-of-care for the replacement of lost teeth. This may also be the reason for the considerably varied success and survival rates for autotransplanted teeth reported in the literature. An overall comparison of the survival rates for autotransplanted teeth with those for single-tooth osseointegrated implants demonstrated comparable results. Czochrowska et al.¹⁷ showed a 90% survival rate for autotransplanted teeth over a follow-up period of 17 to 41 years and a 94.5% survival rate for single-tooth implants over a period of 5 years.¹⁸

CONCLUSION

The findings from our case suggest that tooth autotransplantation is an excellent biological treatment alternative for the restoration of missing teeth, particularly in growing children.

REFERENCES

1. Glendor U. Epidemiology of traumatic dental injuries--a 12 year review of the literature. *Dent Traumatol* 2008;24:603-11.
2. Petersson EE, Andersson L, Sörensen S. Traumatic oral vs non-oral injuries. *Swed Dent J* 1997;21:55-68.
3. Andreasen JO. The dental trauma guide [Internet]. Copenhagen: Copenhagen University Hospital and the International Association of Dental Traumatology (IADT); 2010 [cited 2016 Feb 28]. Available from: <http://dentaltraumaguide.org>.
4. Cross D, El-Angbawi A, McLaughlin P, Keightley A, Brocklebank L, Whitters J, et al. Developments in autotransplantation of teeth. *Surgeon* 2013;11:49-55.
5. Hale ML. Autogenous transplants. *J Am Dent Assoc* 1954;49:193-8.
6. Miller HM. Transplantation and reimplantation of teeth. *Oral Surg Oral Med Oral Pathol* 1956;9:84-95.
7. Slagsvold O, Bjercke B. Autotransplantation of premolars with partly formed roots. A radiographic study of root growth. *Am J Orthod* 1974;66:355-

- 66.
8. Andreasen JO, Kristerson L. Evaluation of different types of autotransplanted connective tissues as potential periodontal ligament substitutes. An experimental replantation study in monkeys. *Int J Oral Surg* 1981;10:189-201.
 9. Andreasen JO. Periodontal healing after replantation and autotransplantation of incisors in monkeys. *Int J Oral Surg* 1981;10:54-61.
 10. Andreasen JO, Paulsen HU, Yu Z, Ahlquist R, Bayer T, Schwartz O. A long-term study of 370 autotransplanted premolars. Part I. Surgical procedures and standardized techniques for monitoring healing. *Eur J Orthod* 1990;12:3-13.
 11. Park JH, Tai K, Hayashi D. Tooth autotransplantation as a treatment option: a review. *J Clin Pediatr Dent* 2010;35:129-35.
 12. Tsukiboshi M. Autotransplantation of teeth: requirements for predictable success. *Dent Traumatol* 2002;18:157-80.
 13. Czochrowska EM, Stenvik A, Album B, Zachrisson BU. Autotransplantation of premolars to replace maxillary incisors: a comparison with natural incisors. *Am J Orthod Dentofacial Orthop* 2000;118:592-600.
 14. Zaleckiene V, Peciuliene V, Brukiene V, Drukteinis S. Traumatic dental injuries: etiology, prevalence and possible outcomes. *Stomatologija* 2014;16:7-14.
 15. Chesterman J, Chauhan R, Patel M, Chan MF. The management of traumatic tooth loss with dental implants: part 1. *Br Dent J* 2014;217:627-33.
 16. Bernard JP, Schatz JP, Christou P, Belser U, Kiliaridis S. Long-term vertical changes of the anterior maxillary teeth adjacent to single implants in young and mature adults. A retrospective study. *J Clin Periodontol* 2004;31:1024-8.
 17. Czochrowska EM, Stenvik A, Bjercke B, Zachrisson BU. Outcome of tooth transplantation: survival and success rates 17-41 years posttreatment. *Am J Orthod Dentofacial Orthop* 2002;121:110-9; quiz 193.
 18. Jung RE, Pjetursson BE, Glauser R, Zembic A, Zwahlen M, Lang NP. A systematic review of the 5-year survival and complication rates of implant-supported single crowns. *Clin Oral Implants Res* 2008;19:119-30.