

# Implications of miR-148a-3p/p35/PTEN signaling in tau hyperphosphorylation and autoregulatory feedforward of Akt/CREB in Alzheimer's disease

Li Zeng, Hailun Jiang, Ghulam Md Ashraf, Jianghong Liu, Linlin Wang, Kaiyue Zhao, Mimin Liu, Zhuorong Li, and Rui Liu

Correspondence: [lizhuorong@imb.pumc.edu.cn](mailto:lizhuorong@imb.pumc.edu.cn), [liurui@imb.pumc.edu.cn](mailto:liurui@imb.pumc.edu.cn)  
<https://doi.org/10.1016/j.omtn.2023.102059>

(Molecular Therapy: Nucleic Acids 27, 256–275; March 2022)

In the originally published version of this article, in Figure 9C, the fluorescence image of PTEN and the merged image of PTEN with phosphorylated tau PHF-1 in the hippocampus of WT control group were inadvertently duplicated for those of the PTEN siRNA-treated APP/PS1 group. The figure has been corrected to show the actual images. These changes have no impact on the results and conclusions presented in this paper. The authors regret this error.




Figure 9C. miR-148a-3p overexpression or PTEN silencing protects miR-148a-3p/p35/PTEN signaling and PTEN/Akt/CREB feedforward in AD mice (original)




Figure 9C. miR-148a-3p overexpression or PTEN silencing protects miR-148a-3p/p35/PTEN signaling and PTEN/Akt/CREB feedforward in AD mice (corrected)

