

Original Article
Medicine General & Policy



Evaluating the Outcome of Multi-Morbid Patients Cared for by Hospitalists: a Report of Integrated Medical Model in Korea

Jung Hwan Lee ,¹ Ah Jin Kim ,¹ Tae Young Kyong ,² Ji-Hun Jang ,¹ Jeongmi Park ,¹ Jeong Hoon Lee ,¹ Man-Jong Lee ,¹ Jung-Soo Kim ,¹ Young Ju Suh ,³ Seong-Ryul Kwon ,⁴ and Cheol-Woo Kim ⁴

OPEN ACCESS

Received: Mar 13, 2019

Accepted: Jun 6, 2019

Address for Correspondence:

Ah Jin Kim, MD

Department of Hospital Medicine, Inha University Hospital, Inha University School of Medicine, 27 Inhang-ro, Jung-gu, Incheon 22332, Republic of Korea.

E-mail: emjin23@naver.com

© 2019 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Jung Hwan Lee

<https://orcid.org/0000-0001-7567-0664>

Ah Jin Kim

<https://orcid.org/0000-0001-5689-5227>

Tae Young Kyong

<https://orcid.org/0000-0003-4106-9306>

Ji-Hun Jang

<https://orcid.org/0000-0002-4946-2199>

Jeongmi Park

<https://orcid.org/0000-0003-3206-4589>

Jeong Hoon Lee

<https://orcid.org/0000-0002-8159-8192>

Man-Jong Lee

<https://orcid.org/0000-0002-4533-2629>

Jung-Soo Kim

<https://orcid.org/0000-0001-6603-6768>

Young Ju Suh

<https://orcid.org/0000-0002-6816-8067>

¹Department of Hospital Medicine, Inha University Hospital, Inha University School of Medicine, Incheon, Korea

²Department of Hospital Medicine, Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul, Korea

³Department of Biomedical Sciences, Inha University Hospital, Inha University School of Medicine, Incheon, Korea

⁴Department of Internal Medicine, Inha University Hospital, Inha University School of Medicine, Incheon, Korea

ABSTRACT

Background: The lack of medical personnel has led to the employment of hospitalists in Korean hospitals to provide high-quality medical care. However, whether hospitalists' care can improve patients' outcomes remains unclear. We aimed to analyze the outcome in patients cared for by hospitalists.

Methods: A retrospective review was conducted in 1,015 patients diagnosed with pneumonia or urinary tract infection from March 2017 to July 2018. After excluding 306 patients, 709 in the general ward who were admitted via the emergency department were enrolled, including 169 and 540 who were cared for by hospitalists (HG) and non-hospitalists (NHG), respectively. We compared the length of hospital stay (LOS), in-hospital mortality, readmission rate, comorbidity, and disease severity between the two groups. Comorbidities were analyzed using Charlson comorbidity index (CCI).


Results: HG LOS (median, interquartile range [IQR], 8 [5–12] days) was lower than NHG LOS (median [IQR], 10 [7–15] days), ($P < 0.001$). Of the 30 (4.2%) patients who died during their hospital stay, a lower percentage of HG patients (2.4%) than that of NHG patients (4.8%) died, but the difference between the two groups was not significant ($P = 0.170$). In a subgroup analysis, HG LOS was shorter than NHG LOS (median [IQR], 8 [5–12] vs. 10 [7–16] days, respectively, $P < 0.001$) with CCI of ≥ 5 points.

Conclusion: Hospitalist care can improve the LOS of patients, especially those with multiple comorbidities. Further studies are warranted to evaluate the impact of hospitalist care in Korea.

Keywords: Hospitalist; Length of Stay; Multiple Chronic Conditions

Seong-Ryul Kwon 

<https://orcid.org/0000-0003-1262-2790>

Cheol-Woo Kim 

<https://orcid.org/0000-0003-4306-1367>

Funding

This study was supported by a grant from the Inha University Hospital 2019.

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Lee JH,¹ Data curation:

Kyong TY, Jang JH, Park J, Lee JH,² Lee MJ.

Formal analysis: Kim AJ, Suh YJ. Investigation:

Kim JS. Methodology: Lee JH,¹ Kim AJ.

Software: Lee JH,¹ Validation: Lee JH,¹ Writing -

original draft: Lee JH,¹ Kim AJ. Writing - review

& editing: Kwon SR, Kim CW.

Lee JH,¹ Lee Jung Hwan; Lee JH,² Lee Jeong Hoon.

INTRODUCTION

Over the past two decades, hospitalists have emerged as medical specialists, worldwide; and are rapidly expanding and creating a new area of “hospital medicine.”¹⁻³ Through numerous studies, the benefits of hospitalists have been proven not only in patients in general medicine,^{4,5} but also in pediatrics,⁶ acute stroke,⁷ and surgical patients.⁸ The potential advantages of the hospitalist system include greater expertise in inpatient care and better availability of physicians during hospitalizations.⁹

In the United States, hospitalists have increased spontaneously in response to the need for medical efficiency and safety of inpatients, particularly for the elderly multi-morbid patients who have increased.¹⁰ Currently, more than 50,000 physicians (5% of all doctors) work as hospitalists.¹ In Korea, the need for inpatient care has led to the increased number of hospitalists due to the lack of medical personnel in tertiary hospitals.¹¹ Therefore, the hospitalist system, introduced institutionally in 2016, is supported by national reimbursement financial sources. Thus far, the training hospitals have recruited about 100 hospitalists nationwide. However, stability, identity, and rewards for hospitalists are not well-established. Hence, there are not enough hospitalists to spread to a lot of hospitals.^{12,13}

The acute medical care unit was first introduced in Korea. It involves the provision of short-term care to patients who present at the emergency departments (ED).¹⁴ The integrated medical model is a recently emerging model, which involves the collaboration of different subspecialists or hospitalists in order to provide quality care to patients in an independent department.¹³ A similar model with an integrated medical model has been used in foreign countries.¹⁵

Despite the high expectations of improvement in patients' outcomes, the effect of the hospitalist system remains unproven. Our institute has operated a hospitalist-centered system involving internal and surgical hospitalists, critical care team, and rapid response team since March 2017. This study therefore aimed to compare the outcome (including length of hospital stay [LOS], in-hospital mortality [IHM], and readmission rate) in patients with pneumonia or urinary tract infection (UTI) in the general wards admitted via the ED between hospitalists and non-hospitalists.

METHODS

Setting of the hospitalist system

The Department of Hospital Medicine at our institute consists of the internal medicine team (one gastroenterologist and one general internist), surgical team (three general surgeons and one obstetrician), and critical care team (one pulmonologist, one cardiologist, and one emergency medicine doctor). Two attending physicians from the internal medicine team independently cared for and discharged patients without serving as specialists. One of them had been serving as a physician for 17 years and had worked at a private local clinic and in a general hospital. The other doctor underwent 2-year fellowship training and 3 years of military service after obtaining the physician certificate. Both of them were the first to start as hospitalists. Most of the patients cared for by the hospitalists were admitted via the ED. Some patients were referred from the intensive care unit (ICU), while the others were from the outpatient clinics. Patients who were initially admitted in the ED were allocated mainly to the hospitalists, and those who had been followed up by subspecialists were allocated to

the non-hospitalists. Hospitalist care was performed during the daytime 6 days a week. Patients assigned to hospitalists were admitted through the ward where the hospitalists worked. However, this ward was shared with other medical patients cared for by non-hospitalists. Non-hospitalists inpatient care was done by subspecialists and residents. Pneumonia or UTI were mainly cared for in the pulmonology, infectious disease, nephrology, or urology units. However, the other divisions of internal medicine, neurology, rehabilitation medicine, or surgical department have also had to treat the disease. The median career period of non-hospitalists was 19 years (interquartile range [IQR] 7–24 years), since their specialty certification.

Patients

In total, 1,501 patients with pneumonia or UTI were admitted from March 2017 to July 2018 in our institute. We excluded 486 patients admitted from outpatient clinics. Moreover, 300 patients were excluded as they were admitted directly in the ICU from ED. Of the remaining patients, six who remained undischarged until the study cut-off date were also excluded. Finally, 709 patients admitted in the general ward during the study period were enrolled and dichotomized into two groups: 169 cared for by hospitalists and 540 cared for by non-hospitalists (Fig. 1). The hospitalists group (HG) was defined as inpatients that were under the care of attending physicians from the internal medicine team, while the non-hospitalists group (NHG) was defined as inpatients cared for by attending physicians from other medical departments.

Calculation of comorbidity and disease severity

Comorbidity and disease severity were considered because these can affect LOS, IHM, and readmission rate.¹⁶⁻¹⁹ Chronic comorbid conditions were identified through chart review and were then reported using the Charlson comorbidity index (CCI).²⁰⁻²² The CCI score was then classified into three previously defined grades: 0–2 points (low), 3–4 points (moderate), and ≥ 5 points (high).²³ Disease severity was measured by Mortality in Emergency Department Sepsis (MEDS) score, which is a superior predictive model used in validating a number of severe illnesses.²⁴⁻²⁶ The MEDS score was calculated as follows: terminal illness with possible death within 1 month (6 points), hypoxia or tachypnea (3 points), septic shock (3 points), platelet count $< 150,000$ cells/mm³ (3 points), band neutrophils $> 5\%$ of leukocyte (3 points), age > 65 years (3 points), lower respiratory tract infection (2 points), nursing home resident (2 points), and altered mental status (2 points). In the current study, the variable “terminal illness” was defined as metastatic cancer or disease conditions with at least 50% chance of mortality within 30 days of admission, including cases with “do not resuscitate” orders or those with orders for withdrawal of life-sustaining treatments.^{25,26} “Septic shock” was defined as persisting hypotension requiring vasopressors to maintain a mean arterial pressure of 65 mmHg or higher or having lactate levels of > 4 mmol/L.^{26,27} “Nursing home resident” were regarded as patients who were referred from long-term care hospitals. CURB-65 score, also used to consider the severity of pneumonia, was divided into 3 groups as follows: 0–1 points (low), 2 points (intermediate), and ≥ 3 points (high).²⁸

The LOS was defined as the duration of a single episode of hospitalization and was calculated by subtracting the day of admission from the day of discharge.¹⁴ IHM was defined as the ratio of inpatient deaths to the total number of inpatients.²⁹ Readmissions were identified as unscheduled admission via ED due to any cause within 14 or 30 days after discharge.

Statistical analysis

Baseline characteristics of the HG and NHG groups were described. Continuous variables were summarized as medians and IQR or means and standard deviations, and analyzed




Fig. 1. Selection of patients with pneumonia or UTI.
UTI = urinary tract infection.

using Mann-Whitney U test. For categorical variables, frequencies and percentages were determined, and differences in distribution were estimated using Pearson's χ^2 test. We performed subgroup analyses according to disease category, comorbidity (based on CCI), and severity (based on MEDS score). When the response variable was treated as a binary variable, logistic regression was used to predict whether the LOS exceeded nine days. Multiple linear regression was also used to determine whether CCI, MEDS score, gender, and hospitalist care were associated with LOS. The distribution of LOS was positively skewed, and the logarithm function was used to transform the variables into linear data.³⁰ *P* values < 0.05 were considered significant. We used the Bonferroni adjustment for multiple comparisons in the subgroup analyses. All statistical analyses were performed using SPSS 19.0 software (SPSS, Chicago, IL, USA).

Ethics statement

The Institutional Review Board of Inha University Hospital reviewed and approved this study (approval No. INHAUH 2019-01-003). Informed consent was waived due to the retrospective nature of the study.

RESULTS

Baseline characteristics

Table 1 presents the baseline characteristics of the HG and NHG groups. The percentage of women in the HG group was significantly higher than that in the NHG group (63.9% vs. 48.1%, respectively; $P < 0.001$). Compared to the HG group, the NHG group had higher proportion of patients with pneumonia (58.6% vs. 71.7%, respectively; $P = 0.001$). Among patients with comorbidities, those in the NHG group had higher proportions of patients with diabetes, chronic kidney disease, and chronic obstructive pulmonary disease than those in the HG group ($P = 0.030$, 0.030 , and 0.010 , respectively). However, there was no significant difference in CCI, which was used to summarize the comorbidities, between the two groups ($P = 0.650$). Disease severity in both groups was compared according to the MEDS score

Table 1. Baseline characteristics of patients cared for by hospitalists group and non-hospitalists group

Characteristics	Hospitalist (n = 169)	Non-hospitalist (n = 540)	P value
Age, yr	68.0 ± 18.2	69.9 ± 15.8	
< 50	26 (15.4)	57 (10.6)	0.11
50–59	23 (13.6)	56 (10.4)	
60–69	21 (12.4)	103 (19.1)	
70–79	42 (26.6)	159 (29.4)	
≥ 80	54 (32.0)	165 (30.6)	
Gender			< 0.001
Men	61 (36.1)	280 (51.9)	
Women	108 (63.9)	260 (48.1)	
Diagnosis			0.001
Pneumonia	99 (58.6)	387 (71.7)	
Urinary tract infection	70 (41.4)	153 (28.3)	
Comorbidities			
Diabetes			0.03
Uncomplicated	41 (24.3)	125 (23.1)	
Complicated	4 (2.4)	45 (8.3)	
Liver disease	8 (4.7)	26 (4.8)	0.97
Malignancy			0.31
Any leukemia, lymphoma, or localized solid tumor	2 (1.2)	12 (2.2)	
Metastatic solid tumor	1 (0.6)	11 (2.0)	
HIV	0 (0)	2 (0.4)	0.428
Chronic kidney disease	4 (2.4)	38 (7.0)	0.03
Congestive heart failure	20 (11.8)	70 (13.0)	0.70
PAOD	0 (0)	9 (1.7)	0.09
COPD	6 (3.6)	57 (10.6)	0.01
Cerebral vascular disease	32 (18.9)	106 (19.6)	0.84
Hemiplegia	1 (0.6)	9 (1.7)	0.30
Rheumatic disease	2 (1.2)	14 (2.6)	0.28
Dementia	17 (10.1)	49 (9.1)	0.70
Peptic ulcer disease	2 (1.2)	3 (0.6)	0.40
Charlson comorbidity index			0.65
0–2	33 (19.5)	89 (16.5)	
3–4	55 (32.5)	186 (34.4)	
> 4	81 (47.9)	265 (49.1)	

(continued to the next page)

Table 1. (Continued) Baseline characteristics of patients cared for by hospitalists group and non-hospitalists group

Characteristics	Hospitalist (n = 169)	Non-hospitalist (n = 540)	P value
Disease severity			
Mental change	13 (7.7)	37 (6.9)	0.71
Long-term care hospital	24 (14.2)	76 (14.1)	0.97
White blood cell, $\times 10^3/\text{mm}^3$	11.4 \pm 6.6	11.6 \pm 5.6	
Normal	77 (45.6)	212 (39.3)	0.15
Leucopenia ($< 4,000/\text{mm}^3$) or leukocytosis ($\geq 10,000/\text{mm}^3$)	92 (54.4)	328 (60.7)	
Bands neutrophil $> 5\%$	1 (0.6)	3 (0.6)	0.96
Tachypnea or hypoxemia	45 (26.6)	210 (38.9)	0.004
Septic shock	13 (7.7)	46 (8.5)	0.73
Platelet $< 150,000/\text{mm}^3$	27 (16)	103 (19.1)	0.36
MEDS score ^a			0.02
Very low	64 (37.9)	148 (27.4)	
Low	57 (33.7)	170 (31.5)	
Moderate	41 (24.3)	193 (35.7)	
Severe	4 (2.4)	24 (4.4)	
Very severe	3 (1.8)	5 (0.9)	
Nurse's working years, ^b median (IQR)	1.5 (0.8–4.5)	1.5 (0.6–4.4)	0.620
Outcomes			
Length of hospital stay	10.2 \pm 8.2	13.2 \pm 12.5	
Median (IQR)	8 (5–12)	10 (7–15)	< 0.001
In-hospital mortality	4 (2.4)	26 (4.8)	0.17
Readmission	22 (13.1)	84 (15.6)	0.43
Readmission within 14 days	4 (2.4)	4 (0.8)	0.15
Same disease	3 (1.8)	2 (0.4)	
Different disease	1 (0.6)	2 (0.4)	
Readmission within 30 days	9 (5.3)	15 (3.8)	0.17
Same disease	7 (4.1)	9 (1.7)	
Different disease	2 (1.2)	6 (1.1)	

Data are presented as mean \pm standard deviation or number (%).

HIV = human immunodeficiency virus, PAOD = peripheral arterial occlusive disease, COPD = chronic obstructive pulmonary disease, MEDS = Mortality in Emergency Department Sepsis score, IQR = interquartile range.

^aMEDS score: very low (0–4), low (5–7), moderate (8–12), severe (13–15), and very severe (> 15); ^bNurses' working years in the wards where hospitalists worked were compared with that the wards where non-hospitalists worked, including respiratory ward, emergency ward, and infectious disease ward.

and its components. Compared to the HG group, the NHG group had higher proportions of patients with tachypnea or hypoxemia (26.6% vs. 38.9%, respectively; $P = 0.004$). Based on the MEDS score, the proportions of patients with very low, low, or very severe MEDS score were higher in the HG group, while the proportions of moderately severe or severe MEDS score were higher in the NHG group. Compared with those in the NHG group, patients in the HG group had shorter LOS (2 days) in the hospital (median [IQR], 8 [5–12] vs. 10 [7–15] days, respectively; $P < 0.001$). The HG group showed lower IHM than the NHG group, but the difference was not significant (2.4% vs. 4.8%, respectively; $P = 0.170$).

Patients cared for by the non-hospitalists were mainly admitted in the respiratory ward (32.4%), emergency ward (15.0%), and infectious disease ward (11.5%). Nurses' working years was not significantly different between HG and NHG groups (median [IQR], 1.5 [0.8–4.5] vs. 1.5 [0.6–4.4], respectively; $P = 0.620$).

Subgroup analysis according to diseases (pneumonia or UTI), comorbidity, and disease severity

Of the 169 patients cared for by hospitalists, 70 (41.4%) were diagnosed with UTI, and 99 (58.7%) were admitted due to pneumonia (Table 2). A subgroup analysis stratified by disease (pneumonia or UTI), LOS, IHM, comorbidity (based on CCI), and disease severity (based on

Table 2. Subgroup analysis according to diseases (pneumonia or UTI) between hospitalist group and non-hospitalist group

Diagnosis	Pneumonia (n = 486)			Urinary tract infection (n = 223)		
	HG (n = 99)	NHG (n = 387)	P value (adjusted P value ^a)	HG (n = 70)	NHG (n = 153)	P value (adjusted P value ^a)
LOS, day	10.0 ± 7.6	13.7 ± 12.3		10.5 ± 9.1	11.7 ± 13.0	
Median (IQR)	8 (5–12)	10 (7–16)	< 0.001 (< 0.001)	7 (5–12)	8 (5–14)	0.49 (0.98)
IHM	7 (4–11)	9 (6–15)	0.62 (1.00)	0 (0)	4 (2.6)	0.17 (0.34)
Readmission						
Within 14 days	4/99 (4)	2/387 (0.5)	0.01 (0.02)	0 (0)	2 (1.3)	0.34 (0.68)
Within 30 days	6/99 (6.1)	13/387 (3.4)	0.26 (0.52)	3 (4.3)	2 (1.3)	0.35 (0.70)
CURB-65 score			0.308 (0.616)			-
0–1	58 (58.6)	205 (53.0)		-	-	
2	30 (30.3)	115 (29.7)		-	-	
≥ 3	11 (11.1)	67 (17.3)		-	-	
Charlson comorbidity index			0.84 (1.00)			0.84 (1.00)
0–2	17 (17.2)	54 (14)		16 (22.9)	35 (22.9)	
3–4	36 (36.4)	139 (35.9)		19 (27.1)	47 (30.7)	
≥ 5	46 (46.5)	194 (50.1)		35 (50)	71 (46.4)	
MEDS ^b			0.07 (0.14)			0.75 (1.00)
Very low (0–4)	21 (21.2)	63 (16.3)		43 (61.4)	85 (55.6)	
Low (5–7)	43 (43.4)	129 (33.3)		14 (20)	41 (26.8)	
Moderate (8–12)	30 (30.3)	170 (43.9)		11 (15.7)	23 (15.0)	
Severe and very severe (≥ 13)	5 (5.1)	25 (6.5)		2 (2.9)	4 (2.6)	

Data are presented as mean ± standard deviation or number (%).

UTI = urinary tract infection, HG = hospitalist group, NHG = non-hospitalist group, LOS = length of hospital stay, IQR = interquartile range, IHM = in-hospital mortality, CURB-65 = a system using Confusion, Urea, Respiratory rate, Blood pressure plus age ≥ 65 years for community-acquired pneumonia management, MEDS = Mortality in Emergency Department Sepsis score.

^aP value using Bonferroni adjustment for subgroup analysis; ^bMEDS score: very low (0–4), low (5–7), moderate (8–12), severe (13–15), and very severe (> 15).

MEDS score) were compared between the NHG and HG groups. In the pneumonia group, HG showed shorter median LOS than the NHG group (difference of 2 days between groups). However, the 14-day readmission rate for the HG group was 4.0%, and was significantly higher than that in the NHG group (0.5%, *P* = 0.020). Other variables did not show any significant difference.

Subgroup analysis stratified by CCI and MEDS score was performed to determine the difference between the two groups (Tables 3 and 4). In all CCI groups, the HG LOS was shorter than that for the NHG group. In the high CCI group with ≥ 5 points, HG LOS was significantly shorter than that of NHG group (median [IQR], 8 [5–12] vs. 10 [7–16] day; *P* < 0.001). IHMs of all CCI groups were not significantly different between the two groups. Patients in the HG group with moderate CCI (3–4) had higher readmission rates than those in the NHG group (14 and 30 days; *P* = 0.020 and *P* = 0.006, respectively).

Table 3. Subgroup analysis according to comorbidity between hospitalist group and non-hospitalist group

CCI	0–2 (n = 121)			3–4 (n = 241)			≥ 5 (n = 346)		
	HG (n = 33)	NHG (n = 88)	P value (adjusted P value ^a)	HG (n = 55)	NHG (n = 186)	P value (adjusted P value ^a)	HG (n = 81)	NHG (n = 265)	P value (adjusted P value ^a)
LOS, day	8.8 ± 5.3	11.5 ± 10.1		11.1 ± 8.9	12.9 ± 13.6		10.1 ± 8.7	14.0 ± 12.4	
Median (IQR)	7 (5–10)	8 (6–13)	0.25 (0.50)	8 (6–14)	9 (7–14)	0.25 (0.50)	8 (5–12)	10 (7–16)	< 0.001 (< 0.001)
IHM	0 (0)	1 (1.1)	0.54 (1.00)	0 (0)	9 (4.8)	0.09 (0.18)	4 (4.9)	16 (6)	0.71 (1.00)
Readmission									
Within 14 days	0 (0)	1 (1.1)	0.39 (0.78)	3 (5.4)	0 (0)	0.01 (0.02)	1 (1.2)	3 (1.2)	0.94 (1.00)
Within 30 days	0 (0)	2 (2.2)	0.75 (1.00)	6 (10.9)	3 (1.6)	0.003 (0.006)	3 (3.7)	10 (3.8)	0.98 (1.00)

Data are presented as mean ± standard deviation or number (%).

CCI = Charlson comorbidity index, HG = hospitalist group, NHG = non-hospitalist group, LOS = length of hospital stay, IQR = interquartile range, IHM = in-hospital mortality.

^aP value using Bonferroni adjustment for subgroup analysis.

Table 4. Subgroup analysis according to diseases severity between hospitalist group and non-hospitalist group

MEDS	Very low (0–4) (n = 212)			Low (5–7) (n = 227)			Moderate (8–12) (n = 234)			Severe and very severe (≥ 13) (n = 36)		
	HG (n = 64)	NHG (n = 148)	P value (adjusted P value)	HG (n = 57)	NHG (n = 170)	P value (adjusted P value)	HG (n = 41)	NHG (n = 193)	P value (adjusted P value)	HG (n = 7)	NHG (n = 29)	P value (adjusted P value)
LOS, day	8.8 ± 6.8	11.1 ± 10.8		9.5 ± 6.3	12.6 ± 10.9		11.7 ± 10.3	14.8 ± 13.0		20.3 ± 13.6	16.3 ± 22.0	
Median (IQR)	6 (5–10)	8 (6–12)	0.09 (0.18)	8 (6–10)	10 (7–15)	0.03 (0.05)	9 (5–14)	11 (8–18)	0.05 (0.09)	18 (12–32)	9 (5–17)	0.14 (0.28)
IHM	0 (0)	1 (0.7)	0.51 (1.00)	2 (3.5)	2 (1.2)	0.25 (0.50)	0 (0)	12 (6.2)	0.10 (0.20)	2 (28.6)	11 (37.9)	0.64 (1.00)
Readmission												
Within 14 days	0 (0)	3 (2.1)	0.25 (0.50)	1 (1.8)	0 (0)	0.08 (0.16)	3 (7.3)	1 (0.5)	0.002 (0.004)	0 (0)	0 (0)	1.00 (1.00)
Within 30 days	2 (3.2)	4 (2.8)	0.87 (1.00)	3 (5.3)	3 (1.8)	0.15 (0.30)	4 (9.7)	7 (3.7)	0.09 (0.18)	0 (0)	1 (3.4)	0.62 (1.00)

Data are presented as mean ± standard deviation or number (%).

MEDS = Mortality in Emergency Department Sepsis score, HG = hospitalist group, NHG = non-hospitalist group, LOS = length of hospital stay, IQR = interquartile range, IHM = in-hospital mortality.

As the MEDS score increased, the LOS in both groups increased (**Table 4**), although not statistically significant. All HG subgroups had lower LOS except the severe MEDS score group. The 14-day readmission rate for the HG group with moderate MEDS score was significantly higher than that for the NHG group (7.3% vs. 0.5%; $P = 0.004$).

Multiple regression analysis of factors associated with LOS

We determined the risk factors associated with LOS longer than 9 days using logistic regression analysis (data not shown). In univariable analysis, age over 65 years, men, cerebrovascular disease, dementia, altered mental status, lower respiratory tract infection, referred from long-term care hospital, tachypnea or hypoxemia, septic shock, band neutrophils over 5% of leukocyte, and hospitalist care were significantly associated with LOS ($P < 0.001$, $P = 0.003$, $P = 0.008$, $P = 0.03$, $P < 0.001$, $P = 0.001$, $P < 0.001$, $P = 0.03$, $P = 0.004$, $P = 0.014$, and $P < 0.001$, respectively). However, multiple logistic regression analysis revealed significant factors, including age over 65 years (odds ratio [OR], 1.68; 95% confidence interval [CI], 1.16–2.39; $P = 0.005$), lower respiratory tract infection (OR, 1.48; 95% CI, 1.01–2.16; $P = 0.046$), long-term care hospital (OR, 2.32; 95% CI, 1.43–3.74; $P = 0.001$), and hospitalist care (OR, 0.59; 95% CI, 0.40–0.87; $P = 0.008$).

Multiple linear regression was conducted to evaluate the association between CCI, MEDS score, gender, and hospitalist care and LOS (**Table 5**). CCI and MEDS score were positively associated with LOS ($P = 0.008$ and $P = 0.041$, respectively), while hospitalist care was negatively associated with LOS ($P = 0.003$). Hospitalist care was found to be associated with a 16.2% decrease in LOS. Considering these factors, it can be described using the following model:

$$\begin{aligned} \ln(\text{LOS}) &= 2.021 + 0.039 \times \text{CCI} + 0.017 \times \text{MEDS} - 0.177 \times \text{hospitalist} \\ \text{LOS} &= \exp(2.021 + 0.039 \times \text{CCI} + 0.017 \times \text{MEDS} - 0.177 \times \text{hospitalist}) \\ &= 7.7 \times (1.04)^{\text{CCI}} \times (1.017)^{\text{MEDS}} \times (0.838)^{\text{hospitalist}} \end{aligned}$$

Table 5. Multiple linear regression analysis of factors associated with the length of hospital stay (log-transformed)

Variables	Coefficients	Standard error	P value
CCI	0.039	0.015	0.008
MEDS	0.017	0.008	0.041
Gender (Men = 1, women = 0)	0.060	0.052	0.245
Hospitalist (Yes = 1, no = 0)	-0.177	0.060	0.003

CCI = Charlson comorbidity index, MEDS = Mortality in Emergency Department Sepsis score, IHM = in-hospital mortality.

Adjusted multiple $R^2 = 5.2\%$.

DISCUSSION

This is the first study to report the outcomes of a Korean independent hospitalist model responsible for the care of patients from admission to discharge. This study showed shorter LOS with hospitalist care than for the non-hospitalist care, and that the care of patients by hospitalists was associated with reduced overall LOS. The same trend was observed in the subgroup and multiple regression analyses. In particular, patients in the HG group with comorbidity had shorter LOS with CCI of ≥ 5 . HG group LOS was shown to be shorter than the national average LOS in patients with pneumonia or UTI with complications or comorbidities, (pneumonia, 8.27–17.27 days; UTI, 8.83–16.67 days), based on the health insurance review and assessment (HIRA) service data in 2017 and 2018.³¹ However, the readmission rates of patients in the HG group with moderate CCI and MEDS score were significantly increased. Despite the small sample size and lack of significant difference, the overall IHM in the HG group remained low compared to that of the NHG group, and the same trend was reported after the subgroup analysis.

Previous studies investigating the impact of hospitalists reported a reduction in LOS with hospitalist care.^{5,7,14,32-35} Similarly, hospitalist care had the strongest association with LOS in patients with pneumonia, UTI, or sepsis.³⁴ The reductions in LOS associated with hospitalist care was more prominent in older patients and those who developed complications.^{9,32,34} The implementation of hospitalist-operated system, like that utilized in our institute, was effective in patients visiting the ED.^{14,15,36} Unlike with pneumonia, no statistical difference in LOS was found with UTI between HG and NHG in the current study. This finding suggests that the HG group with UTI had higher proportions of patients referred from long-term care and that these patients could have had antibiotic-resistant strains that should have been treated with intravenous antibiotics for a prolonged period. In addition, the effect of individual diseases on LOS seems to be minimal as shown in the multiple logistic regression. This is because change in MEDS score and CCI was associated with LOS in multiple linear regression, ultimately it was thought that the patients' overall condition influenced the LOS.

Patients cared for by hospitalists could have had a shorter LOS due to several reasons. Because the hospitalists are better trained medical personnel than residents, it is possible that they are able to address the disease management from various perspectives and may be more likely to solve the patients problems.³⁷ Hospitalists also manage patients with similar diseases repeatedly; thus, they are more efficient due to their length of experience. Disease-specific physicians and hospitalists who are more experienced are effective in improving patient outcomes.^{4,38} To solve various patients' problems, from time to time, hospitalists communicated with the members of our department or with specialists. In our hospitalist system, emergency medical issues are monitored by the rapid response team, this is immediately addressed by our critical care team if the patient require intensive care treatment.

Although results were conflicting, higher readmission rates associated with hospitalist care were reported in previous studies.^{7,39} One possible explanation for the higher readmission rates reported here could be the disruption in the inpatient to outpatient continuity of care.^{7,12,40-43} In the current study, 3 of 4 patients who were re-admitted to the hospital within 14 days did not visit the outpatient clinic after discharge. In one of the patients that were readmitted within 30 days, the changes applied during the hospital stay were not properly communicated to the outpatient clinic specialist. The other reason was the post-discharge management of multi-comorbid and elderly patients; in the analysis of patients who were

readmitted within 30 days, 7 (77.8%) were patients with pneumonia, while 2 (22.2%) had UTI. Most patients that were readmitted with pneumonia were of old age and had neurological problems such as dementia, Parkinson's disease, or sequelae of cerebrovascular disease. Four of the patients were re-admitted with repeated aspiration.⁴⁴ To prevent this problem, short-term outpatient follow-up and careful management of discharged patients are needed.⁴⁵ Hospitalists should communicate adequately with specialists about referred patients and educate the patients regarding the importance of follow-up care.^{42,43} Hospitalists who provide post-discharge care may help patients in preventing re-hospitalization.⁴⁶⁻⁴⁸

Recently, the allocation of hospitalists to acute medical units in Korean tertiary hospitals resulted in a reduction in the LOS and ED waiting time.¹⁴ This study did not include other quality indicators, such as the morbidity, mortality, and readmission rate due to the short-term transit care. In our study, IHM, readmission rate, and LOS were obtained, and subgroup analysis using CCI and MEDS score was performed to determine the confounding factors that may affect LOS. We also analyzed the causes of readmissions. However, our study had some limitations. It was a retrospective study, used a relatively small sample size, and had a short-term study period. Secondly, patients who had been previously diagnosed with disease were included in the NHG group; therefore, a selection bias might have occurred during the assignment of patients in the ED, which may have affected the LOS. For adjustment of this bias, stratification analysis was performed in consideration of comorbidity and disease severity, and no statistical differences in leukocyte counts and previous long-term care hospitalization were found between HG and NHG groups. Thirdly, other quality indicators, such as discharge destination, and patient or staff satisfaction, were not evaluated in the present study. Finally, only two hospitalists worked during the day in our institute. Hence, further study is needed to evaluate the results of several hospitalists working full-time including those working for 24 hours a day, 7 days a week.

In conclusion, hospitalist care included in the integrated medical model can help reduce the LOS, especially in patients with multiple comorbidities. The interventions should be developed to prevent the readmission of patients having moderate comorbidity or severe disease. Further studies on various hospitalist-operated models and patients hospitalized with other diseases are warranted to evaluate the impact of hospitalist care in Korea.

REFERENCES

1. Wachter RM, Goldman L. Zero to 50,000 - The 20th anniversary of the hospitalist. *N Engl J Med* 2016;375(11):1009-11.
[PUBMED](#) | [CROSSREF](#)
2. Shu CC, Lin JW, Lin YF, Hsu NC, Ko WJ. Evaluating the performance of a hospitalist system in Taiwan: a pioneer study for nationwide health insurance in Asia. *J Hosp Med* 2011;6(7):378-82.
[PUBMED](#) | [CROSSREF](#)
3. Kisuule F, Howell E. Hospital medicine beyond the United States. *Int J Gen Med* 2018;11:65-71.
[PUBMED](#) | [CROSSREF](#)
4. Meltzer D, Manning WG, Morrison J, Shah MN, Jin L, Guth T, et al. Effects of physician experience on costs and outcomes on an academic general medicine service: results of a trial of hospitalists. *Ann Intern Med* 2002;137(11):866-74.
[PUBMED](#) | [CROSSREF](#)
5. Auerbach AD, Wachter RM, Katz P, Showstack J, Baron RB, Goldman L. Implementation of a voluntary hospitalist service at a community teaching hospital: improved clinical efficiency and patient outcomes. *Ann Intern Med* 2002;137(11):859-65.
[PUBMED](#) | [CROSSREF](#)

6. Bellet PS, Whitaker RC. Evaluation of a pediatric hospitalist service: impact on length of stay and hospital charges. *Pediatrics* 2000;105(3 Pt 1):478-84.
[PUBMED](#) | [CROSSREF](#)
7. Howrey BT, Kuo YF, Goodwin JS. Association of care by hospitalists on discharge destination and 30-day outcomes after acute ischemic stroke. *Med Care* 2011;49(8):701-7.
[PUBMED](#) | [CROSSREF](#)
8. Tadros RO, Faries PL, Malik R, Vouyouka AG, Ting W, Dunn A, et al. The effect of a hospitalist comanagement service on vascular surgery inpatients. *J Vasc Surg* 2015;61(6):1550-5.
[PUBMED](#) | [CROSSREF](#)
9. Landefeld CS. Care of hospitalized older patients: opportunities for hospital-based physicians. *J Hosp Med* 2006;1(1):42-7.
[PUBMED](#) | [CROSSREF](#)
10. Jang SI, Jang SY, Park EC. Trends of US hospitalist and suggestions for introduction of Korean hospitalist. *Korean J Med* 2015;89(1):1-5.
[CROSSREF](#)
11. Eom JS. Operating the hospitalist system. *J Korean Med Assoc* 2016;59(5):342-4.
[CROSSREF](#)
12. Lee DK. Launching the hospitalist in Korea. *Korean J Med* 2016;91(3):241-4.
[CROSSREF](#)
13. Kim HW. The current status of hospital medicine in Korea, 2019. *Korean J Med* 2019;94(2):139-44.
[CROSSREF](#)
14. Ohn JH, Kim NH, Kim ES, Baek SH, Lim Y, Hur J, et al. An Acute medical unit in a Korean tertiary care hospital reduces the length of stay and waiting time in the emergency department. *J Korean Med Sci* 2017;32(12):1917-20.
[PUBMED](#) | [CROSSREF](#)
15. Weng TC, Huang CC, Zhang CC, Huang HK, Wu MC, Wsu CC. Application of integrated medical care model (hospitalist) in emergency department and its efficiency and outcome: 4 years' experience in Taiwan. *Health Care Curr Rev* 2017;5(4):211-6.
[CROSSREF](#)
16. Matsui K, Goldman L, Johnson PA, Kuntz KM, Cook EF, Lee TH. Comorbidity as a correlate of length of stay for hospitalized patients with acute chest pain. *J Gen Intern Med* 1996;11(5):262-8.
[PUBMED](#) | [CROSSREF](#)
17. Rosenthal GE, Harper DL, Quinn LM, Cooper GS. Severity-adjusted mortality and length of stay in teaching and nonteaching hospitals. Results of a regional study. *JAMA* 1997;278(6):485-90.
[PUBMED](#) | [CROSSREF](#)
18. Ofori-Asenso R, Zomer E, Chin KL, Si S, Markey P, Tacey M, et al. Effect of comorbidity assessed by the Charlson comorbidity index on the length of stay, costs and mortality among older adults hospitalised for acute stroke. *Int J Environ Res Public Health* 2018;15(11):E2532.
[PUBMED](#) | [CROSSREF](#)
19. Hijazi HH, Alyahya MS, Hammouri HM, Alshraideh HA. Risk assessment of comorbidities on 30-day avoidable hospital readmissions among internal medicine patients. *J Eval Clin Pract* 2017;23(2):391-401.
[PUBMED](#) | [CROSSREF](#)
20. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40(5):373-83.
[PUBMED](#) | [CROSSREF](#)
21. Quan H, Li B, Couris CM, Fushimi K, Graham P, Hider P, et al. Updating and validating the Charlson comorbidity index and score for risk adjustment in hospital discharge abstracts using data from 6 countries. *Am J Epidemiol* 2011;173(6):676-82.
[PUBMED](#) | [CROSSREF](#)
22. Stavem K, Hoel H, Skjaker SA, Haagensen R. Charlson comorbidity index derived from chart review or administrative data: agreement and prediction of mortality in intensive care patients. *Clin Epidemiol* 2017;9:311-20.
[PUBMED](#) | [CROSSREF](#)
23. Yang Y, Yang KS, Hsann YM, Lim V, Ong BC. The effect of comorbidity and age on hospital mortality and length of stay in patients with sepsis. *J Crit Care* 2010;25(3):398-405.
[PUBMED](#) | [CROSSREF](#)
24. Shapiro NI, Howell MD, Talmor D, Donnino M, Ngo L, Bates DW. Mortality in Emergency Department Sepsis (MEDS) score predicts 1-year mortality. *Crit Care Med* 2007;35(1):192-8.
[PUBMED](#) | [CROSSREF](#)

25. Shapiro NI, Wolfe RE, Moore RB, Smith E, Burdick E, Bates DW. Mortality in Emergency Department Sepsis (MEDS) score: a prospectively derived and validated clinical prediction rule. *Crit Care Med* 2003;31(3):670-5.
[PUBMED](#) | [CROSSREF](#)
26. Williams JM, Greenslade JH, Chu K, Brown AF, Lipman J. Severity scores in emergency department patients with presumed infection: a prospective validation study. *Crit Care Med* 2016;44(3):539-47.
[PUBMED](#) | [CROSSREF](#)
27. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016;315(8):801-10.
[PUBMED](#) | [CROSSREF](#)
28. Lim WS, van der Eerden MM, Laing R, Boersma WG, Karalus N, Town GI, et al. Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation study. *Thorax* 2003;58(5):377-82.
[PUBMED](#) | [CROSSREF](#)
29. Hock Lee K, Yang Y, Soong Yang K, Chi Ong B, Seong Ng H. Bringing generalists into the hospital: outcomes of a family medicine hospitalist model in Singapore. *J Hosp Med* 2011;6(3):115-21.
[PUBMED](#) | [CROSSREF](#)
30. Feng C, Wang H, Lu N, Tu XM. Log transformation: application and interpretation in biomedical research. *Stat Med* 2013;32(2):230-9.
[PUBMED](#) | [CROSSREF](#)
31. Health Insurance Review & Assessment Service. *Hospital Average Length of Stay by Diagnostic Categories in 2017 and 2018. Indicator Status*. Wonju: Health Insurance Review & Assessment Service; 2019.
32. Rachoin JS, Skaf J, Cerceo E, Fitzpatrick E, Milcarek B, Kupersmith E, et al. The impact of hospitalists on length of stay and costs: systematic review and meta-analysis. *Am J Manag Care* 2012;18(1):e23-30.
[PUBMED](#)
33. Kuo YF, Goodwin JS. Effect of hospitalists on length of stay in the medicare population: variation according to hospital and patient characteristics. *J Am Geriatr Soc* 2010;58(9):1649-57.
[PUBMED](#) | [CROSSREF](#)
34. Southern WN, Berger MA, Bellin EY, Hailpern SM, Arnsten JH. Hospitalist care and length of stay in patients requiring complex discharge planning and close clinical monitoring. *Arch Intern Med* 2007;167(17):1869-74.
[PUBMED](#) | [CROSSREF](#)
35. Lindenaier PK, Rothberg MB, Pekow PS, Kenwood C, Benjamin EM, Auerbach AD. Outcomes of care by hospitalists, general internists, and family physicians. *N Engl J Med* 2007;357(25):2589-600.
[PUBMED](#) | [CROSSREF](#)
36. Howell EE, Bessman ES, Rubin HR. Hospitalists and an innovative emergency department admission process. *J Gen Intern Med* 2004;19(3):266-8.
[PUBMED](#) | [CROSSREF](#)
37. Freed DH. Hospitalists: Evolution, evidence, and eventualities. *Health Care Manag (Frederick)* 2004;23(3):238-56.
[PUBMED](#) | [CROSSREF](#)
38. Goodwin JS, Salameh H, Zhou J, Singh S, Kuo YF, Nattinger AB. Association of hospitalist years of experience with mortality in the hospitalized medicare population. *JAMA Intern Med* 2018;178(2):196-203.
[PUBMED](#) | [CROSSREF](#)
39. Kuo YF, Goodwin JS. Association of hospitalist care with medical utilization after discharge: evidence of cost shift from a cohort study. *Ann Intern Med* 2011;155(3):152-9.
[PUBMED](#) | [CROSSREF](#)
40. Hamel MB, Drazen JM, Epstein AM. The growth of hospitalists and the changing face of primary care. *N Engl J Med* 2009;360(11):1141-3.
[PUBMED](#) | [CROSSREF](#)
41. Sharma G, Fletcher KE, Zhang D, Kuo YF, Freeman JL, Goodwin JS. Continuity of outpatient and inpatient care by primary care physicians for hospitalized older adults. *JAMA* 2009;301(16):1671-80.
[PUBMED](#) | [CROSSREF](#)
42. Bell CM, Schnipper JL, Auerbach AD, Kaboli PJ, Wetterneck TB, Gonzales DV, et al. Association of communication between hospital-based physicians and primary care providers with patient outcomes. *J Gen Intern Med* 2009;24(3):381-6.
[PUBMED](#) | [CROSSREF](#)
43. Moore C, Wisnivesky J, Williams S, McGinn T. Medical errors related to discontinuity of care from an inpatient to an outpatient setting. *J Gen Intern Med* 2003;18(8):646-51.
[PUBMED](#) | [CROSSREF](#)

44. Cabré M, Serra-Prat M, Force L, Almirall J, Palomera E, Clavé P. Oropharyngeal dysphagia is a risk factor for readmission for pneumonia in the very elderly persons: observational prospective study. *J Gerontol A Biol Sci Med Sci* 2014;69(3):330-7.
[PUBMED](#) | [CROSSREF](#)
45. Tong L, Arnold T, Yang J, Tian X, Erdmann C, Esposito T. The association between outpatient follow-up visits and all-cause non-elective 30-day readmissions: a retrospective observational cohort study. *PLoS One* 2018;13(7):e0200691.
[PUBMED](#) | [CROSSREF](#)
46. Burke RE, Ryan P. Postdischarge clinics: hospitalist attitudes and experiences. *J Hosp Med* 2013;8(10):578-81.
[PUBMED](#) | [CROSSREF](#)
47. Mahant S, Weinstein M. Hospitalists and the outpatient clinic: time to reconsider. *J Pediatr* 2016;168:3-4.
[PUBMED](#) | [CROSSREF](#)
48. Shu CC, Hsu NC, Lin YF, Wang JY, Lin JW, Ko WJ. Integrated postdischarge transitional care in a hospitalist system to improve discharge outcome: an experimental study. *BMC Med* 2011;9(1):96.
[PUBMED](#) | [CROSSREF](#)