

Genetic prion disease: D178N with 129MV disease modifying polymorphism – a clinical phenotype

Tracie Huey-Lin Tan ¹, Richard J Stark,^{1,2} John A Waterston,¹ Owen White,² Dominic Thyagarajan,² Mastura Monif^{1,2}

To cite: Tan TH-L, Stark RJ, Waterston JA, *et al.* Genetic prion disease: D178N with 129MV disease modifying polymorphism—a clinical phenotype. *BMJ Neurology Open* 2020;**2**:e000074. doi:10.1136/bmjno-2020-000074

Accepted 17 August 2020

ABSTRACT

Background Human prion diseases are a group of rare neurological diseases with a minority due to genetic mutations in the prion protein (PRNP) gene. The D178N mutation is associated with both Creutzfeldt-Jakob disease and fatal familial insomnia with the phenotype modified by a polymorphism at codon 129 with the methionine/valine (MV) polymorphism associated with atypical presentations leading to diagnostic difficulty.

Case We present a case of fatal familial insomnia secondary to a PRNP D178N mutation with 129MV disease modifying polymorphism who had no family history, normal MRI, electroencephalography (EEG), cerebrospinal fluid (CSF) and positron emission tomography findings and a negative real-time quaking-induced conversion result.

Conclusion Patients with genetic prion disease may have no known family history and normal EEG, MRI brain and CSF findings. PRNP gene testing should be considered for patients with subacute progressive neurological and autonomic dysfunction.

CASE REPORT

A Caucasian, left-handed, office worker in her early 40s, presented with a 12-month history of progressive symptoms which began with intermittent diplopia and light-headedness with an initially normal neurological examination. Over the following months, she developed cerebellar and brainstem dysfunction with macrosaccadic oscillations, saccadic smooth pursuit and bilaterally inaccurate saccades with significant leftwards hypermetric saccades ([figure 1](#)). Worsening dysarthria, limb and truncal ataxia resulted in numerous falls and communication issues forcing her to leave her job. Focal seizures began 4 months post disease onset with speech arrest preceded by an autoscopic phenomenon where she felt ‘to the right’ of herself. Focal to bilateral tonic-clonic seizures occurred 7 months post disease onset with post ictal left hemiparesis and expressive dysphasia.

She was admitted 8 months into the disease course at which point strength and sensory

testing was unremarkable. However, there were upper motor neuron signs of upgoing plantar reflexes, generalised hyperreflexia and lower limb spasticity in addition to severe cerebellar findings resulting in an inability to mobilise independently. While she was alert and able to hold an appropriate conversation, an Addenbrooke’s cognitive assessment revealed diffuse cortical dysfunction with a total score of 73/100 9 months post disease onset. In particular, there was executive dysfunction with deficits in letter more than category fluency and poor planning with visuospatial tasks. Memory was affected with impaired delayed recall, but there was relative sparing of language domains. Autonomic dysfunction with persistent borderline tachycardia of 90–100 bpm and a postural systolic blood pressure drop of 50–60 mm Hg was noted throughout her inpatient stay. Furthermore, she had a 6 kg loss of weight over 6 months despite a normal oral intake and no symptoms of malabsorption.

Her medical history was significant for long-standing mild insomnia, not worse in the recent months, with a normal sleep-wake cycle in hospital although formal sleep studies were not performed. She also had a history of depression and anxiety since her teenage years. Family history was incomplete as the patient’s father was adopted and she had lost contact with her maternal extended family. However, her parents, and her two older brothers and one older maternal half-brother, along with her four nieces and nephews, had no neurological issues.

Given the presentation with subacute ataxia, dysautonomia, cognitive dysfunction and seizures, autoimmune, atypical infectious and rapidly progressive neurodegenerative conditions were considered. MRI brain, positron emission tomography (PET) brain and cerebrospinal fluid (CSF) studies, including 14-3-3 protein, tau and real-time



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Neurology, Alfred Health, Melbourne, Victoria, Australia
²Neuroscience, Monash University Faculty of Medicine Nursing and Health Sciences, Clayton, Victoria, Australia

Correspondence to

Dr Tracie Huey-Lin Tan;
t.tan@alfred.org.au




Figure 1 (A) Saccadic testing showing bilateral gaze evoked nystagmus. (B) Pursuit testing showing left sided ocular dysmetria. (C) Testing of saccadic accuracy (to 10° targets), right and left saccades overlaid showing normal rightward saccades. The leftward saccades are hypermetric, by comparison, consistent with the observed overshoot. (D) A routine interictal electroencephalography showing a normal background. The electro-oculography leads (top trace) shows direction changing nystagmus (circled).

quaking-induced conversion (RT-QuIC) were unremarkable as was testing for anti-neuronal, glutamate decarboxylase, tissue transglutaminase and anti-gliadin antibodies. Interictal electroencephalography (EEG) was normal and post ictal EEG showed non-specific frontal intermittent rhythmic delta activity. A small bowel biopsy did not have evidence of Whipple's disease and syphilis antibody testing was negative. Genetic testing for spinocerebellar ataxias 1, 2, 3, 6, 7 and 15, Friedreich's Ataxia and Huntington's disease was unremarkable. 24 hours urinary copper and organic and amino acid screens for inherited metabolic disorders was also negative.

Given the clinical features and rapidly progressive course, a prion protein (PRNP) genetic analysis was performed and revealed a heterogenic pathogenic variant, D178N (aspartic acid to asparagine substitution), in the PRNP gene combined with a heterozygous MV (methionine/valine) disease modifying polymorphism at codon 129 which is associated with Creutzfeldt-Jakob disease (CJD) and fatal familial insomnia (FFI).

DISCUSSION

Prions are pathogenic misfolded proteins which replicate by causing conformational change and misfolding in neighbouring proteins. This leads to an exponential increase in prion formation with consequent neuronal damage.¹ Since their discovery, several rapidly progressive human neurological diseases have been shown to be due to prions. These human prion diseases include CJD, Kuru and various genetic prion diseases.¹ In more recent times, the pathogenesis of other neurodegenerative diseases including the α -synucleinopathies and tauopathies, are also hypothesised to involve prion like spread.² In addition,

multiple systems atrophy, which can present with rapidly progressive cerebellar and autonomic dysfunction akin to the classic human prion diseases, has been shown to be transmissible following animal inoculation with brain homogenate from deceased patients.³ While the implications of prions in human disease is broad, this discussion will focus on the classic human prion diseases.

Human prion diseases are rare, with an annual incidence of around 1–1.5 per million people.^{4 5} Genetic prion diseases account for around 8%–15% of human prion disease^{1 4 5} and are classified into three clinicopathological subtypes: genetic CJD, FFI and Gerstmann-Straussler-Scheinker syndrome (GSS syndrome). There are over 30 PRNP gene mutations¹ that have been associated with human genetic prion diseases with significant genotype-phenotype variability and differing age dependent penetrance.

The D178N mutation on the PRNP gene is associated with the CJD or FFI phenotypes. The clinical presentation can be modified by a polymorphism in codon 129 of the PRNP gene—methionine homozygosity (129MM) at this position is associated with an FFI phenotype and valine homozygosity (129VV) is associated with a CJD phenotype, although significant clinical phenotypic overlap exists.^{6 7}

CJD characteristically presents with a rapidly progressive cognitive impairment, cerebellar dysfunction, behavioural or psychiatric disturbance and visual changes with later development of extrapyramidal and pyramidal symptoms and myoclonus.¹ Compared with sporadic CJD, familial cases tend to present earlier with slower progression.⁸ FFI typically presents with sleep disturbance, dysautonomia and visual deficits with subsequent development of extrapyramidal signs, hallucinations, disorientation and cerebellar signs and later myoclonus and pyramidal signs.⁹

Our patient presented with an atypical clinical syndrome with prominent early ataxia and less pronounced cognitive deficit and sleep disturbance in association with heterozygosity at codon 129. A D178N mutation with 129 heterozygosity (129MV) is less commonly reported but can also present clinically as CJD or FFI, typically with a longer duration of disease than the codon 129 homozygotes.^{6 10 11} In terms of the clinical phenotype, a study by Krasnianski *et al* suggested that 129MV patients with FFI present with visual changes and ataxia earlier than their 129MM counterparts (6 vs 13 weeks and 9 vs 21 weeks, respectively) and tend to have a later onset of sleep disturbance (15 vs 3 weeks), hallucinations (57.5 vs 16 weeks), spatial disorientation (38.5 vs 20 weeks) and myoclonus (32 vs 16 weeks).¹⁰ Montagna *et al* found similar phenotypic differences but also noted that the 129MV patients were more likely to suffer from tonic-clonic seizures.¹¹ This phenotype of FFI would align with our case's presentation with early ataxia and visual disturbance with relatively preserved cognition and sleep wake cycle.

Despite being an autosomal dominant condition, our case did not have a known family history. While a family

history may be identified in 76%–92% of patients with a D178N mutation,^{7 12} not all patients will have a positive family history likely due to a combination of misdiagnosis of family members, variable age-dependent penetrance or a sporadic mutation in the index case.

While FFI patients with the D178N mutation frequently do not have a positive CSF 14-3-3 protein^{7 9} and only have non-specific changes on EEG and MRI brain,^{7 9} they often display thalamic and/or cortical hypometabolism, even pre-symptomatically, on [¹⁸F] fluorodeoxyglucose PET scan^{11 13} which our patient did not.

RT-QuIC is a relatively new technique that detects PrP^{Sc} (pathological scrapie isoform of the prion protein) in tissue and CSF¹⁴ and has a reported specificity of 98% and a sensitivity that varies with genetic mutation.¹⁵ In contrast, CSF 14-3-3 protein and tau have much lower specificities of 63% and 46%, respectively.¹⁵ In FFI cases, only 7%–8.3% of patients demonstrate CSF 14-3-3 protein or tau positivity.^{9 14} In comparison, RT-QuIC positivity is significantly higher but varies widely between studies from 17% in a Chinese study to 57% in a German study and 83% in a Japanese study.^{14 16 17} The significant discrepancy in sensitivities between studies may reflect different populations studied with different disease modifiers or different testing protocols. While it is clear that the PRNP 129 codon affects the clinical expression of the D178N mutation, it is not certain whether it has an effect on the RT-QuIC positivity and most studies only contained 129MM homozygotes or very few 129MV heterozygotes.^{14 16 17} While RT-QuIC is a promising new technique in the diagnosis of prion diseases, it may not be as useful in FFI cases with a PRNP D178N mutation and 129MV polymorphism as highlighted by our case.

This case is an uncommon presentation of a rare genetic prion disease. The patient presented without a known family history and with normal CSF, MRI and PET brain scans and interictal EEG. Promising new techniques such as RT-QuIC, which is highly sensitive in gCJD and GSS syndrome^{14 16} may not be as reliable in the FFI cohort and perhaps especially so in patients with the less common codon 129MV disease modifier mutation, although further studies are required.

The variability in presentation and the insensitivity of investigations in detecting genetic prion diseases, as illustrated by this case, emphasises the importance of maintaining a high level of suspicion for these conditions. PRNP genetic testing is crucial to obtaining the correct diagnosis in this situation which, in turn, allows for the appropriate counselling and management of patients and their families.

Acknowledgements The authors thank Dr Christiane Stehmann, PhD, Coordinator, Australian National CJD Registry, The Florey Institute, Parkville, Australia for assistance with genetic testing and advice.

Contributors TH-LT collected and analysed the data and wrote the manuscript. RJS, JAW, DT, OW and MM provided intellectual review, feedback and data analysis.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data sharing not applicable as no datasets generated and/or analysed for this study. Data sharing not applicable as no datasets generated and/or analysed for this study.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Tracie Huey-Lin Tan <http://orcid.org/0000-0002-1638-7129>

REFERENCES

- 1 Takada LT, Geschwind MD. Prion diseases. *Semin Neurol* 2013;33:348–56.
- 2 Tarutani A, Hasegawa M. Prion-like propagation of α -synuclein in neurodegenerative diseases. *Prog Mol Biol Transl Sci* 2019;168:323–48.
- 3 Woerman AL, Kazmi SA, Patel S, *et al*. MSA prions exhibit remarkable stability and resistance to inactivation. *Acta Neuropathol* 2018;135:49–63.
- 4 Stehmann C, Senesi M, Lewis V, Umami M, *et al*. Creutzfeldt-Jakob disease surveillance in Australia: update to 31 December 2018. *Commun Dis Intell* 2019;43:1–18.
- 5 Ladogana A, Puopolo M, Croes EA, *et al*. Mortality from Creutzfeldt-Jakob disease and related disorders in Europe, Australia, and Canada. *Neurology* 2005;64:1586–91.
- 6 Goldfarb LG, Petersen RB, Tabaton M, *et al*. Fatal familial insomnia and familial Creutzfeldt-Jakob disease: disease phenotype determined by a DNA polymorphism. *Science* 1992;258:806–8.
- 7 Zarranz JJ, Dignon A, Atarés B, *et al*. Phenotypic variability in familial prion diseases due to the D178N mutation. *J Neurol Neurosurg Psychiatry* 2005;76:1491–6.
- 8 Bagyinszky E, Giau VV, Youn YC, *et al*. Characterization of mutations in PRNP (prion) gene and their possible roles in neurodegenerative diseases. *Neuropsychiatr Dis Treat* 2018;14:2067–85.
- 9 Krasnianski A, Bartl M, Sanchez Juan P, *et al*. Fatal familial insomnia: clinical features and early identification. *Ann Neurol* 2008;63:658–61.
- 10 Krasnianski A, Sanchez Juan P, Ponto C, *et al*. A proposal of new diagnostic pathway for fatal familial insomnia. *J Neurol Neurosurg Psychiatry* 2014;85:654–9.
- 11 Montagna P, Cortelli P, Avoni P, *et al*. Clinical features of fatal familial insomnia: phenotypic variability in relation to a polymorphism at codon 129 of the prion protein gene. *Brain Pathol* 1998;8:515–20.
- 12 Kovács GG, Puopolo M, Ladogana A, *et al*. Genetic prion disease: the EUROCJD experience. *Hum Genet* 2005;118:166–74.
- 13 Cortelli P, Perani D, Montagna P, *et al*. Pre-symptomatic diagnosis in fatal familial insomnia: serial neurophysiological and 18FDG-PET studies. *Brain* 2006;129:668–75.
- 14 Sano K, Satoh K, Atarashi R, *et al*. Early detection of abnormal prion protein in genetic human prion diseases now possible using real-time QuIC assay. *PLoS One* 2013;8:e54915.
- 15 Foutz A, Appleby BS, Hamlin C, *et al*. Diagnostic and prognostic value of human prion detection in cerebrospinal fluid. *Ann Neurol* 2017;81:79–92.
- 16 Cramm M, Schmitz M, Karch A, *et al*. Stability and reproducibility underscore utility of RT-QuIC for diagnosis of Creutzfeldt-Jakob disease. *Mol Neurobiol* 2016;53:1896–904.
- 17 Xiao K, Shi Q, Zhou W, *et al*. T188K-Familial Creutzfeldt-Jacob disease, predominant among Chinese, has a reactive pattern in CSF RT-QuIC different from D178N-Fatal familial insomnia and E200K-Familial CJD. *Neurosci Bull* 2019;35:519–21.