

Choice of a suitable diabetes risk assessment tool in Nepal – can we learn from Canada?

Gina Agarwal¹, Brijesh Sathian², Sutapa Agrawal³

Correspondence: Dr. Gina Agarwal, Associate Professor, Departments of Family Medicine, Clinical Epidemiology and Biostatistics, Quality Assurance Program Coordinator for Family Medicine Residency, Residency Program Research Coordinator, Family Medicine Residency Program, McMaster University, David Braley Health Sciences Centre, 100 Main Street West, 5th Floor, Hamilton, ON L8P 1H6, Canada
Email: gina.agarwal@gmail.com

Received 24 June 2016/**Revised** 25 June 2016/**Accepted** 26 June 2016

Citation: Agarwal G, Sathian B, Agarwal S. Choice of a suitable diabetes risk assessment tool in Nepal – can we learn from Canada?. Nepal J Epidemiol. 2016;6(2); 548-550.

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Copyright © 2016 CEA& INEA. Published online by NepJOL-INASP.
www.nepjol.info/index.php/NJE

Diabetes is extremely prevalent globally and increasing in prevalence in low and middle income countries (LMICs) [1, 2], such as Nepal. Indeed, the pooled prevalence of type 2 diabetes in Nepal from a systematic review examining data from 2000 to 2014 was found to be 8.4% (95% CI: 6.2–10.5%) [3] – higher than the current national estimate of 4.5% [4]. The South Asian population (comprising of people from India, Pakistan, Bangladesh, Sri Lanka, Nepal, Bhutan and Tibet) [5], is at high risk of developing type 2 diabetes (T2DM) [6, 7].

The culprits causing this rapid surge in diabetes are many. It is due to a variety of factors including a strong genetic predisposition, a preponderance to be sedentary and certain dietary factors [6, 8, 9,10]. Diet and lifestyle are rapidly changing in LMICs in general, mimicking western lifestyle and diet [11, 12].

Biological theory states that adiposity-induced insulin resistance followed by a subsequent decline in pancreatic-cell function, causes T2DM [13]. Indeed, although South Asians have conventionally “normal” BMI ranges, they have a higher percentage of body fat, increased visceral abdominal fat and greater insulin resistance compared with white people [14,15]. Genetic studies suggest that South Asians carry many of the

specific genetic variants that are required for T2DM to develop, though the mechanism of activation, whether by environmental or physiological factors, is not clear [16]. Intra-uterine stressors leading to low birthweight in infants may cause epigenetic changes in gene expression and a pre-disposition to visceral adiposity, leading to T2DM [17]. South Asians are physically less active. A systematic review of studies in UK South Asians reported physical activity levels that were 50–75% lower than those of Europeans [18].

The South Asian diet itself is not only higher in overall calories and percentage of carbohydrate content, compared with standard European meals [19,20], but there has been a shift from eating traditional to non traditional foods which are also higher in animal proteins, sugar, fats and trans-fats (20). These trends exist not only in South Asians outside of South Asia, but in those residing in their home countries too – and Nepal is no exception [21]. All this to say that diabetes is in Nepal to stay.

Of course, diabetes is an extremely costly condition [1] and the resource implications of a diabetes epidemic in any LMIC such as Nepal are very serious [11, 12]. One way of curbing such an epidemic would be to diagnose people with diabetes and pre diabetes earlier, thus reducing or even preventing

complications of diabetes [22], by helping them gain access to appropriate help. However, the method of identification for those at risk for diabetes and hence needing screening, is not clear amongst LMICs in general, particularly Nepal.

In Canada for example, the use of the CANRISK diabetes risk assessment tool (23) is now more widespread, and in other countries, ethnic specific diabetes risk tools have been developed [24-28]. However, there is no consensus as to which tool would be appropriate for use in Nepal. Though the existing tools have been developed in specific populations of certain ethnicities, but they may be inappropriate for use in other ethnic groups.

The Indian Diabetes Risk Score (IDRS) was developed for a population living in India, though may be suitable for a Nepalese population as well [29]. It has 4 question categories (age, waist circumference, family history and physical activity), while the CANRISK has at least 12 questions (Gender, Parents' ethnicities, Waist circumference, BMI, Physical Activity, Fruit and Vegetable intake, High Blood Pressure, High Sugar, and Family History). A shorter and less intense screening tool may have advantages for use in the practicalities of administration in a busy rural situation or urban office. The IDRS was found to have an area under the receiver operating curve (AUC) of 0.698 (95%confidence interval (CI):0.663 –0.733); and sensitivity of 72.5% and specificity of 60.1% for determining undiagnosed diabetes with a positive predictive value of 17.0%, negative predictive value of 95.1%, and accuracy of 61.3% [28]. This scoring tool has not been tested in a Nepalese population. There is a need for studies to find suitable tools or for the development of a Nepal specific diabetes risk assessment tool ('NEPAL-RISK') to guide a targeted 2 tier approach for screening, in which individuals are assessed for risk en-masse, but only formally screened according to the risk results, thus saving resources.

If the population can be made more aware about diabetes by the use of a risk assessment tool as an educational tool as well, it could help to curb the diabetes epidemic in Nepal. Education of the masses about diabetes risk factors, prevention, and complications is urgently needed, using clear and simple messages. National policy efforts can be strengthened and health outcomes improved when awareness is increased [30]. Perhaps learning from Canada is a start, and Nepal will be able to make progress with something simple like 'NEPAL-RISK'?

Conflict of interest:

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding:

The authors received no financial support for the research, authorship, and/or publication of this article.

Author's affiliations:

¹ Associate Professor, Departments of Family Medicine, Clinical Epidemiology and Biostatistics, Quality Assurance Program Coordinator for Family Medicine Residency,

Residency Program Research Coordinator, Family Medicine Residency Program, McMaster University, David Braley Health Sciences Centre, 100 Main Street West, 5th Floor, Hamilton, ON L8P 1H6, Canada

² Assistant Professor, Community Medicine Department, Manipal College of Medical Sciences, Pokhara, Nepal

³ Epidemiologist, South Asia Network for Chronic Disease, Public Health Foundation of India, Fourth Floor, Plot no 47, Sector 44, Gurgaon (Haryana)-122002, India

References:

1. IDF Atlas: 6th Edition. Diabetes in low-middle-and high-income countries; Accessed May 11, 2015: Available at: <http://www.idf.org/diabetesatlas>
2. Estimates for the year 2000 and projections for 2030. World Health 2004; 27:1047–1053.
3. Gyawali B, Sharma R, Neupane D, Mishra SR, van Teijlingen E, Kallestrup P. Prevalence of type 2 diabetes in Nepal: a systematic review and meta-analysis from 2000 to 2014. Glob Health Action. 2015 Nov 26;8:29088. doi: 10.3402/gha.v8.29088. eCollection 2015. PMID: 26613684
4. International Diabetes Federation. Country estimates table 2011. 2012. [cited 6 May 2015]. Available from: http://www.idf.org/sites/default/files/EN_6E_Atlas_Full_0.pdf
5. Demography Division. Projections of the diversity of the Canadian population, 2006 to 2031. Statistics Canada (2010).Catalogue number 91-551-X. Accessed on June 29th 2016. Available from: <http://www.statcan.gc.ca/pub/91-551-x/91-551-x2010001-eng.pdf>
6. Lee JW, Brancati FL, Yeh HC. Trends in the prevalence of type 2 diabetes in Asians versus whites: results from the United States National Health Interview Survey, 1997-2008. Diabetes Care. 2011 Feb;34(2):353-7. <http://dx.doi.org/10.2337/dc10-0746>
7. Sohal P. Prevention and Management of Diabetes in South Asians. Canadian Journal of Diabetes. 2008; 32(3): 206-210. [http://dx.doi.org/10.1016/S1499-2671\(08\)23011-X](http://dx.doi.org/10.1016/S1499-2671(08)23011-X)
8. Chan JC, Malik V, Jia W, Kadowaki T, Yajnik CS, Yoon KH, Hu FB. Diabetes in Asia: epidemiology, risk factors, and pathophysiology. JAMA. 2009 May 27;301(20):2129-40. <http://dx.doi.org/10.1001/jama.2009.726> PMID:19470990
9. Patel KC, Bhopal R. Diabetes epidemic in the South Asian Diaspora: action before desperation. J R Soc Med. 2007 Mar;100(3):115-6. <http://dx.doi.org/10.1258/jrsm.100.3.115> PMID:17339300 PMCID:PMC1809161

10. Ramachandran A, Ma RC, Snehalatha C. Diabetes in Asia. *Lancet*. 2010 Jan 30;375(9712):408-18. [http://dx.doi.org/10.1016/S0140-6736\(09\)60937-5](http://dx.doi.org/10.1016/S0140-6736(09)60937-5)
11. Popkin BM. The nutrition transition and obesity in the developing world. *J Nutr*. 2001 Mar;131(3):871S-873S. PMID:11238777
12. Bovet P, Paccaud F. Cardiovascular disease and the changing face of global public health: a focus on low and middle income countries. *Public Health Reviews*. 2012;33(2):397-415.
13. Kahn SE. The relative contributions of insulin resistance and beta-cell dysfunction to the pathophysiology of Type 2 diabetes. *Diabetologia*. 2003; 46(1): 3–19
PMid:12637977
14. Rana A, de Souza RJ, Kandasamy S, Lear SA, Anand SS. Cardiovascular risk among South Asians living in Canada: a systematic review and meta-analysis. *CMAJ Open*. 2014 Jul 22;2(3):E183-91. doi: 10.9778/cmajo.20130064. eCollection 2014.
<http://dx.doi.org/10.9778/cmajo.20130064>
PMid: 25295238
15. Raji A, Seely EW, Arky RA, Simonson DC. Body fat distribution and insulin resistance in healthy Asian Indians and Caucasians. *J Clin Endocrinol Metab*. 2001 Nov;86(11):5366-71.
<http://dx.doi.org/10.1210/jcem.86.11.7992>
PMid:11701707
16. Gujral UP, Pradeepa R, Weber MB, Narayan KV, Mohan V. Type 2 diabetes in South Asians: similarities and differences with white Caucasian and other populations. *Annals of the New York Academy of Sciences*. 2013;1281(1):51-63.
<http://dx.doi.org/10.1111/j.1749-6632.2012.06838.x>
17. Chan JC, Malik V, Jia W, Kadowaki T, Yajnik CS, Yoon KH, Hu FB. Diabetes in Asia: epidemiology, risk factors, and pathophysiology. *JAMA*. 2009 May 27;301(20):2129-40. doi: 10.1001/jama.2009.726.
<http://dx.doi.org/10.1001/jama.2009.726>
PMid:19470990
18. Fischbacher CM, Hunt S, Alexander L. How physically active are South Asians in the United Kingdom? A literature review. *J. Public Health (Oxf.)*. 2004;26: 250–258.
<http://dx.doi.org/10.1093/pubmed/fdh158>
PMid:15454592
19. Burden ML, Samanta A, Spalding D, Burden AC. A comparison of the glycemic and insulinaemic effects of an Asian and a European meal. *Pract. Diab. Int*. 1994;11: 208–211.
<http://dx.doi.org/10.1002/pdi.1960110508>
20. Raj S, Ganganna P, Bowering J. Dietary habits of Asian Indians in relation to length of residence in the United States. *J Am Diet Assoc*. 1999 Sep;99(9):1106-8.
[http://dx.doi.org/10.1016/S0002-8223\(99\)00266-7](http://dx.doi.org/10.1016/S0002-8223(99)00266-7)
21. Subedi YP, Marais D, Newlands D. Where is Nepal in the nutrition transition? *Asia Pac J Clin Nutr*. 2015;published online November 2015. doi:10.6133/apjcn.112015.10.
22. Lindström J, Tuomilehto J. The Diabetes Risk Score: a practical tool to predict type 2 diabetes risk. *Diabetes Care* 2003;26:725–731.
<http://dx.doi.org/10.2337/diacare.26.3.725>
PMid:12610029
23. Agarwal G, Kaczorowski J, Hanna S. Effect of a Community-Based Diabetes Awareness Program on the Detection of Diabetes. 2014, 1:1–8.
24. Robinson CA, Agarwal G, Nerenberg K. Validating the CANRISK prognostic model for assessing diabetes risk in Canada 's multi-ethnic population. 2011; 32:19–31.
25. Taksande B, Ambade M, Joshi R. External validation of Indian diabetes risk score in a rural community of central India. *J Diabetes Mellit*. 2012; 02:109–113.
<http://dx.doi.org/10.4236/jdm.2012.21017>
26. Bumrerraj S, Kaczorowski J, Kessomboon P, Thinkhamrop B, Rattarasarn C: Simple Risk Score to Identify Population at Risk of Impaired Glucose Tolerance in the Thai Population. *Austin J Endocrinol Diabetes*. 2014; 1(4): 4.
27. Pramono L a, Setiati S, Soewondo P, Subekti I, Adisasmita A, Kodim N, Sutrisna B. Prevalence and predictors of undiagnosed diabetes mellitus in Indonesia. *Acta Med Indones* 2010, 42:216–223.
PMid:21063043
28. Neatrour G, Aloï J, Graves A. Using a Risk Assessment Questionnaire to Identify Prediabetics and Diabetics in Tandag, Philippines. (In Press)
29. Mohan V, Deepa R, Deepa M, Somannavar S, Datta M. A simplified Indian Diabetes Risk Score for screening for undiagnosed diabetic subjects. *J Assoc Physicians India*. 2005 Sep; 53:759-63.
PMid:16334618
30. Yadav NK, Sathian B, Kalai RS. Assessment of Diabetes Mellitus in India and Nepal. *Webmed Central BIOCHEMISTRY* 2012;3(6):WMC00354