

RESEARCH ARTICLE

Open Access

Lavender oil suppresses indoleamine 2,3-dioxygenase activity in human PBMC

Johanna M Gostner¹, Markus Ganzera², Kathrin Becker³, Simon Geisler³, Sebastian Schroecksnadel³, Florian Überall¹, Harald Schennach⁴ and Dietmar Fuchs^{3*}

Abstract

Background: Lavender remedies have been used in traditional medicine because of antimicrobial, anti-inflammatory and mood alleviating effects, but underlying molecular mechanisms are not yet fully elucidated. Recently, studies investigating the effects of lavender oil in the context of psychiatric disorders have indicated potent pharmacological properties. Metabolism of tryptophan by indoleamine 2,3-dioxygenase (IDO) was found to provide a biochemical link between immunology and neuroendocrinology and to be a frequent target of natural products.

Methods: In this *in vitro* study, interferences of lavender oil and constituents (–)-linalool, (+)- α -pinene and (+)-limonene with tryptophan catabolism by IDO and formation of neopterin via guanosine triphosphate (GTP)-cyclohydrolase-I and of interferon- γ have been investigated using unstimulated and phytohemagglutinin (PHA)-stimulated human peripheral blood mononuclear cells (PBMC).

Results: Treatment with lavender oil dose-dependently suppressed PHA-induced tryptophan breakdown and kynurenine formation. Similar effects were observed for the three constituents. In parallel, formation of neopterin and interferon- γ was diminished upon lavender oil treatment. In unstimulated PBMC, effect of lavender oil treatment was similar, but less pronounced.

Conclusion: Data from this *in vitro* study suggest that lavender oil treatment might contribute to the modulation of the immune and neuroendocrine system by interfering with activation-induced tryptophan breakdown and IDO activity.

Keywords: Lavender oil, Tryptophan, Indoleamine 2,3-dioxygenase, Neopterin, Kynurenine

Background

Lavender (*Lavandula angustifolia*) essential oil consists of a mixture of mono- and sesquiterpenoid alcohols, esters, oxides and ketones, containing linalool, linalyl acetate, 1,8-cineole, terpinen-4-ol, β -caryophyllene and camphor as primary components [1,2].

Lavender and its essential oil have been used since centuries due to antiseptic, antimicrobial and sedative effects. In today's folk and complementary medicine, the oil is applied also for the treatment of conditions such as anxiety, restlessness, insomnia and depression. Administration routes include absorption via the respiratory tract (aromatherapy) or oral ingestion [3]. Although there is

evidence-based information on the pharmaceutical efficacy of lavender oil for the treatment of anxiety-related disturbances [4], its therapeutic significance was little appreciated for a long time, due to the lack of larger clinical trials, but also due to methodological problems in constituent identification and standardization of such complex multicomponent preparations.

Recently, Kasper *et al.* demonstrated the therapeutic efficacy of the lavender oil preparation Silexan for the treatment of subsyndromal anxiety disorder in a randomized, double-blind, placebo controlled trial [5]. Lavender oil treatment was found to alleviate anxiety related symptoms such as restlessness, disturbed sleep as well as somatic complaints, whereby the product demonstrated good tolerability without provoking greater adverse effects [6].

However, only few reports on the specific neurobiochemical actions of lavender oil are available. Best discussed

* Correspondence: dietmar.fuchs@i-med.ac.at

³Division of Biological Chemistry, Biocenter, Medical University of Innsbruck, Innsbruck, Austria

Full list of author information is available at the end of the article

in the literature are the anxiolytic, analgesic and anti-inflammatory effects of one of the oil's principal components, linalool and its derivatives like linalyl acetate [7]. The antinociceptive potential of linalool was studied in several animal studies, e.g. it has been shown to interfere with glutamatergic transmission in mice [8,9] and to modify the nicotinic receptor-ion channel kinetics at the neuromuscular junction [10].

Importantly, immune activation and inflammation are strongly associated with an increase of mood disorders [11,12]. Several biochemical links between psychoneuroimmunology and neuropsychopharmacology have been dissected in the past. The catabolism of the essential amino acid tryptophan, known for its essential role in antimicrobial defence, has turned out as an important link between the immunological network and neuroendocrine functions [12,13]. The enzyme indoleamine 2,3-dioxygenase (IDO, EC 1.13.11.52) catalyses the rate-limiting step in the conversion of tryptophan to kynurenine and becomes highly activated in macrophages and is induced also in many other cell types upon exposure to pro-inflammatory cytokine interferon- γ (IFN- γ) signaling in the course of the cellular immune response [12]. Tryptophan depletion creates an anti-proliferative environment against target cells and contributes to the antimicrobial effects of activated macrophages [14]. However, a reduction in plasma tryptophan leads in consequence to low serotonin (5-HT) synthesis and further, several tryptophan breakdown products are known to exert neuroactive effects [11].

During the cellular immune response, in parallel to IDO, guanosine triphosphate (GTP)-cyclohydrolase-I (GTP-CH-I, EC 3.5.4.16) is induced by IFN- γ . GTP-CH-I is the key enzyme in the biosynthesis of neopterin, a marker molecule for immune system activation [15,16]. As tryptophan metabolism may occur not only via IDO but also via hepatic tryptophan 2,3-dioxygenase (TDO), a concomitant determination of immune activation marker neopterin is suitable to judge the contribution of inflammation in changes of tryptophan levels [12]. Neopterin levels and kynurenine to tryptophan ratio (Kyn/Trp) have turned out as useful markers for a variety of diseases that are associated with chronic immune activation such as infections, autoimmune syndromes, malignancies or neurodegeneration [12,16].

The aim of this study was to evaluate the effects of lavender essential oil and some of its constituents on tryptophan catabolism, by using the well established model system of freshly isolated human peripheral blood mononuclear cells (PBMC), stimulated or not with the mitogen phytohemagglutinin (PHA) [17]. Determination of Kyn/Trp and neopterin levels in cell culture supernatants is used as sensitive and reliable read-out for the activation status of PBMCs. The terpene alcohol (-)-linalool, a major constituent of lavender oil that is also contained in

several edible plant species, as well as two minor lavender oil constituents (+)- α -pinene and (+)-limonene were chosen as reference compounds for analysis, due to their reported anti-inflammatory properties [18,19].

Methods

Chemicals

Phytohemagglutinin (PHA), (+)- α -pinene, (-)-linalool and (+)-limonene were purchased from Sigma Aldrich (Vienna, Austria). Freshly prepared lavender oil (*Aetheroleum Lavanduli*) was purchased from a local pharmacy (Tiroler Adler Apotheke, Innsbruck, Austria).

Analytical methods

Analyses of lavender oil were performed on a Perkin-Elmer Autosystem gas chromatograph (Norwalk, USA) equipped with FID, split-splitless injector and a Perma-bond SE-54-DF capillary column (50 m x 0.32 ID; 0.25 μ m film thickness) from Macherey-Nagel (Düren, Germany). Helium (1 mL/min) was used as carrier gas, and injector and detector temperatures were set to 220 and 240°C, respectively. The injected sample volume was 0.5 μ L with a split ratio of 1:4. The following temperature gradient was applied for all separations: for the first 10 min isotherm at 85°C, then the temperature was increased to 180°C (10°C/min), to keep this setting for 5.5 min. For GC analysis, 60.0 mg of essential oil was dissolved in 5.00 mL dichloromethane (Chromasolv; Riedel-de-Haen, Seelze, Germany). For preparation of the standard solutions 30.0 mg of each compound ((+)- α -pinene, (-)-linalool and (+)-limonene) was placed in one 5.00 mL volumetric flask, which was filled up to volume with DCM (level 1). Further concentration levels were prepared by serial dilution using the same solvent, so that the covered linear range was from 6.0 to 0.08 mg/mL for all compounds. External calibration curves were constructed for the three available reference compounds, and the following regression equations were obtained: $y = 14429x - 336.59$ ((+)- α -pinene), $y = 16024x - 902.95$ ((+)-limonene) and $y = 15780x - 1407.70$ ((-)-linalool). The respective correlation coefficients were 0.9997 or higher.

Isolation of human PBMC

The study was performed in accordance with the Helsinki declaration. PBMC were isolated from whole blood obtained from healthy donors of whom informed consent was obtained that their donated blood might be used for scientific purposes in case when it was not selected for transfusion. The local ethics committee confirmed that no further approval is required for using anonymized leftover specimens from blood donations of the local blood bank for scientific purposes. Separation of blood cells was performed by density centrifugation (Lymphoprep, Nycomed

Pharma AS, Oslo, Norway) as described [17,20]. After isolation, PBMC were washed three times in phosphate buffered saline containing 1 mM EDTA. Cells were cultivated in RPMI 1640 supplemented with 10% heat-inactivated fetal calf serum (Biochrom, Berlin, Germany), 2 mM L-glutamine (Serva, Heidelberg, Germany) and 50 µg/mL gentamicin (Bio-Whittaker, Walkersville, MD) at 37°C in a humidified atmosphere containing 5% CO₂.

Cell treatment

In brief, isolated PBMC were seeded at a density of 1.5×10^6 cells/mL in supplemented RPMI 1640 medium and pre-incubated for 30 min with or without different concentrations of lavender oil [0.1 – 5%] or constituents ((+)- α -pinene [31.25 – 250 µM], (-)-linalool [25 – 1000 µM] and (+)-limonene [50 – 250 µM]). Then, cells were stimulated or not with 10 µg/mL PHA for 48 h. For each of the experiments run in duplicates, PBMC were prepared freshly from blood of at least three different healthy donors [20].

Cell viability

CellTiter-Blue assay (Promega, Germany) was used to determine cell viability. The half maximal (50% inhibitory) concentration (IC₅₀) was calculated by using the CalcuSyn software (Biosoft, UK) [21].

Measurement of tryptophan, kynurenine, neopterin and IFN- γ concentrations

After 48 h of incubation, the accumulated tryptophan breakdown and neopterin formation reach a plateau [17,20]. PBMC supernatants were harvested by centrifugation. Tryptophan and kynurenine concentrations were measured by high performance liquid chromatography (HPLC) using 3-nitro-L-tyrosine as an internal standard [22]. To estimate the activity of IDO, Kyn/Trp was calculated and expressed as µmol kynurenine/mmol tryptophan. Neopterin and IFN- γ concentrations were measured by ELISA (BRAHMS, Hennigsdorf/Berlin, Germany and R&D, Biomedica, Vienna, Austria) according to the manufacturers' instructions with a detection limit of 2 nmol/L neopterin and 8 pg/mL IFN- γ . Due to limited sample volumes, IFN- γ measurements were performed in a smaller set of 32 lavender oil treated samples only.

Statistical analysis

Data were analysed by using the Statistical Package for the Social Sciences (version 19, SPSS, Chicago, III, USA). To take into account that not all collected data followed a normal distribution, non-parametric Friedman and Wilcoxon signed-rank test were applied. P-values below 0.05 were considered to indicate significant differences.

Results

GC analysis

The three compounds (+)- α -pinene, (-)-linalool and (+)-limonene could be separated with GC and assigned in lavender essential oil (Figure 1). The respective retention times were found to be 5.72 min ((+)- α -pinene), 8.34 min ((+)-limonene) and 11.38 min ((-)-linalool). Their content in the essential oil sample was 3.70% ($\sigma_{rel} = 0.29$; $n = 3$) for (+)- α -pinene, 1.29% ($\sigma_{rel} = 0.11$) for (+)-limonene, and 33.09% ($\sigma_{rel} = 0.64$) for (-)-linalool.

Effect of lavender oil and its constituents on cell viability

Cytotoxicity was evaluated after 48 h incubation with lavender oil in a concentration range from 0.2 to 5.0% in unstimulated and PHA-stimulated PBMC. Treatment with the mitogen PHA [10 µg/mL] resulted in a significant increase of cell number due to the stimulated cell growth (Figure 2A). Reduced viability in the presence of lavender oil was measured with 0.5% addition in unstimulated and with 1.0% in PHA-stimulated cells, indicating the beginning of cytotoxic effects for higher treatment concentrations (Figure 2B). Viability decreased in a dose dependent manner, with IC₅₀ values of 2.29% in unstimulated and 1.79% in stimulated cells.

No cytotoxic effects in the tested concentration ranges could be observed for (-)-linalool (IC₅₀ \geq 1000 µM) and (+)-limonene (IC₅₀ \geq 250 µM), while for (+)- α -pinene IC₅₀ values of 195.41 µM in unstimulated PBMC and 127.36 µM in PHA-stimulated PBMC were determined.

Effect of lavender oil and its constituents on tryptophan breakdown

After an incubation period of 48 h, the average tryptophan concentration in supernatants of unstimulated PBMC was 31.5 ± 1.7 µmol/L (mean \pm SEM; $85.2 \pm 4.7\%$ of initial medium content) and the mean kynurenine concentration was 2.3 ± 0.7 µmol/L, resulting in Kyn/Trp of 79.0 ± 28.0 µmol/mmol (Figure 3). Upon stimulation of cells with PHA, tryptophan levels decreased to 11.3 ± 3.8 µmol/L, while the mean kynurenine concentration increased to 13.1 ± 3.3 µmol/L. The Kyn/Trp was increased in PHA-treated samples approximately 100-fold in comparison to the unstimulated supernatants (all $P < 0.05$).

By addition of lavender oil [0.1 – 5.0%] to unstimulated PBMC, no change of tryptophan concentrations but a significant decline of kynurenine and Kyn/Trp levels was apparent with the treatment concentrations of 0.2 to 5.0% (Figure 4). In PHA-stimulated PBMC, lavender oil suppressed tryptophan breakdown significantly at all treatment concentrations (0.1 to 5.0%), Kyn/Trp and kynurenine levels dropped dose-dependently and significantly (Figure 4). Addition of 1.0% of lavender oil





Figure 1 Determination of (+)- α -pinene, (-)-linalool and (+)-limonene in lavender oil. GC analysis of a standard mixture (A) and lavender essential oil (B).

completely inhibited tryptophan breakdown and restored tryptophan levels comparable to the concentration in unstimulated cell supernatants ($79.4 \pm 3.0\%$ of medium content).

While in PHA-treated cells, the incubation with 0.5% of lavender oil resulted in a suppression of kynurenine to $19.9 \pm 5.7\%$ of PHA control, the same concentration decreased kynurenine levels only to $50.2 \pm 10.1\%$ in unstimulated supernatants (compared to unstimulated control). The effect of lavender oil on IDO activity as indicated by Kyn/Trp was also significantly reduced in both, mitogen-stimulated and unstimulated PBMC. Likewise, this effect was more striking in PHA-stimulated cells, where 0.5% of lavender oil addition resulted in a reduction of IDO activity to $8.5 \pm 2.2\%$ compared to the PHA control. The same lavender concentration decreased Kyn/Trp only to $51.4 \pm 11.3\%$ in unstimulated PBMC (compared to unstimulated controls). Overall, lavender oil

induced effects were more pronounced in PHA-stimulated cells than in unstimulated ones.

Neither (-)-linalool treatment (up to $1000 \mu\text{M}$) nor (+)-limonene addition (up to $250 \mu\text{M}$) resulted in a significant change of the Kyn/Trp ratio in unstimulated PBMC (Figure 5A). (+)- α -pinene did not affect IDO activity up to a concentration of $125 \mu\text{M}$. Treatment of unstimulated PBMC with (+)- α -pinene at already cytotoxic concentrations ($250 \mu\text{M}$) resulted in a Kyn/Trp decrease of $19.4 \pm 8.0\%$ (Figure 5A). In PHA-stimulated cells, all constituents were able to strongly and significantly inhibit IDO activity in a dose-dependent manner at non-toxic concentrations (Figure 5B). With (+)-limonene and (+)- α -pinene, lower treatment concentrations led to a decrease of the Kyn/Trp ratio in stimulated cells ($25 \mu\text{M}$ and $62.5 \mu\text{M}$, respectively) than with (-)-linalool, where the inhibitory effect started with $125 \mu\text{M}$.





Effect of lavender oil on neopterin and IFN-γ production
 After 48 h, neopterin concentrations were significantly higher (10.5 ± 1.8 nmol/L) in culture supernatant of PHA-stimulated cells than in supernatants of unstimulated PBMC (3.8 ± 0.3 nmol/L) ($p < 0.05$; Figure 6A). Also, IFN-γ levels were increased in mitogen stimulated cells (209.7 ± 73.4 pg/mL) in comparison to untreated cells (19.3 ± 14.5 pg/mL) ($P < 0.05$).

The incubation of unstimulated PBMC with lavender oil resulted in the reduction of neopterin levels at all

treatment concentrations (Figure 6B). At a concentration of 0.5% lavender oil and higher, neopterin formation was maximally decreased to $85.4 \pm 3.4\%$ in comparison to the untreated control cells. In PHA-stimulated cells, neopterin concentrations were significantly and dose-dependently lowered in the concentration range from 0.5 - 5.0% of lavender oil, with a reduction to $67.6 \pm 15.5\%$ at 0.5% lavender oil treatment, in comparison to the PHA-treated control.

IFN-γ concentrations were not affected in unstimulated PBMC upon lavender oil treatment, while in stimulated







cells, IFN- γ levels decreased in a dose-dependent manner (data not shown). Addition of 0.5% of lavender oil resulted in a reduction of IFN- γ production until baseline levels of unstimulated cells (22.4 ± 5.9 pg/mL).

Discussion

Anti-inflammatory properties of lavender oil and its constituents have been reported in several *in vitro* and *in vivo* studies. Lavender essential oil and constituents have been shown to interfere with key immunological pathways, e.g. nuclear factor kappa B (NF- κ B) and p38 mitogen-activated protein kinase (MAPK) signaling as

well as cytokine secretion [19,23]. E.g., (+)- α -pinene, (-)-linalool and (+)-limonene were able to decrease interleukin-2 (IL-2) secretion and to increase the IL-10/IL-2 ratio in mouse primary splenocytes, which indicates their property to repress Th1 immune activation and suggest a potential inclination towards Th2 [19]. Furthermore, (-)-linalool was able to attenuate the production of lipopolysaccharide (LPS)-induced tumor necrosis factor α (TNF α) and IL-6 both in RAW 264.7 macrophages and in mice, and has been discussed as potential anti-inflammatory agent for preventing lung injury [19,23].





The impact of the reference substances in attenuating Th1 immune response agrees with results of our study, which showed that non-toxic concentrations of (+)- α -pinene, (-)-linalool and (+)-limonene were able to inhibit mitogen-stimulated IDO activity in a model system of freshly isolated PBMC.

Also, lavender oil treatment was able to dose-dependently inhibit both tryptophan breakdown and kynurenine formation in supernatants of mitogen-stimulated PBMC. This inhibitory effect could already be detected at lavender oil concentrations that affected cell viability only slightly (0.1 to 0.5%). At higher treatment concentrations, effects on tryptophan and kynurenine were even stronger, however also cytotoxic effects of lavender oil increased. Interestingly, it has been shown that kynurenine metabolites are able to induce Th1 cell apoptosis [24]. Thus, we suggest that at low concentrations, lavender oil might beneficially influence cell viability by counteracting pro-apoptotic signaling, while at higher concentrations toxicity effects become prevalent. In studies with several compounds in the PBMC model [17,20], IDO inhibition preceded substance toxicity phenomena, thus probably being a more sensitive indicator of cell death. Of note, the viability assay used in this study is based on the reduction of resazurin to fluorescent resorufin. Increased conversion rates may also indicate enhanced metabolic activity of cells, which does not always correlate with an increase in proliferation [25,26].

Importantly, in mitogen-stimulated cells, a suppressive effect of lavender oil treatment on neopterin and IFN- γ

concentrations could be observed. In unstimulated cells, lavender oil treatment had no influence on tryptophan and IFN- γ levels, but the formation of kynurenine and neopterin was suppressed to some extent. As PBMC were preincubated with the lavender oil before PHA addition, we suggest that the oil interferes mainly with IDO and GTP-CH-I stimulation. A basal activity of both enzymes is suggested to be present also in unstimulated cells, probably initiated due to the preceding cell isolation procedure. In unstimulated PBMC, tryptophan levels remained unaffected upon lavender oil treatment, e.g. with a 0.5% oil addition, 86.8 \pm 3.2% of the initial medium content of tryptophan, corresponding to \sim 32 μ mol/L, was still detectable after 48 h, while a significant reduction of kynurenine levels was observed. For 0.5% lavender oil treatment, kynurenine levels were reduced to 50.2 \pm 10.1% compared to the untreated control, which corresponds to a reduction from 2.3 \pm 0.7 μ mol/L to 0.8 \pm 0.1 μ mol/L.

Of note, changes in immune parameters, such as impaired activities of immunocompetent cells, and involvement of inflammatory mediators and pro-inflammatory cytokines have been reported to be associated with behavioural alterations by several studies, and cell-mediated immune activation is suggested to be an important factor in distinct mental disturbances [11]. Behavioural changes can be induced by altered cytokine levels, e.g. studies of IFN- α treated patients showed therapy-induced depressive symptoms associated with activation of neuroendocrine pathways and altered serotonin metabolism [11,27]. Within

the cellular immune response, pro-inflammatory pathways are strongly induced, including neopterin production via GTP-CH-I and tryptophan catabolism via IDO, and the concentrations of these biomarkers have been found to be altered in mental disorders or diseases associated mood disturbances [12]. Enhanced neopterin concentrations together with low serum levels of tryptophan caused by increased tryptophan breakdown were shown to correlate with neuropsychiatric abnormalities like cognitive decline and depressive symptoms especially in long-lasting and chronic diseases [28].

Beside the important role of tryptophan catabolism in the regulation of inflammatory responses [29], tryptophan is a source for the production of 5-hydroxytryptophan, an intermediate in the biosynthesis of neurotransmitter serotonin. In states of persistent immune activation, availability of free serum tryptophan is diminished and as a consequence of reduced serotonin production, serotonergic functions may as well be affected [12].

About 95% of the body's serotonin resides in the gut [30]. Furthermore, the gastrointestinal tract is rich in lymphocytes. Lavender oil treatment concentrations used for this *in vitro* study may appear relatively high, however *in vivo*, initial effects on IDO are suggested to be initiated already in the gastrointestinal tract, were such concentrations can be readily reached. In the study of Kasper et al., a treatment concentration of 80 mg/day was able to induce clinically meaningful and statistically significant anxiolytic effects [5].

Of note, deciphering specific bioactivities of isolated essential oil components is challenging because of the great number of constituents with similar physicochemical properties (e.g. lipophilic, high vapor pressure). In general, the major components reflect quite well the features of the essential oils from which they derive [31]. However minor constituent may contribute to the overall activity profile by modulating these activities and synergisms can play a major role [32]. (-)-Linalool is the most studied monoterpene regarding analgesic effects, a more detailed elucidation of its impact on the GABAergic system would help to dissect molecular details on its anti-convulsant, analgesic as well as anxiolytic activities [33]. For both (+)-limonene [34] and (+)- α -pinene [35] antinociceptive effects have been reported, however these effects are suggested to be strongly associated with their anti-inflammatory activities [33].

The here reported effect on tryptophan breakdown is not a unique property of lavender oil or the analysed constituents. In earlier studies using the identical cell-biological assay, similar effects on tryptophan metabolism have been found by investigating *Hypericum perforatum* extracts as well as Δ^9 -tetrahydrocannabinol and cannabidiol, indicating that the suppression of tryptophan breakdown and neopterin production might be an important

but a more general aspect in the action of psychoactive compounds [36,37]. Thus, although our findings are from *in vitro* experiments only, they might be relevant also for the *in vivo* situation.

Furthermore, kynurenine derivatives such as kynurenic and quinolinic acid and 3-hydroxykynurenine are known to be neuroactive and their hyper- or hypofunction is associated with neurological disorders and psychiatric diseases such as depression and schizophrenia [38]. The quinolinic acid to kynurenic acid ratio in the brain is discussed as a potential measure for conditions linked to excitotoxicity. Although both substances must be synthesized locally, because they are not able to cross the blood-brain barrier, other kynurenine pathway components such as tryptophan, kynurenine and 3-hydroxykynurenine can enter the brain, thus establishing a link between peripheral inflammation and brain tryptophan metabolism [39]. Additionally, also microglial cells and blood-borne cells within the brain can be stimulated to activate the kynurenine pathway in states of peripheral immune activation [38].

Scientific reports on the impact of IDO activity for different pathological conditions, including neuropsychiatric disturbances, account for IDO as a potential key pharmacological target. Several IDO inhibitors have been identified yet, and much effort will be necessary to evaluate their *in vivo* efficacy. The most prominent IDO inhibitor 1-methyl tryptophan (1-MT) has been shown to counteract microbial-induced depressive-like symptoms in animal studies [40]. Beside the synthetic design of IDO antagonists via rational design strategies, also a variety of endogenous and exogenous antioxidants, such as vitamins, food supplements or preservatives, have been shown to suppress tryptophan catabolism in cellular model systems [17,20]. Thereby, the modulation of tryptophan metabolism is suggested to be due to the interference of the test substances mostly with immune activation cascades, rather than directly with IDO enzyme, as often other immune-relevant molecules such as neopterin and IFN- γ are affected additionally. Also in this study, lavender oil treatment was able to reduce neopterin and IFN- γ levels in mitogen-treated PBMC.

Moreover, the increased production of neopterin during inflammation could also indirectly affect neurotransmitter concentrations. In human macrophages, neopterin is produced at the expense of tetrahydrobiopterin (BH₄), an essential cofactor for several monooxygenases including tryptophan, phenylalanine and tyrosine hydroxylase [15]. Thus, beside serotonin synthesis, also catecholamine formation depends on BH₄ availability. Interestingly, in depressed patients with a history of seasonal affective disorder, significantly lower plasma biopterin and tryptophan levels but elevated neopterin levels were found in comparison to healthy controls [41]. Thus, reduced BH₄ levels

in inflammatory conditions might negatively influence neurotransmitter production.

Conclusion

We could show that lavender oil can suppress mitogen-induced tryptophan degradation and IFN- γ production *in vitro*, and influence on kynurenine and neopterin formation in activated as well as to a lower extent in unstimulated PBMC.

Also, the constituents (-)-linalool, (+)- α -pinene and (+)-limonene showed dose-dependent inhibitory effects on tryptophan breakdown in PHA-stimulated PBMC. Thus, the IDO suppressing activities of lavender oil might at least partially result from the concerted action of the analysed constituents, where also minor components may play an essential role.

The finding that lavender essential oil, a medicinal plant-derived natural multicomponent preparation, may be a source of pharmacological active substances that interfere with key immune activation cascades such as the IDO and GTP-CH-I pathway, is of central relevance for the understanding of its therapeutic efficacy.

However, as the reported effects might not reflect the whole activity spectrum of lavender oil. Further research is necessary to elucidate other neuro-immunological relevant activities and to confirm the *in vivo* relevance of our findings.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Compounds characterization: MG; blood donations: HS; idea: JMG, FÜ, DF; cell culture work and read outs: JMG, KB, SS; HPLC measurements: SG, SS, DF; manuscript draft: JMG, FÜ, DF. All authors have read and contributed to the final version of the article. All authors read and approved the final manuscript.

Grants

This work was supported by the Austrian Research Promotion Agency (FFG) grant FFG 840590 and by the Austrian Science Fund (FWF) grant T 703. The content of this article does not necessarily reflect the views or policies of the funding sources.

Author details

¹Division of Medical Biochemistry, Biocenter, Medical University of Innsbruck, Innsbruck, Austria. ²Institute of Pharmacy, Pharmacognosy, University of Innsbruck, Innsbruck, Austria. ³Division of Biological Chemistry, Biocenter, Medical University of Innsbruck, Innsbruck, Austria. ⁴Central Institute of Blood Transfusion and Immunology, University Hospital, Innsbruck, Austria.

Received: 10 January 2014 Accepted: 9 December 2014

Published: 16 December 2014

References

1. Woronuk G, Demissie Z, Rheault M, Mahmoud S: **Biosynthesis and therapeutic properties of Lavandula essential oil constituents.** *Planta Med* 2011, **77**:7–15.
2. Da Porto C, Decorti D, Kikic I: **Flavour compounds of Lavandula angustifolia L. To use in food manufacturing: comparison of three different extraction methods.** *Food Chem* 2009, **112**:1072–1078.
3. Perry N, Perry E: **Aromatherapy in the management of psychiatric disorders: clinical and neuropharmacological perspectives.** *CNS Drugs* 2006, **20**:257–280.
4. Cavanagh HM, Wilkinson JM: **Biological activities of lavender essential oil.** *Phytother Res* 2002, **16**:301–308.
5. Kasper S, Gastpar M, Muller WE, Volz HP, Moller HJ, Dienel A, Schlafke S: **Silexan, an orally administered Lavandula oil preparation, is effective in the treatment of 'subsyndromal' anxiety disorder: a randomized, double-blind, placebo controlled trial.** *Int Clin Psychopharmacol* 2010, **25**:277–287.
6. Doroshenko O, Rokitta D, Zadoyan G, Klement S, Schlafke S, Dienel A, Gramatte T, Luck H, Fuhr U: **Drug cocktail interaction study on the effect of the orally administered lavender oil preparation silexan on cytochrome p450 enzymes in healthy volunteers.** *Drug Metab Dispos* 2013, **41**:987–993.
7. Hajhashemi V, Ghannadi A, Sharif B: **Anti-inflammatory and analgesic properties of the leaf extracts and essential oil of Lavandula angustifolia Mill.** *J Ethnopharmacol* 2003, **89**:67–71.
8. Silva Brum LF, Emanuelli T, Souza DO, Elisabetsky E: **Effects of linalool on glutamate release and uptake in mouse cortical synaptosomes.** *Neurochem Res* 2001, **26**:191–194.
9. Hossain SJ, Hamamoto K, Aoshima H, Hara Y: **Effects of tea components on the response of GABA(A) receptors expressed in Xenopus Oocytes.** *J Agric Food Chem* 2002, **50**:3954–3960.
10. Re L, Barocci S, Sonnino S, Mencarelli A, Vivani C, Paolucci G, Scarpantonio A, Rinaldi L, Mosca E: **Linalool modifies the nicotinic receptor-ion channel kinetics at the mouse neuromuscular junction.** *Pharmacol Res* 2000, **42**:177–182.
11. Haroon E, Raison CL, Miller AH: **Psychoneuroimmunology meets neuropsychopharmacology: translational implications of the impact of inflammation on behavior.** *Neuropsychopharmacology* 2012, **37**:137–162.
12. Widner B, Laich A, Sperner-Unterwieser B, Ledochowski M, Fuchs D: **Neopterin production, tryptophan degradation, and mental depression—what is the link?** *Brain Behav Immun* 2002, **16**:590–595.
13. Muller N, Myint AM, Schwarz MJ: **Inflammatory biomarkers and depression.** *Neurotox Res* 2011, **19**:308–318.
14. Murray HW, Szuro-Sudol A, Wellner D, Oca MJ, Granger AM, Libby DM, Rothermel CD, Rubin BY: **Role of tryptophan degradation in respiratory burst-independent antimicrobial activity of gamma interferon-stimulated human macrophages.** *Infect Immun* 1989, **57**:845–849.
15. Werner ER, Werner-Felmayer G, Fuchs D, Hausen A, Reibnegger G, Yim JJ, Pfeleiderer W, Wachter H: **Tetrahydrobiopterin biosynthetic activities in human macrophages, fibroblasts, THP-1, and T 24 cells. GTP-cyclohydrolase I is stimulated by interferon-gamma, and 6-pyruvoyl tetrahydropterin synthase and sepiapterin reductase are constitutively present.** *J Biol Chem* 1990, **265**:3189–3192.
16. Murr C, Widner B, Wirleitner B, Fuchs D: **Neopterin as a marker for immune system activation.** *Curr Drug Metab* 2002, **3**:175–187.
17. Becker K, Schroecksnadel S, Gostner J, Zaknun C, Schennach H, Uberall F, Fuchs D: **Comparison of in vitro tests for antioxidant and immunomodulatory capacities of compounds.** *Phytomedicine* 2013, **21**:164–171.
18. Huo M, Cui X, Xue J, Chi G, Gao R, Deng X, Guan S, Wei J, Soromou LW, Feng H, Wang D: **Anti-inflammatory effects of linalool in RAW 264.7 macrophages and lipopolysaccharide-induced lung injury model.** *J Surg Res* 2013, **180**:e47–e54.
19. Ku CM, Lin JY: **Anti-inflammatory effects of 27 selected terpenoid compounds tested through modulating Th1/Th2 cytokine secretion profiles using murine primary splenocytes.** *Food Chem* 2013, **141**:1104–1113.
20. Jenny M, Klieber M, Zaknun D, Schroecksnadel S, Kurz K, Ledochowski M, Schennach H, Fuchs D: **In vitro testing for anti-inflammatory properties of compounds employing peripheral blood mononuclear cells freshly isolated from healthy donors.** *Inflamm Res* 2011, **60**:127–135.
21. Chou TC, Talalay P: **Quantitative analysis of dose-effect relationships: the combined effects of multiple drugs or enzyme inhibitors.** *Adv Enzyme Regul* 1984, **22**:27–55.
22. Widner B, Werner ER, Schennach H, Wachter H, Fuchs D: **Simultaneous measurement of serum tryptophan and kynurenine by HPLC.** *Clin Chem* 1997, **43**:2424–2426.
23. Huang MY, Liao MH, Wang YK, Huang YS, Wen HC: **Effect of lavender essential oil on LPS-stimulated inflammation.** *Am J Chin Med* 2012, **40**:845–859.
24. Fallarino F, Grohmann U, Vacca C, Bianchi R, Orabona C, Spreca A, Fioretti MC, Puccetti P: **T cell apoptosis by tryptophan catabolism.** *Cell Death Differ* 2002, **9**:1069–1077.

25. O'Brien J, Wilson I, Orton T, Pognan F: **Investigation of the Alamar Blue (resazurin) fluorescent dye for the assessment of mammalian cell cytotoxicity.** *Eur J Biochem* 2000, **267**:5421–5426.
26. Rampersad SN: **Multiple applications of Alamar Blue as an indicator of metabolic function and cellular health in cell viability bioassays.** *Sensors (Basel)* 2012, **12**:12347–12360.
27. Capuron L, Neurauder G, Musselman DL, Lawson DH, Nemeroff CB, Fuchs D, Miller AH: **Interferon-alpha-induced changes in tryptophan metabolism. Relationship to depression and paroxetine treatment.** *Biol Psychiatry* 2003, **54**:906–914.
28. Schrocksnadel K, Wirleitner B, Winkler C, Fuchs D: **Monitoring tryptophan metabolism in chronic immune activation.** *Clin Chim Acta* 2006, **364**:82–90.
29. Mellor AL, Munn DH: **Tryptophan catabolism and regulation of adaptive immunity.** *J Immunol* 2003, **170**:5809–5813.
30. Gershon MD, Tack J: **The serotonin signaling system: from basic understanding to drug development for functional GI disorders.** *Gastroenterology* 2007, **132**:397–414.
31. Ipek E, Zeytinoglu H, Okay S, Tuylu BA, Kurkcuoglu M, Husnu Can Baser K: **Genotoxicity and antigenotoxicity of Origanum oil and carvacrol evaluated by Ames Salmonella/microsomal test.** *Food Chem* 2005, **93**:551–556.
32. Bakkali F, Averbeck S, Averbeck D, Idaomar M: **Biological effects of essential oils—a review.** *Food Chem Toxicol* 2008, **46**:446–475.
33. Guimaraes AG, Quintans JS, Quintans LJ Jr: **Monoterpenes with analgesic activity—a systematic review.** *Phytother Res* 2013, **27**:1–15.
34. do Amaral JF, Silva MI, Neto MR, Neto PF, Moura BA, de Melo CT, de Araujo FL, de Sousa DP, de Vasconcelos PF, de Vasconcelos SM, de Sousa FC: **Antinociceptive effect of the monoterpene R-(+)-limonene in mice.** *Biol Pharm Bull* 2007, **30**:1217–1220.
35. Him A, Ozbek H, Turel I: **Antinociceptive activity of alpha-pinene and fenchone.** *Pharmacology Online* 2008, **3**:363–369.
36. Winkler C, Wirleitner B, Schrocksnadel K, Schennach H, Fuchs D: **St. John's wort (*Hypericum perforatum*) counteracts cytokine-induced tryptophan catabolism in vitro.** *Biol Chem* 2004, **385**:1197–1202.
37. Jenny M, Santer E, Pirich E, Schennach H, Fuchs D: **Delta9-tetrahydrocannabinol and cannabidiol modulate mitogen-induced tryptophan degradation and neopterin formation in peripheral blood mononuclear cells in vitro.** *J Neuroimmunol* 2009, **207**:75–82.
38. Schwarcz R, Bruno JP, Muchowski PJ, Wu HQ: **Kynurenines in the mammalian brain: when physiology meets pathology.** *Nat Rev Neurosci* 2012, **13**:465–477.
39. Saito K, Nowak TS Jr, Suyama K, Quearry BJ, Saito M, Crowley JS, Markey SP, Heyes MP: **Kynurenine pathway enzymes in brain: responses to ischemic brain injury versus systemic immune activation.** *J Neurochem* 1993, **61**:2061–2070.
40. O'Connor JC, Lawson MA, Andre C, Moreau M, Lestage J, Castanon N, Kelley KW, Dantzer R: **Lipopolysaccharide-induced depressive-like behavior is mediated by indoleamine 2,3-dioxygenase activation in mice.** *Mol Psychiatry* 2009, **14**:511–522.
41. Hoekstra R, Fekkes D, van de Wetering BJ, Pepplinkhuizen L, Verhoeven WM: **Effect of light therapy on biopterin, neopterin and tryptophan in patients with seasonal affective disorder.** *Psychiatry Res* 2003, **120**:37–42.

doi:10.1186/1472-6882-14-503

Cite this article as: Gostner et al.: Lavender oil suppresses indoleamine 2,3-dioxygenase activity in human PBMC. *BMC Complementary and Alternative Medicine* 2014 **14**:503.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

