

RESEARCH ARTICLE

SARS-CoV-2 S Protein Subunit 1 Elicits Ca²⁺ Influx – Dependent Ca²⁺ Signals in Pancreatic Stellate Cells and Macrophages *In Situ*


Julia V Gerasimenko , Ole H Petersen , Oleg V Gerasimenko *

School of Biosciences, Sir Martin Evans Building, Cardiff University, Wales CF10 3AX, UK

*Address correspondence to O.G. (e-mail: GerasimenkoOV@cardiff.ac.uk)

Abstract

The S protein subunit 1 (S1) of SARS-CoV-2 is known to be responsible for the binding of the virus to host cell receptors, but the initial intracellular signalling steps following receptor activation of cells in the exocrine pancreas are unknown. Using an intact live mouse pancreatic lobule preparation, we observed that S1 elicited Ca²⁺ signals in stellate cells and macrophages, but not in the dominant acinar cells. The Ca²⁺ signals occurred mostly in the form of repetitive Ca²⁺ spikes. The probability of observing Ca²⁺ signals depended on the S1 concentration. The threshold was close to 70 nM, whereas at 600 nM, all cells responded. The SARS-CoV-2 nucleocapsid protein did not elicit any Ca²⁺ signals in any of the three cell types tested. The S1-induced Ca²⁺ signals in stellate cells started much faster (122 ± 37s) than those in macrophages (468 ± 68s). Furthermore, the interleukin-18 binding protein (IL-18BP) abolished the responses in macrophages without affecting the Ca²⁺ signals in stellate cells. The S1-elicited Ca²⁺ signals were completely dependent on the presence of external Ca²⁺ and were abolished by a selective inhibitor (CM4620) of Orai1 Ca²⁺ Release Activated Ca²⁺ channels. SARS-CoV-2 may contribute to acute pancreatitis, an often fatal inflammatory human disease. The S1-elicited Ca²⁺ signals we have observed in the pancreatic stellate cells and endogenous macrophages may play an important part in the development of the inflammatory process.



Key words: SARS-COV-2; S1 spike protein; calcium signalling; pancreatic macrophages; pancreatic stellate cells; interleukin-18; IL-18BP; ORAI1 CRAC channels; CM4620

Introduction

SARS-CoV-2 enters cells via receptor-mediated endocytosis¹, using binding of its S protein subunit to ACE2 receptors thereby

allowing endocytic uptake of the virus². SARS-CoV-2 is known to enter a number of cell types in the human body including the respiratory tract², the central nervous system³ and the gastrointestinal tract, including the exocrine pancreas⁴, with several

Submitted: 3 January 2022; Revised: 21 January 2022; Accepted: 24 January 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of American Physiological Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

reported cases of acute pancreatitis (AP)⁴. The ACE2 receptor is present in the pancreas⁴, but the mechanism by which SARS-CoV-2 can induce AP as well as the type of cells targeted by the virus are currently unknown.

Pancreatic acinar cells dominate the exocrine pancreas and are largely responsible for the initiation of AP⁵. However, other cell types in this tissue also play important roles, in both physiological and pathophysiological processes, including AP⁵⁻¹¹. Our novel method of using isolated lobules of the exocrine pancreas^{7,9,10}, allowing preservation of the normal microscopic structure of the pancreatic environment, enables simultaneous recordings of intracellular signals i.e., cytosolic Ca²⁺⁶⁻¹¹ and nitric oxide¹¹ in several different cell types, including stellate cells⁹, intrinsic nerves¹⁰ and pancreatic macrophages⁷.

Whereas physiological and pathological Ca²⁺ signals in pancreatic acinar cells are well understood⁵, their roles in the stellate cells⁹ and the macrophages⁷ are less clear. There is, however, evidence that Ca²⁺ and NO signals in stellate cells amplify the pathological effects in AP⁸⁻¹¹. In our recent work⁷ we have studied immune cell Ca²⁺ signalling in response to stimulation with ATP and identified the responding cells as macrophages by immunohistochemistry with external markers specific for this cell type¹²⁻¹⁴. Macrophages are involved in the development of both AP and pancreatic cancer¹⁵, leading to inflammation and exacerbation of the patients' conditions¹⁶, but the evidence for this has largely been limited to work on cell cultures or fixed tissue samples of the pancreas. Taking advantage of the pancreatic lobule preparation, we now report for the first time Ca²⁺ signals induced by the SARS-CoV-2 S1 protein in pancreatic cells.

Methods

Animals

Ethical Approval: All animal studies were ethically reviewed and conducted according to the UK Animals (Scientific Procedures) Act, 1986. All experimental protocols were performed under a Project Licence granted by the UK Home Office and approved by the Animal Care and Ethics Committees at Cardiff School of Biosciences, Cardiff University. Animals were maintained in plastic cages supplied with fresh corn cob bedding, tap water, and commercial pelleted diet.

Lobule Preparation

Pancreatic lobules were freshly isolated from the pancreas of 5- to 7-week-old male C57BL6/J mice as previously described^{7,10}. The pancreas was rapidly removed, injected with standard Na⁺-Hepes-based solution containing collagenase and incubated for 5-6 min at 37°C. The standard solution was composed of (in mM): NaCl, 140; KCl, 4.8; Hepes, 10; MgCl₂, 1; glucose, 10; CaCl₂, 1 (unless stated otherwise), pH 7.3 (NaOH). All experiments were carried out with pancreatic lobules attached to the coverslip of a perfusion chamber at room temperature (22°C).

Ca²⁺ Measurements

Pancreatic lobules were loaded with Fluo-4 acetoxymethyl (AM) ester at room temperature as described previously⁷. The lobules were transferred into a flow chamber and perfused with the standard solution alone or containing different chemicals as described in the experimental protocols of the result section. Cells were visualized using a Leica SP5 MPEII two-photon confocal

microscope, with an x63 1.2NA objective lens. The Fluo-4 excitation wavelength was 488 nm and emission was collected at 500-560 nm with resolution of 256 × 256 pixels and speed of 0.7 frames/s. Images were analysed using Leica Confocal Software (Leica, Mannheim, Germany). Fluorescence signals were plotted as normalized F/F₀. ANOVA or Student's t-test were performed for statistical analysis.

Immunostaining in ExVivo Pancreatic Lobules

Immunostaining of live pancreatic lobules was performed as previously described⁷. Mouse F4/80 Alexa Fluor 647-conjugated monoclonal rat antibodies were used to label specific surface proteins of immune cells, usually at the end of Ca²⁺ measurement experiments, unless otherwise stated. After blocking with 1% BSA and 10% goat serum containing PBS, the isolated pancreatic lobules were incubated for 1 h at room temperature with the selected antibody. Antibody staining was visualized by exciting Alexa Fluor 647 with 633 nm laser at 10% power and emitted light was collected at 640-700 nm. Conjugated antibody fluorescence was also overlaid with Fluo-4 staining as described in the Ca²⁺ measurements section. Lobules were attached to the glass coverslips covered with poly-L-lysine.

Materials

SARS-CoV-2 S1 protein and Nuclear Capsid protein were purchased from GenScript or Abcam; S1 RBD (Receptor Binding Domain) was obtained from Abcam. IL-18BP was from R&D. Bradykinin (BK) was purchased from Tocris Biosciences (UK). Fluo-4 AM and Hoechst 33342 were purchased from ThermoFisher Scientific (UK). Anti-mouse F4/80 monoclonal rat antibody (CI-A3-1) [Alexa Fluor 647] were obtained from Novus Biologicals. CM4620 was provided by CalciMedica Inc., US. Other chemicals were purchased from Sigma or Calbiochem (Merck, UK).

Results

As previously reported^{7,10}, pancreatic macrophages are typically sensitive to ATP or ADP and display Ca²⁺ signals when stimulated by these agents, whereas pancreatic stellate cells are typically insensitive to ATP, but are always responsive to bradykinin (BK), displaying Ca²⁺ signals when stimulated by the nona-peptide. Pancreatic acinar cells are always sensitive to acetylcholine and cholecystokinin, but never respond to stimulation with BK. They can occasionally display Ca²⁺ signals in response to stimulation with ATP. Nevertheless, ATP-sensitive cells are most likely to be macrophages and this can be verified after the recording of Ca²⁺ signals by staining with fluorescently labelled antibodies for macrophages (F4/80) (Figure 1A-D). In the macrophages, untagged S1 protein (200 nM) elicited either a single Ca²⁺ spike (Figure 1A, 38% of cells tested, n = 17) or a train of such spikes (Figure 1B, 40% of cells tested, n = 18). Similar results were obtained from the BK-sensitive stellate cells (Figures 1C, D, n = 23). A lower concentration of S1 (70nM) evoked Ca²⁺ signals in a smaller proportion of cells (33% of macrophages and 25% of stellate cells, Figure 1E), whereas a higher concentration of S1 (600nM) induced Ca²⁺ signals in every stellate cell and macrophage tested (Figure 1E). The latency after S1 application, before the first Ca²⁺ signal appeared, was much longer in the case of the macrophages than for the stellate cells (Figure 1F, P < 0.006). The S1-elicited Ca²⁺ signals in




Figure 1. Pancreatic immune and stellate cells exhibit Ca^{2+} signals in response to the addition of SARS-CoV2 S1 protein. A-B. Representative traces demonstrate single and multiple Ca^{2+} signals from pancreatic immune cells in response to S1 addition (Pancreatic macrophages (PM), green traces). Lobules were stimulated with $2.5 \mu\text{M}$ ATP, followed by application of 200 nM S1 (for 20 min). At the end of each experiment $2.5 \mu\text{M}$ ATP was consecutively added ($n = 50$). At the end of Ca^{2+} measurements, lobules were stained under the microscope with fluorescently labelled antibodies for macrophages (F4/80) and nuclear staining with Hoechst 33342. Images of nuclear staining (Hoechst 33342, Fig 1Aa), calcium dye (Fluo-4, Fig 1Ab) and antibody staining (AlexaFluo 647, F4/80, Fig 1Ac) for the same area as for the trace in Fig 1A (images in Fig 1Aa-c with overlay presented in Fig 1Ad; scale bar $10 \mu\text{m}$). C-D. Representative traces of single (C) and multiple (D) Ca^{2+} responses from pancreatic stellate cells (PSC, red traces) in pancreatic lobules. Lobules were stimulated with 200 nM S1 for 20 min. At the end of each experiment 5 nM and/or 30 nM BK were sequentially added ($n = 23$). (E) Concentration-dependence of PSCs and PMs responding to S1 stimulation by Ca^{2+} signals (red and green data bars, respectively). No responses were observed from acinar cells (PAC, blue data bars). The number of responses was calculated including single and multiple Ca^{2+} spikes to different concentrations of S1 (70 nM , 200 nM , 600 nM). (F) Time between application of S1 (200 nM) and beginning of elevation of the cytosolic Ca^{2+} concentration in PSCs ($122 \pm 37 \text{ s}$) and PMs ($468 \pm 68 \text{ s}$). PSCs responded significantly faster, $P < 0.006$. (G) Representative responses to S1 in the presence of IL-18 binding protein (IL-18BP) in PMs and PSCs. (H) No Ca^{2+} signals to S1 were observed in the presence of IL-18BP (600 nM) in PMs and PACs ($n = 7$ and $n = 33$, respectively) while 63.5% of PSCs did respond ($n = 11$), similarly to results shown in Fig 1E.




Figure 2. SARS-CoV2 RBD protein elicits Ca^{2+} signals in macrophages and stellate cells. A-B. Representative traces demonstrate single and multiple Ca^{2+} signals from PMs in response to RBD addition (PM, green traces). Lobules were stimulated with $2.5 \mu\text{M}$ ATP followed by application of 200 nM RBD (for 20 min). At the end of each experiment $2.5 \mu\text{M}$ ATP and 5 nM BK were consecutively added ($n = 16$). C-D. Representative traces of single (C) and multiple (D) Ca^{2+} spikes from PSCs (red traces). Lobules were stimulated with 200 nM RBD for 20 min. At the end of each experiment $2.5 \mu\text{M}$ ATP and 5 nM BK were sequentially added ($n = 17$). (E) The percentage of cells responding to 200 nM RBD with single or multiple Ca^{2+} signals. PACs, blue; PMs, green and PSC, red.

macrophages were abolished by the IL-18 binding protein IL-18BP (Fig 1G, H), whereas the responses to S1 in the stellate cells were unchanged (63.5%, $n = 11$). S1 never elicited any Ca^{2+} signals in the acinar cells ($n > 100$). We have also tested the effect of the S1 receptor binding domain (RBD). As seen in Figure 2A-E, RBD at a concentration of 200 nM, elicited Ca^{2+} signals in 69% of the macrophages ($n = 16$) and 71% of the stellate cells tested ($n = 17$).

We did not observe any spontaneous (in the absence of S1 or RBD) Ca^{2+} signals in any of the stellate cells or macrophages

tested (Supplementary Figure A and B; $n = 4$ and $n = 9$, respectively). In another series of control experiments, the potential effects of SARS-CoV-2 nuclear capsid protein (200 nM) was tested, but this protein never induced any Ca^{2+} signals in any of the cell types (Supplementary Figure C and D; $n = 8$ and $n = 5$, respectively).

We investigated whether the S1-elicited Ca^{2+} signals were dependent on the presence of Ca^{2+} in the external solution. In the absence of external Ca^{2+} , as seen in Figures 3A and B, S1 did not elicit any change in the cytosolic Ca^{2+} concentration in




Figure 3. Removal of external Ca^{2+} or application of the Orai1 inhibitor CM4620 block S1-induced Ca^{2+} signals in pancreatic immune and stellate cells. A-B. Representative traces showing that there were no Ca^{2+} signals in response to 200 nM S1 in PMs ($n = 11$, green trace) and PSCs ($n = 9$, red trace) in the absence of external Ca^{2+} . 25 nM BK and 2.5 μM ATP were added at the end of each experiment. C-D. Representative traces demonstrating the absence of Ca^{2+} signals in response to 200 nM S1 in PMs ($n = 11$, green trace) and PSCs (red trace; only 2 out of the 8 tested cells responded with Ca^{2+} signals of reduced amplitude and duration), when treated with the Orai1 CRAC channel inhibitor CM4620 (10 μM). 2.5 μM ATP was added at the end of each experiment.

macrophages that were responsive to ATP ($n = 11$) or in stellate cells ($n = 9$) that displayed Ca^{2+} signals in response to stimulation with BK. In both these cell types, it has previously been shown that an inhibitor of the opening of Orai1 Ca^{2+} Release Activated Ca^{2+} (CRAC) channels markedly reduces the duration

of Ca^{2+} signals evoked by ATP or BK^{7,9}. As shown in Figure 3C and D, the Orai1 inhibitor CM4620 abolished S1-elicited Ca^{2+} signal generation in the macrophages ($n = 11$) and markedly inhibited or abolished Ca^{2+} signals in 75% of the stellate cells tested ($n = 8$).

Discussion

There is general agreement that ACE2, the SARS-CoV-2 receptor, is expressed in the exocrine pancreas, but little is known about its distribution between the different cell types^{17–19}. Specifically, there is currently no information on the presence or absence of ACE2 in the pancreatic stellate cells or macrophages. Our finding that S1 and its RBD elicits Ca²⁺ signals in both stellate cells and macrophages provides the first evidence for functional receptors, but not necessarily on both cell types. The much longer delay in the response to S1 in the macrophages (Figure 1F), as compared to the stellate cells, might indicate an indirect action of S1 on the macrophages. This conclusion is strengthened by the experiments (Figure 1G) showing that the IL-18 binding protein IL-18BP abolished S1-elicited Ca²⁺ signal generation in the macrophages but not in the stellate cells. The simplest explanation would be that S1-elicited Ca²⁺ signals in the stellate cells stimulate IL-18 secretion, which then in turn elicits Ca²⁺ signals in the macrophages. Stellate cells are capable of IL-18 production and, in the case of chronic pancreatitis, it has been shown that levels of IL-18 are increased following activation by lipopolysaccharide (LPS) or TGF- β ²⁰. Although plausible, and compatible with our data, the hypothesis that the S1-elicited Ca²⁺ signals in the macrophages are mediated by IL-18 from the stellate cells needs to be tested further in future experiments. The absence of Ca²⁺ signals in the acinar cells in response to S1 does not rule out the existence of receptors on these cells, as receptor activation could elicit effects other than Ca²⁺ signals but, so far, there are no data providing evidence for functional receptors on the acinar cells.

At this stage there is no information about the intracellular steps leading to the generation of the Ca²⁺ signals, but the fact that these signals, in contrast to those evoked in the same cells by ATP or BK, are completely dependent on the presence of external Ca²⁺ and can be inhibited by the Orai1 CRAC channel inhibitor CM4620, suggests a mechanism of generation that is somewhat different from that of the physiological agonists.

Based on our previous studies of Ca²⁺ and NO signalling in pancreatic stellate cells^{9,10} and Ca²⁺ signalling in pancreatic macrophages⁷, we have proposed that AP, although initiated in the acinar cells, progresses to a severe and often fatal disease state through necrotic amplification loops, sustained by excessive Ca²⁺ entry through CRAC channels in stellate cells, macrophages and acinar cells. We have specifically provided evidence for a number of interaction pathways between acinar cells and stellate cells and between acinar cells and macrophages⁵. The new data presented here suggest a further pathway, not hitherto recognized, between stellate cells and macrophages in which IL-18—secreted by stellate cells in response to activation of SARS-CoV-2 receptors—acts on macrophages to generate Ca²⁺ signals in these cells. IL-18 levels are elevated in AP, and it has previously been suggested that this agent may play a role in acute and chronic pancreatitis together with several other interleukins^{21–23}.

The S1 and RBD-elicited Ca²⁺ signals in macrophages and stellate cells, demonstrated in this study, would be likely to exacerbate an otherwise mild case of AP initiated, for example, by the primary action of bile acids or a combination of ethanol and fatty acids on the acinar cells, and could help to explain the known cases of severe AP associated with SARS-CoV-2 infection^{4,24}. Our finding that the S1- and RBD-elicited Ca²⁺ signals can be inhibited by an Orai1 CRAC channel blocker, is a further argument in favour of this pharmacological treatment against AP^{5,25}.

Funding

The work was supported by grants from the Medical Research Council (UK) (MR/J002771/1 and G19/22/2 to OHP) and Children with Cancer UK grants (17/248 and 19/288 to OVG and JVG).

Conflict of Interest Statement

OHP holds the position of Editor-in-Chief for Function and is blinded from reviewing or making decisions on the manuscript. JVG holds the position of Editorial Board Member for Function and is blinded from reviewing or making decisions for the manuscript. OVG has no conflicts to disclose.

Data Availability

The data underlying this article will be shared on reasonable request to the corresponding author.

References

- Ou X, Liu Y, Lei X, et al. Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Nat Commun* 2020;11(1):1620.
- Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020;181(2):271–280.e8.
- Steardo L, Steardo L, Jr, Verkhatsky A, A. Neuroinfection may contribute to pathophysiology and clinical manifestations of COVID-19. *Acta Physiologica* 2020;229(3):e13473
- Hadi A, Werge M, Kristiansen KT, et al. Coronavirus disease-19 (COVID-19) associated with severe acute pancreatitis: case report on three family members. *Pancreatol*, 2020;20(4):665–667
- Petersen OH, Gerasimenko JV, Gerasimenko OV, et al. The roles of calcium and ATP in the physiology and pathology of the exocrine pancreas. *Physiol Rev*, 2021,101(4):1691–1744
- Pallagi P, Madacsi T, Varga A, et al. Intracellular Ca²⁺ signalling in the pathogenesis of acute pancreatitis: recent advances and translational perspectives. *Int J Mol Sci* 2020;21(11):4005
- Gryshchenko O, Gerasimenko JV, Petersen OH, Gerasimenko OV. Calcium signaling in pancreatic immune cells in situ. *FUNCTION* 2020;2(1):zqaa026.
- Ferdek PE, Jakubowska MA, Gerasimenko JV, et al. Bile acids induce necrosis in pancreatic stellate cells dependent on calcium entry and sodium-driven bile uptake. *J Physiol* 2016;594(21):6147–6164.
- Gryshchenko O, Gerasimenko JV, Gerasimenko OV, et al. Ca²⁺ signals mediated by bradykinin type 2 receptors in normal pancreatic stellate cells can be inhibited by specific Ca²⁺ channel blockade. *J Physiol* 2016;594(2):281–293.
- Gryshchenko O, Gerasimenko JV, Peng S, et al. Calcium signalling in the acinar environment of the exocrine pancreas: physiology and pathophysiology. *J Physiol* 2018;596(14):2663–2678.
- Jakubowska MA, Ferdek PE, Gerasimenko OV, et al. Nitric oxide signals are interlinked with calcium signals in normal pancreatic stellate cells upon oxidative stress and inflammation *Open Biology* 2016;6(8):160149.
- Gundra UM, Girgis NM, Ruckerl D, et al. Alternatively activated macrophages derived from monocytes and tissue macrophages are phenotypically and functionally distinct. *Blood* 2014;123(20): e110–e122.

- 13 Schulz C, Gomez Perdiguero E, Chorro L, et al. A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science* 2012;**336**(6077):86–90.
14. Weisberg SP, Carpenter DJ, Chait M, et al. Tissue-resident memory T cells mediate immune homeostasis in the human pancreas through the PD-1/PD-L1 pathway. *Cell Rep* 2019;**29**(12):3916–3932.e5
15. Calderon B, Carrero JA, Ferris ST, et al. The pancreas anatomy conditions the origin and properties of resident macrophages. *J Exp Med* 2015;**212**(10):1497–1512
- 16 Russo MW, Wei JT, Thiny MT, et al. Digestive and liver diseases statistics. *Gastroenterology* 2004;**126**(5):1448–1453
17. Coate, et al. SARS-CoV-2 Cell Entry Factors ACE2 and TMPRSS2 Are Expressed in the Microvasculature and Ducts of Human Pancreas but Are Not Enriched in b Cells. *Cell Metab* 2020;**32**(6):1028–1040.e4
18. Liu F, Long X, Zhang B, et al. ACE2 Expression in Pancreas May Cause Pancreatic Damage After SARS-CoV-2 Infection. *Clin Gastroenterol Hepatol* 2020;**18**(9):2128–2130.e2.
19. Müller JA, Groß R, Conzelmann C, et al. SARS-CoV-2 infects and replicates in cells of the human endocrine and exocrine pancreas. *Nature Metabolism* 2021;**3**(2):149–165
20. Li CX, Cui LH, Zhang LQ, et al. Role of NLR family pyrin domain-containing 3 inflammasome in the activation of pancreatic stellate cells. *Exp Cell Res* 2021;**404**(2):112634
21. Ueda T, Takeyama Y, Yasuda T, et al. Significant elevation of serum interleukin-18 levels in patients with acute pancreatitis. *J Gastroenterol* 2006;**41**(2):158–165
22. Schneider A, Haas SL, Hildenbrand R, et al. Enhanced expression of interleukin-18 in serum and pancreas of patients with chronic pancreatitis. *World J Gastroenterol* 2006;**12**(40):6507–6514
23. Pastor CM, Morel DR, Vonlaufen A, et al. *Pancreatology* 2011;**10**(6):752–757
24. Petersen OH, Gerasimenko OV, Gerasimenko JV. Endocytic uptake of SARS-CoV-2: the critical roles of pH, Ca²⁺, and NAADP. *FUNCTION* 2020;**1**(1):zqaa003
25. Bruen C, Miller J, Wilburn J, et al. Auxora for the Treatment of Patients With Acute Pancreatitis and Accompanying Systemic Inflammatory Response Syndrome. *Pancreas* 2021;**50**(4):537–543