

High-throughput SNP-genotyping analysis of the relationships among Ponto-Caspian sturgeon species

Sergey M. Rastorguev^{1,2}, Artem V. Nedoluzhko¹, Alexander M. Mazur³, Natalia M. Gruzdeva¹, Alexander A. Volkov², Anna E. Barmintseva², Nikolai S. Mugue² & Egor B. Prokhortchouk^{1,3}

¹National Research Center "Kurchatov Institute", Kurchatov sq. 1, 123182 Moscow, Russia

²Russian Federal Research Institute of Fisheries and Oceanography (VNIRO), V. Krasnoselskaya str. 17, 107140 Moscow, Russia

³Center of Bioengineering, Russian Academy of Sciences, 60-letiya Oktyabrya av. 7-1, 117312 Moscow, Russia

Keywords

Genotyping, glaciation, population genetics, SNP, species relationships, sturgeon.

Correspondence

Sergey M. Rastorguev, National Research Center "Kurchatov Institute," Laboratory of Genome Analysis, NBIC Center, Kurchatov sq. 1, 123182 Moscow, Russia. Tel: +7-(499)-1967937; Fax: +7-(499)-1967723; E-mail: rastorgueff@gmail.com

Funding Information

No funding information provided.

Received: 28 March 2013; Revised: 30 May 2013; Accepted: 31 May 2013

Ecology and Evolution 2013; 3(8): 2612–2618

doi: 10.1002/ece3.659

Introduction

The Russian sturgeon (*Acipenser gueldenstaedtii*) is one of the most commercially valuable fish species in the Caspian, Azov, and Black Seas due to its role as a major caviar producer (Vlasenko et al. 1989a,b; Birstein and Bemis 1997) and important aquaculture species (Fig. 1). Three other species are closely related to *A. gueldenstaedtii*: the Persian sturgeon (*A. persicus*), the Adriatic sturgeon (*A. naccarii*), and the Siberian sturgeon (*A. baerii*), the least abundant and hence most valuable species. Together, these four species make up the Ponto-Caspian clade of sturgeons. However, several difficulties have complicated efforts to distinguish between the various species in this clade. *Acipenser persicus* inhabits the Caspian Sea and cannot be readily distinguished from *A. gueldenstaedtii* by mitochondrial markers (Birstein et al. 2000; Birstein and Doukakis 2001), or by morphological or anatomical differences (Vlasenko et al. 1989a,b; Ruban

Abstract

Legally certified sturgeon fisheries require population protection and conservation methods, including DNA tests to identify the source of valuable sturgeon roe. However, the available genetic data are insufficient to distinguish between different sturgeon populations, and are even unable to distinguish between some species. We performed high-throughput single-nucleotide polymorphism (SNP)-genotyping analysis on different populations of Russian (*Acipenser gueldenstaedtii*), Persian (*A. persicus*), and Siberian (*A. baerii*) sturgeon species from the Caspian Sea region (Volga and Ural Rivers), the Azov Sea, and two Siberian rivers. We found that Russian sturgeons from the Volga and Ural Rivers were essentially indistinguishable, but they differed from Russian sturgeons in the Azov Sea, and from Persian and Siberian sturgeons. We identified eight SNPs that were sufficient to distinguish these sturgeon populations with 80% confidence, and allowed the development of markers to distinguish sturgeon species. Finally, on the basis of our SNP data, we propose that the *A. baerii*-like mitochondrial DNA found in some Russian sturgeons from the Caspian Sea arose via an introgression event during the Pleistocene glaciation.

et al. 2011). Thus, the taxonomic rank of *A. persicus* is disputed, with some authors considering it to be a separate species (Luk'yaneko et al. 1974; Putilina 1983) and others regarding it to be a subspecies of *A. gueldenstaedtii* (Berg 1934; Birstein and Bemis 1997; Birstein et al. 2000; Birstein and Doukakis 2001; Ruban et al. 2011). Although the Adriatic sturgeon uniquely inhabits the northern Adriatic Sea, it also morphologically resembles *A. gueldenstaedtii* (Tortonese 1989). Unlike its anadromous relatives, the Siberian sturgeon is a freshwater species. Its geographical distribution includes Siberian Rivers and Lake Baikal, and does not overlap with Russian sturgeon. All these species are highly endangered, and are under the control and protection of the Convention for International Trade in Endangered Species (CITES).

The unusual polyphyly of Russian sturgeons' mitochondrial DNA further complicates their unique identification. Roughly a third of Russian sturgeon specimens from the Caspian Sea have mitochondrial DNA that is similar to




Figure 1. The Russian sturgeon (*Acipenser gueldenstaedtii*) at the Astrakhan fish farm.

that of Siberian sturgeons, which is often referred to as the “*baerii*-like mitotype” (Birstein et al. 2000; Jenneckens et al. 2000; Rastorguev et al. 2008). However, *baerii*-like haplotypes found in the Caspian population of *A. gueldenstaedtii* are different from those of *A. baerii* in one substitution in the highly variable D-loop and can be distinguished by PCR-based test (Mugue et al. 2008). Two percent of Russian sturgeons in the Azov Sea have the *baerii*-like mitotype (Timoshkina et al. 2009). In contrast, no Persian sturgeons have been found to possess the *baerii*-like mitotype (Mugue et al. 2008), supporting that *A. persicus* is genetically isolated entity. Nevertheless, mitochondrial DNA analysis can distinguish only the Siberian sturgeon from among its relatives (Mugue et al. 2008).

There are several explanations of the “*baerii*-like mitotype” phenomenon. Some authors propose that the *baerii*-like mitotype arose from farmed *A. baerii* having escaped into the Volga River (Jenneckens et al. 2000), whereas others suggest that the *baerii*-like mitotype and the majority *A. gueldenstaedtii* mitotype are ancestral mitotypes for the Siberian, Russian, and Persian sturgeon species (Birstein et al. 2005). It is also possible that hybridization between Siberian and Russian sturgeons during the Pleistocene glaciation led to Siberian sturgeon mitochondrial DNA introgression, as considerable changes in the species distributions of other organisms occurred in that period (Hewitt 2000).

The identification of nuclear genome markers associated with specific Ponto-Caspian sturgeon species has generally lagged behind due to the polyploidy of the sturgeon genome. It is widely accepted that polyploidization took place several times within the order Acipenseriformes. First whole-genome duplication presumably occurred in the ancestral lineage and led to formation of $2n = 120\text{--}140$ (“low chromosome number”) group of species. Two independent second rounds of polyploidization were proposed: one in Pacific clade, and another one in Atlantic clade of sturgeons ($2n = 240\text{--}260$), leading to formation of two phylogenetically unrelated groups of “high chromosome

number” sturgeon species (Birstein et al. 1993). Evidence suggests that low chromosome number species are tetraploids with functionally diploid genome, and high chromosome number species are octoploids, with two sets of paralogous tetrasomic loci.

The species relationship between Russian and Persian sturgeons has been addressed recently by restriction site-associated DNA (RAD) sequencing (Baird et al. 2008), but the issue remains unresolved (Ogden et al. 2013).

Meanwhile, high-throughput single-nucleotide polymorphism (SNP) genotyping provides precise sample assignment to its origin therefore it becomes highly valuable tool against illegal fishing and mislabeling. Effectiveness of such approach was shown for four commercial marine fish species (Nielsen et al. 2012).

In this study, we performed high-throughput genotyping analysis of individuals from several populations of Russian, as well as Persian, and Siberian sturgeons to find SNP markers for species identification. The resulting data should aid in sturgeon conservation efforts and help prevent illegal fishing. Additionally, we suggest an explanation of the *baerii*-like mitotype found in some Russian sturgeons in the Caspian Sea region.

Material and Methods

RNA isolation and cDNA library construction

A several-month-old Russian sturgeon fingerling from an Astrakhan fish farm (Ltd Astrakhan Fish Breeding Company, also known as “Beluga”) was suspended in RNAlater reagent (Invitrogen/Life Technologies, Carlsbad, CA) for the preparation of a cDNA library. RNA extraction from different tissues (representing total body RNA content) was done with the TRIzol reagent according to the provided RNA extraction protocol (Invitrogen). Recent international FP6 supported project SturSNiP revealed that there is no fixed segregating genetic markers between these species, and all informative markers are frequency dependent due to shared polymorphism (Ogden et al. 2013). Our approach was different from that one of SturSNiP because our aim was to reveal highly polymorphic loci in the Caspian population of Russian sturgeon; therefore, we collected transcriptome information from a single individual rather than from pooled samples from different populations of different species. There was twofold rationale behind taking one fingerling for transcriptome analysis; first, we attempted to analyze the entire mRNA repertoire, so we needed artificially propagated fingerling to be sacrificed and the diversity of tissues to be sampled, and second, the data of allelic composition from one individual allowed us to avoid problem of paralogous (resulted from previous whole-genome duplication) loci and to discrimi-

nate paralogous loci from true polymorphic sites (see below). cDNA was prepared from total RNA by a M-MLV reverse transcriptase approach (Schmidt and Mueller 1999) using the MINT cDNA synthesis kit (Evrogen, catalog #SK001). The resulting “SMART” cDNA was amplified and normalized (reducing the levels of highly abundant transcripts), using the duplex-specific nuclease (DSN) normalization method (Zhulidov et al. 2004).

Transcriptome sequencing and mapping

Approximately 5 μg of nonnormalized cDNA and 10 μg of normalized cDNA were used for SOLiD library construction and sequencing using the SOLiD™ 4.0 system (Life Technologies) according to the standard protocol (SOLiD User Guide). Reads from the normalized and nonnormalized libraries were mapped against the sequence of Russian sturgeon mitochondrial DNA (NCBI accession number NC_012576.1), and all reads apparently derived from mitochondrial DNA were excluded from further mapping. The remaining reads were mapped against the sturgeon EST database from GenBank using the Bowtie software package (Langmead et al. 2009). An EST-derived reference database was created using sequences bearing the “Acipenser” tag in the “species name” field (6219 sequences in total), and was then used for mapping. SNP calling was performed using the SAMtools package (Li et al. 2009; Li 2011). Mismatches between mapped reads were catalogued in VCF files, and heterozygous positions were then identified with a custom Perl script. SNP positions were sorted by coverage, and those that were covered by more than 100 reads were selected for the further analysis.

Sturgeon samples and genotyping

Sturgeon samples were obtained from The Russian National Collection of Reference Genetic Materials (RNCGRM), developed by the Russian CITES scientific authority on sturgeon fishes and maintained by VNIRO. Population samples of Caspian basin Russian sturgeons originated from the Volga and Ural Rivers (14 specimens each), with five additional specimens of Russian sturgeons originating from the Azov Sea; 28 Persian sturgeon specimens originated from the northern Caspian Sea; and five Siberian sturgeon specimens originated from the Lena and Irtysh rivers. DNA was isolated using the Wizard® SV 96 Genomic DNA Purification System (Promega, Fitchburg, WI).

Custom GoldenGate Genotyping Panel was constructed in Illumina Inc., San Diego, CA using loci, selected from SNP calling procedure described above (Table S1). The GoldenGate assay was performed using the Illumina GoldenGate platform according to the manufacturer’s

instructions and analyzed by the genotyping module of Illumina’s Genome Studio data analysis software (Illumina Inc.). Loci for which all specimens showed the same genotype or genotypes could not be distinguished by this software, and were excluded from further analysis.

Genotyping data analysis

An UPGMA (unweighted pair group method with arithmetic mean) tree of sturgeon populations was constructed using the POPTREE2 software (Takezaki et al. 2010), with F_{ST} distances according to Latter (1972). We used the Whichloci program to identify SNP loci with the highest statistical power for population assignments (Banks et al. 2003). For the Whichloci analysis, we generated four simulated datasets by random resampling the SNP loci (three with sample size 100, one with sample size 500), and we then ran Whichloci on each dataset and chose loci that were present in the outcomes from all four simulated datasets. Population structure was inferred using the STRUCTURE software (Pritchard et al. 2000), with parameters “prior sample location information” and “allele frequency model correlated among populations” included in the admixture model. F_{ST} statistics – the effect of subpopulations (s) compared to the total population (t) – were determined using the GenePop web service (<http://genepop.curtin.edu.au/>; Raymond and Rousset 1995; Rousset 2008), with the “ F_{ST} and other correlation” options.

Results and Discussion

RNA-Seq analysis of the sturgeon transcriptome

Transcriptome data for the Russian sturgeon fingerling were generated from approximately 250 million 50-base reads of a normalized cDNA library and 80 million reads of a nonnormalized cDNA library using the SOLiD (Life Technology) platform. After removing mitochondrial DNA reads, approximately 1.7 million reads from the normalized library and 3 million reads from the nonnormalized library mapped to the GenBank sturgeon expressed sequence tag (EST) database (Table 1; note that the reduced number of mapped reads for the normalized library was expected due to the removal of ribosomal and tRNA transcripts during the normalization process; 0.3% [821,892 of 252,186,716] of normalized library reads and 4.6% [3,624,935 of 79,030,219] of control library reads were mapped on ribosomal database). From 6219 totals, 4300 ESTs were covered by at least one read. The overall low numbers of mapped reads could be explained by the fact that the database contains mainly *A. sinensis* and *A. transmontanus* ESTs, which are distantly related to *A. gueldenstaedtii*.

SNP discovery for sturgeon genotyping and population identification

Several thousand SNP candidates were identified by screening heterozygous positions using the following criteria of read sequence quality, read mapping quality, and quality score at the mismatch position. Additionally, we required that the presumptive SNPs have allele ratios

Table 1. SOLiD read mapping and SNP calling statistics for two *Acipenser gueldenstaedtii* cDNA libraries.

	Normalized cDNA library	Nonnormalized cDNA library
Total reads	252,186,716	79,030,219
Mapped reads	1,787,162	3,159,679
Total SNPs called	1,014,088	755,147
Heterozygous SNP	50,124	30,682
SNPs (coverage ≥ 10)	611,971	415,411
Heterozygous (coverage ≥ 10)	45,645	27,143
SNPs (coverage ≥ 100)	181,683	138,058
Heterozygous (coverage ≥ 100)	17,765	9812

Read mapping was performed using the sturgeon EST database as a reference. SNP calling was done with default SNP caller settings (using SAMtools) and with minimum coverage depths of 10 and 100.

close to 3:1 based on the following assumptions: “high chromosome number” sturgeons, including three species in our study, have two sets of paralogous tetrasomic loci (total eight similar genomic regions, segregated as two tetrasomic loci), special precaution was applied to avoid erroneous considering alternatively fixed paralogous loci as a polymorphic locus. Thus, alleles with 1:1 ratios would likely represent divergent paralogous loci which would not be polymorphic across individuals in a population. Therefore, we sought to identify genotypes where one tetrasomic set of alleles was heterozygous and the other locus (paralog) was homozygous. Allelic ratio 3:1 would likely represent situation when one paralog is fixed (i.e., AAAA) and the other one is polymorphic (AABB). Total 384 of the best-scoring mismatch positions were chosen for a GoldenGate custom SNP panel (Illumina) for sturgeon genotyping (Table S1). The GoldenGate assay revealed that 123 of the 384 variable loci were successfully genotyped; the others appeared homozygous or heterozygous across all individuals (fixed at the same or alternative alleles at paralogous loci) and were excluded from further analysis. Loci that appeared heterozygous across all individuals most likely reflected the presence of a paralogous locus, creating the appearance of two different alleles.

Table 2. DNA sequences of the 12 SNP candidate markers used for sturgeon population assignment.

Marker name	DNA sequence
EV824350.93	TTCGATACGATAAGCCGCAATGCTGTCAGGAACACTGGCTAAACCCGCAATGCGCGGTCT[A/G] CTCGGTAACGCTACGATTCCACCTGGCTGTTGCTTTCACGCTGCTTTGGCTGCAGCA
ES698489.358	ATTTTAAAGTGGARCCCTTTGTTGTTGATTCTGTTTAAATTCCAGCACCTTCACAAA[AV]TC CAGTAAGAAAGACATGTGAAGTAGACMSTGGCCGTGATTTTCAGATGTGCTCTGATAA
DR977078.72	TTGCCAGGCTTTGAACAATATGCTATCAAGAAGTTTGCAGAGGCTTTTGAAGCCATTCC[A/G]C GTGCTCTTGAGAGAACTCTGGTGCAAGGGCAATGAACCTATCTCCAAACTGTATGCT
DR976209.314	GAACAGTGCCTCCACAAGAATTGTTTTAGCCGTATAACTCAAGAACAGAAAATAAG[T/C] TGCAATTAGAGTGGAGGAAGACAGATGACTGCGCTACTGTACACCCCTTTTACAACAASA
DR977006.239	CTCTTTCACCTTTGAGCTTCGTGATACTGGCAGATATGGATTCTCTTCCAGAGTGC[A/G]A TCAGGCCAAGTTCGAAGAACAATGCTGGCAGTCAATACATTGCCAAACATGTTTCAG
EV824380.474	CCAAGGGCAATGTACCGCAACTCTGGAGCCATCCATACGAGTTCAAGGCGTAAAGCA[T/C] GGTCAAGCCAGAATAAAGTCTGTTTTCAACAAAAAAAAAAAAAAAAAAAAAAAAANNA
DR976578.395	TTTCCCATCAATGTGGTCGTTTCAGGAACTGGGTCCTGGTGGAAATCAGGAATTTCTT[A/G]G GAGAAAAGTACATCCGCAGAGTACGCATGAGGCAGGGTGTGACCTGTAATGTTTCCAG
DR975536.188	TAGAAGAGGGCGCTGACCAAGCACACCAAGTGTTCGGGACATGATCCGCGAGGTGTG[T/C] GGCTTCGCCCTACGAGAGACGCGCCATGGAGCTGCTGAAGGCTCCAAGGATAAGCGM
DR977561.396	GAAGTTTGCCTGCAATGGGACTGTGATTGAGCATCCAGAATATGGTGAAGTGGTTCAGCT[A/G] CAGGGTGATCAGCGCAAGAATATCTGCCAGTTCCTACTGAGATTGGCTGGGTAAGGAG
ES698197.350	AACAAGCCAAGGACAAGCCAGGAGAAACCAAATAACATGGCAAGAGRACTAACCATC[A/G] CTTACAAGAGAAGTTTCAATCTCCGACACAAGCTTCTGTCTGGGATTTTCATTTTCMTT
DR974886.648	TTGGTACCGTGTATCTCTCTGCTCTGGCTTTATAATGATGGGTGTACCGTATATGAAGG[A/G]G GCACCTGTAATGCASATGGTAAACCAATCAGTAATGTAAGAATTGGATGCAATAAAG
ES697699.316	GAGAAGATYGACCTGAAGTTCAACCACCTCAAGTTCGGACACGGACGCTTCCAGACCGC[T/C] GAAGAGAAGAAGCGTTCATGGGACCACTCAAAAAGGACCGAATCTCAAGGAGGAGACT

Each polymorphic position is shown in square brackets. Sequence names consist of the GenBank accession number of the candidate EST sequence, with the SNP position within the EST given after the period.

Analysis of sturgeon population structure

To perform a population structure analysis, genotype calls were filtered from the Illumina Genome Studio report into the STRUCTURE (Pritchard et al. 2000) and GenePop (Raymond and Rousset 1995) programs. The Which-loci program (Banks et al. 2003) guided the choices of the most informative loci for population assignment. We detected few genetic differences at the 123 diagnostic positions between the Volga River and Ural River populations of Russian sturgeons. Eight of the most informative SNP loci were sufficient to assign specimens to the Caspian Russian (Volga and Ural River) populations, Russian Azov Sea population, and Persian populations with 80% accuracy. A total of 12 loci were sufficient to make these population assignments with 90% accuracy (Table 2).

Using the allele frequencies within each population for all 123 polymorphic SNP loci (taken from the Genome Studio report), we constructed a population tree (Fig. 2). As expected, the Siberian sturgeons appeared more distant from the other species of the Ponto-Caspian clade. Our data also indicated that the Persian sturgeon was separated from all three samples of the Russian sturgeon, suggesting that *A. persicus* is genetically different from all *A. gueldenstaedtii* populations. These data do not support an opinion of Ruban et al. (2011) that *A. persicus* is an ecological morph within the Caspian Sea population of Russian sturgeon, but not a valid species. The population assignment analysis (STRUCTURE) also supported the genetic separation of the Persian sturgeon, as a homogeneous cluster consisted exclu-

sively of *A. persicus* specimens (Fig. 3). The Siberian and Persian populations were clearly separated into two distinct groups, whereas the Russian sturgeons appeared as intermediate and shared population structure elements of both the Siberian and Persian sturgeons. This is likely due to nature of our SNP loci when polymorphic loci were obtained from Russian sturgeon specimen.




Figure 2. UPGMA tree showing the relationships of sturgeon populations (POPTREE2; Takezaki et al. 2010). Genetic distances (F_{ST}) between the populations (Latter 1972) were estimated using allele frequency data for 123 loci. Bootstrap values are indicated next to corresponding nodes. GUE_azov, *Acipenser gueldenstaedtii* from the Azov Sea; GUE_ur, *A. gueldenstaedtii* from the Ural River; GUE_vol, *A. gueldenstaedtii* from the Volga River; PERS, *A. persicus*; BAE, *A. baerii*. (The Ural and Volga River sturgeons are considered to be representative of Caspian Sea Russian sturgeons.)




Figure 3. Population structure analysis of various sturgeon samples using the STRUCTURE software. Three runs of STRUCTURE with the cluster number (K) set to two, three, or four are shown. Siberian (*Acipenser baerii*) and Persian (*A. persicus*) sturgeons form their own distinct clusters, whereas all three Russian sturgeon populations have mixed origins.

Siberian sturgeon introgression into Caspian Russian sturgeon population during the Pleistocene glaciation

Frequency of several SNPs of the Caspian (Volga and Ural Rivers) populations of Russian sturgeon shows an intermediate value compared to those in Siberian and *Persicus* populations, indicating that Caspian population of the Russian sturgeon retains traces of ancient introgression of Siberian sturgeon into ancestral Caspian population of the Russian sturgeon during the Pleistocene glaciation. These data support the hypothesis that the *baerii*-like mitotype is a result of this ancient introgression. The climatic oscillations during the glaciation caused great changes in species distribution, and considerable evidence suggests that other organisms dispersed to new locations (Hewitt 2000). An investigation of 41 North American fish species concluded that the Pleistocene glaciation had a great impact on fish population structure (Bernatchez and Wilson 1998). The presence of distinct mitochondrial DNA sequences within a species has been explained as a consequence of intraspecific lineage reorganization, whereby the dispersed lineages of an ancestral set of refugees reunite in one formerly glaciated region (Taberlet and Bouvet 1994; Santucci et al. 1998).

Notably, the hypothesis of the interspecific hybridization is required to explain abundance of the *baerii*-like mitotype and nuclear markers in the Caspian population of *A. gueldenstaedtii*. This scenario is supported by evidence that hybridization between Russian and Siberian sturgeons leads to the formation of fertile hybrids (Jennekens et al. 2000), which are widely used in aquaculture. Considering that the glaciation allowed the Siberian sturgeons into the Caspian Sea, there may have been nothing to prevent their hybridization with Caspian sturgeons. Previously published data that “*baerii*-like” mitotypes are present in the Caspian Russian sturgeon population and absent from the Persian population are consistent with this notion (Mugue et al. 2008).

Conclusions

In this study, we discovered SNP markers for distinguishing between Caspian and Azov Sea populations of Russian sturgeons, as well as between Russian, Persian, and Siberian sturgeons. These data allow the development of a genetic test that can be used for law enforcement in sturgeon fishery, aquaculture industry, and trading, as well as in conservation efforts for endangered sturgeon species. Finally, our SNP analysis supported the idea that the mitochondrial DNA polyphyly of Russian sturgeons in the Caspian Sea resulted from an introgression event during the Pleistocene glaciation.

Acknowledgments

We thank Anthon Teslyuk for useful discussions and for technical support with the computer cluster of the NRC Kurchatov Institute, Maria Tolochkova for help in obtaining samples from the RNCSSGM genetic material collection, and Aleksey Sergeev and Nikolay Chekanov for comments and discussion.

Data Accessibility

The SOLiD sequence data in .csfasta + .quality formats, mapped data in SAM/BAM formats, and SNP variants in .vcf format are available on request.

Conflict of Interest

None declared.

References

- Baird, N. A., P. D. Etter, T. S. Atwood, M. C. Currey, A. L. Shiver, Z. A. Lewis, et al. 2008. Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLoS ONE* 3:e3376.
- Banks, M. A., W. Eichert, and J. B. Olsen. 2003. Which genetic loci have greater population assignment power? *Bioinformatics* 19:1436–1438.
- Berg, L. S. 1934. *Acipenser gueldenstadti persicus*, a sturgeon from the south Caspian Sea. *Ann. Mag. Hist. London Ser.* 10 13:317–318.
- Bernatchez, L., and C. C. Wilson. 1998. Comparative phylogeography of nearctic and palearctic fishes. *Mol. Ecol.* 7:431–452.
- Birstein, V. J., and W. E. Bemis. 1997. How many species are there within the genus *Acipenser*? Pp. 157–163 in V. J. Birstein, J. R. Waldman, and W. E. Bemis, eds. *Sturgeon biodiversity and conservation*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Birstein, V. J., and P. Doukakis. 2001. Molecular identification of sturgeon species: science, bureaucracy, and the impact of environmental agreements. Pp. 47–63 in K. Riede, ed. *New perspectives for monitoring for migratory animals*. International Workshop on Behalf of the 20th Anniversary of the Bonn Convention. Federal Agency for Nature Conservation, Bonn, Germany.
- Birstein, V. J., A. I. Poletaev, and B. F. Goncharov. 1993. DNA content in Eurasian sturgeon species determined by flow cytometry. *Cytometry* 14:377–383.
- Birstein, V. J., P. Doukakis, and R. DeSalle. 2000. Polyphyly of mtDNA lineages in the Russian sturgeon, *Acipenser gueldenstaedtii*: forensic; evolutionary implications. *Conserv. Genet.* 1:81–88.

- Birstein, V. J., G. Ruban, A. Ludwig, P. Doukakis, and R. DeSalle. 2005. The enigmatic Caspian Sea Russian sturgeon: how many cryptic forms does it contain? *Syst. Biodivers.* 3:203–218.
- Hewitt, G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405:907–913.
- Jenneckens, I., J. N. Meyer, L. Debus, C. Pitra, and A. Ludwig. 2000. Evidence of mitochondrial DNA clones of Siberian sturgeon, *Acipenser baerii*, within Russian sturgeon, *Acipenser gueldenstaedtii*, caught in the River Volga. *Ecol. Lett.* 3:503–508.
- Langmead, B., C. Trapnell, M. Pop, and S. L. Salzberg. 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10:R25.
- Latter, B. D. 1972. Selection in finite populations with multiple alleles. 3. Genetic divergence with centripetal selection and mutation. *Genetics* 70:475–490.
- Li, H. 2011. Improving SNP discovery by base alignment quality. *Bioinformatics* 27:1157–1158.
- Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, et al. 2009. 1000 Genome Project Data Processing Subgroup. The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078–2079.
- Luk'yanenko, V. I., Z. G. Umerov, and B. B. Karataeva. 1974. The Southern sturgeon is a separate species within the genus *Acipenser*. *Izv. Akad. Nauk SSSR Ser. Biol.* 5:736–739. (in Russian)
- Mugue, N. S., A. E. Barmintseva, S. M. Rastorguev, V. N. Mugue, and V. A. Barmintsev. 2008. Polymorphism of the mitochondrial DNA control region in eight sturgeon species and development of a system for DNA-based species identification. *Rus. J. Genetics* 44:793–798.
- Nielsen, E. E., A. Cariani, E. Mac Aoidh, G. E. Maes, I. Milano, R. Ogden, et al. 2012. Gene-associated markers provide tools for tackling illegal fishing and false eco-certification. *Nat. Commun.* 3:851.
- Ogden, R., K. Gharbi, N. Mugue, J. Martinsohn, H. Senn, J. W. Davey, et al. 2013. Sturgeon conservation genomics: SNP discovery and validation using RAD sequencing. *Mol. Ecol.* 22:3112–3123. doi:10.1111/mec.12234.
- Pritchard, J. K., M. Stephens, and P. Donnelly. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155:945–959.
- Putilina, L. A. 1983. Morphological characteristics of the Persian sturgeon in the Volga River. Pp. 70–71 in V. N. Belyaeva, ed. Complex utilization of biological resources of the Caspian and Azov Seas. VNIRO, Moscow, Russia. (in Russian)
- Rastorguev, S., N. Mugue, A. Volkov, and V. Barmintsev. 2008. Complete mitochondrial DNA sequence analysis of Ponto-Caspian sturgeon species. *J. Appl. Ichthyol.* 24:46–49.
- Raymond, M., and F. Rousset. 1995. GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *J. Hered.* 86:248–249.
- Rousset, F. 2008. genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Mol. Ecol. Resour.* 8:103–106.
- Ruban, G. I., M. V. Kholodova, V. A. Kalmykov, and P. A. Sorokin. 2011. A review of the taxonomic status of the Persian sturgeon (*Acipenser persicus* Borodin). *J. Appl. Ichthyol.* 27:470–476.
- Santucci, F., B. Emerson, and G. M. Hewitt. 1998. Mitochondrial DNA phylogeography of European hedgehogs. *Mol. Ecol.* 7:1163–1172.
- Schmidt, W. M., and M. W. Mueller. 1999. CapSelect: a highly sensitive method for 5' CAP dependent enrichment of full-length cDNA in PCR mediated analysis of mRNAs. *Nucleic Acids Res.* 27:e31.
- Taberlet, P., and J. Bouvet. 1994. Mitochondrial DNA polymorphism, phylogeography, and conservation genetics of the brown bear (*Ursus arctos*) in Europe. *Proc. Biol. Sci.* 255:195–200.
- Takezaki, N., M. Nei, and K. Tamura. 2010. POPTREE2: software for constructing population trees from allele frequency data and computing other population statistics with Windows interface. *Mol. Biol. Evol.* 27:747–752.
- Timoshkina, N. N., A. E. Barmintseva, A. V. Usatov, and N. S. Miuge. 2009. Intraspecific genetic polymorphism of Russian sturgeon *Acipenser gueldenstaedtii*. *Genetika* 45:1250–1259. (in Russian)
- Tortonese, E. 1989. *Acipenser naccarii* Bonaparte, 1836. Pp. 284–293 in J. Holcik, ed. The freshwater fishes of Europe. Vol. 1, Part II. General introduction to fishes, Acipenseriformes. AULA-Verlag, Wiesbaden, Germany.
- Vlasenko, A. D., A. V. Pavlov, L. I. Sokolov, and V. P. Vasilev. 1989a. *Acipenser gueldenstaedti* Brandt, 1833. Pp. 294–344 in J. Holcik, ed. The freshwater fishes of Europe. General introduction to fishes, Acipenseriformes. 1 II. AULA-Verlag, Wiesbaden, Germany.
- Vlasenko, A. D., A. V. Pavlov, and V. P. Vasil'ev. 1989b. *Acipenser persicus* Borodin, 1897. Pp. 344–366 in J. Holcik, ed. The freshwater fishes of Europe. Vol. 1, Part II. General introduction to fishes. Acipenseriformes. AULA-Verlag, Wiesbaden, Germany.
- Zhulidov, P. A., E. A. Bogdanova, A. S. Shcheglov, L. L. Vagner, G. L. Khaspekov, V. B. Kozhemyako, et al. 2004. Simple cDNA normalization using kamchatka crab duplex-specific nuclease. *Nucleic Acids Res.* 32:e37.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. List of loci for GoldenGate.